

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
21 March 2002 (21.03.2002)

PCT

(10) International Publication Number  
**WO 02/21996 A2**

(51) International Patent Classification<sup>7</sup>:

**A61B**

(74) Agents: **BASTIAN, Kevin, L.** et al.; Townsend and Townsend and Crew LLP, Two Embarcadero Center, Eighth Floor, San Francisco, CA 94111 (US).

(21) International Application Number: PCT/US01/28716

(81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(22) International Filing Date:

14 September 2001 (14.09.2001)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

09/663,733 15 September 2000 (15.09.2000) US  
09/930,020 14 August 2001 (14.08.2001) US

(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(71) Applicant (for all designated States except US): **EOS BIOTECHNOLOGY, INC.** [US/US]; 225 A Gateway Boulevard, South San Francisco, CA 94080 (US).

**Published:**

— without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.



**WO 02/21996 A2**

(54) Title: METHODS OF DIAGNOSIS OF COLORECTAL CANCER, COMPOSITIONS AND METHODS OF SCREENING FOR COLORECTAL CANCER MODULATORS

(57) Abstract: Described herein are methods that can be used for diagnosis and prognosis of colorectal cancer. Also described herein are methods that can be used to screen candidate bioactive agents for the ability to modulate colorectal cancer. Additionally, methods and molecular targets (genes and their products) for therapeutic intervention in colorectal and other cancers are described.

## Methods of Diagnosis of Colorectal Cancer, Compositions and Methods of Screening for Colorectal Cancer Modulators

### 5 CROSS-REFERENCES TO RELATED APPLICATIONS

[01] This application is a continuation in part of US Patent Application USSN 09/663,733 filed September 15, 2000, and US Patent Application filed August 14, 2001 USSN, not yet known, which are incorporated herein by reference in their entirety.

### 10 FIELD OF THE INVENTION

[02] The invention relates to the identification of expression profiles and the nucleic acids involved in colorectal cancer, and to the use of such expression profiles and nucleic acids in diagnosis and prognosis of colorectal cancer. The invention further relates to methods for identifying and using candidate agents and/or targets which modulate colorectal 15 cancer.

### BACKGROUND OF THE INVENTION

[03] Cancer of the colon and/or rectum (referred to as "colorectal cancer") are significant in Western populations and particularly in the United States. Cancers of the colon and rectum occur in both men and women most commonly after the age of 50. These 20 develop as the result of a pathologic transformation of normal colon epithelium to an invasive cancer. There have been a number of recently characterized genetic alterations that have been implicated in colorectal cancer, including mutations in two classes of genes, tumor-suppressor genes and proto-oncogenes, with recent work suggesting that mutations in DNA repair genes may also be involved in tumorigenesis. For example, inactivating mutations of 25 both alleles of the adenomatous polyposis coli (APC) gene, a tumor suppressor gene, appears to be one of the earliest events in colorectal cancer, and may even be the initiating event. Other genes implicated in colorectal cancer include the MCC gene, the p53 gene, the DCC (deleted in colorectal carcinoma) gene and other chromosome 18q genes, and genes in the TGF- $\beta$  signaling pathway. For a review, see *Molecular Biology of Colorectal Cancer*, pp. 30 238-299, in *Curr. Probl. Cancer*, Sept/Oct 1997; see also Williams, *Colorectal Cancer* (1996); Kinsella & Schofield, *Colorectal Cancer: A Scientific Perspective* (1993); *Colorectal*

*Cancer: Molecular Mechanisms, Premalignant State and its Prevention* (Schmiegel & Scholmerich eds., 2000); *Colorectal Cancer: New Aspects of Molecular Biology and Their Clinical Applications* (Hanski et al., eds 2000); McArdle et al., *Colorectal Cancer* (2000); Wanebo, *Colorectal Cancer* (1993); Levin, *The American Cancer Society: Colorectal Cancer* (1999); *Treatment of Hepatic Metastases of Colorectal Cancer* (Nordlinger & Jaeck eds., 1993); *Management of Colorectal Cancer* (Dunitz et al., eds. 1998); *Cancer: Principles and Practice of Oncology* (Devita et al., eds. 2001); *Surgical Oncology: Contemporary Principles and Practice* (Kirby et al., eds. 2001); Offit, *Clinical Cancer Genetics: Risk Counseling and Management* (1997); *Radioimmunotherapy of Cancer* (Abrams & Fritzberg eds. 2000); Fleming, *AJCC Cancer Staging Handbook* (1998); *Textbook of Radiation Oncology* (Leibel & Phillips eds. 2000); and *Clinical Oncology* (Abeloff et al., eds. 2000).

[04] Imaging of colorectal cancer for diagnosis has been problematic and limited. In addition, metastasis of the tumor to the lumen, and metastasis of tumor cells to regional lymph nodes are important prognostic factors (see, e.g., *PET in Oncology: Basics and Clinical Application* (Ruhlmann et al. eds. 1999). For example, five year survival rates drop from 80 percent in patients with no lymph node metastases to 45 to 50 percent in those patients who do have lymph node metastases. A recent report showed that micrometastases can be detected from lymph nodes using reverse transcriptase-PCR methods based on the presence of mRNA for carcinoembryonic antigen, which has previously been shown to be present in the vast majority of colorectal cancers but not in normal tissues. Liefers et al., *New England J. of Med.* 339(4):223 (1998).

[05] Thus, methods that can be used for diagnosis and prognosis of colorectal cancer would be desirable. Accordingly, provided herein are methods that can be used in diagnosis and prognosis of colorectal cancer. Further provided are methods that can be used to screen candidate bioactive agents for the ability to modulate colorectal cancer. Additionally, provided herein are molecular targets for therapeutic intervention in colorectal and other cancers.

#### BRIEF SUMMARY OF THE INVENTION

[06] The present invention provides novel methods for diagnosis and prognosis evaluation for colorectal cancer, as well as methods for screening for compositions which modulate colorectal cancer. Methods of treatment of colorectal cancer, as well as compositions, are also provided herein.

[07] In one aspect, a method of screening drug candidates comprises providing a cell that expresses an expression profile gene selected from those of Table I. The method further includes adding a drug candidate to the cell and determining the effect of the drug candidate on the expression of the expression profile gene.

5 [08] In one embodiment, the method of screening drug candidates includes comparing the level of expression in the absence of the drug candidate to the level of expression in the presence of the drug candidate, wherein the concentration of the drug candidate can vary when present, and wherein the comparison can occur after addition or removal of the drug candidate. In a preferred embodiment, the cell expresses at least two expression profile genes. The profile genes may show an increase or decrease.

10 [09] Also provided herein is a method of screening for a bioactive agent capable of binding to a colorectal cancer modulator protein, the method comprising combining the colorectal cancer modulator protein and a candidate bioactive agent, and determining the binding of the candidate agent to the colorectal cancer modulator protein.

15 Preferably the colorectal cancer modulator protein is a product encoded by a gene of Table 1 or Table 2.

20 [10] Further provided herein is a method for screening for a bioactive agent capable of modulating the activity of a colorectal cancer modulator protein. In one embodiment, the method comprises combining the colorectal cancer modulator protein and a candidate bioactive agent, and determining the effect of the candidate agent on the bioactivity of the colorectal cancer modulator protein. Preferably the colorectal cancer modulator protein is a product encoded by a gene of Table 1 or Table 2.

25 [11] Also provided is a method of evaluating the effect of a candidate colorectal cancer drug comprising administering the drug to a transgenic animal expressing or over-expressing the colorectal cancer modulator protein, or an animal lacking the colorectal cancer modulator protein, for example as a result of a gene knockout.

30 [12] Additionally, provided herein is a method of evaluating the effect of a candidate colorectal cancer drug comprising administering the drug to a patient and removing a cell sample from the patient. The expression profile of the cell is then determined. This method may further comprise comparing the expression profile to an expression profile of a healthy individual. In a preferred embodiment, said expression profile includes a gene of Table 1 or Table 2.

[13] Moreover, provided herein is a biochip comprising one or more nucleic acid segments of Table 1 or Table 2, wherein the biochip comprises fewer than 1000 nucleic acid probes. Preferably at least two nucleic acid segments are included.

[14] Furthermore, a method of diagnosing a disorder associated with colorectal cancer is provided. The method comprises determining the expression of a gene of Table 1 or Table 2, in a first tissue type of a first individual, and comparing the distribution to the expression of the gene from a second normal tissue type from the first individual or a second unaffected individual. A difference in the expression indicates that the first individual has a disorder associated with colorectal cancer.

[15] In another aspect, the present invention provides an antibody which specifically binds to a protein encoded by a nucleic acid of Table 1 or Table 2 or a fragment thereof. Preferably the antibody is a monoclonal antibody. The antibody can be a fragment of an antibody such as a single stranded antibody as further described herein, or can be conjugated to another molecule. In one embodiment, the antibody is a humanized antibody.

[16] In one embodiment a method for screening for a bioactive agent capable of interfering with the binding of a colorectal cancer modulating protein (colorectal cancer modulator protein) or a fragment thereof and an antibody which binds to said colorectal cancer modulator protein or fragment thereof. In a preferred embodiment, the method comprises combining a colorectal cancer modulator protein or fragment thereof, a candidate bioactive agent and an antibody which binds to said colorectal cancer modulator protein or fragment thereof. The method further includes determining the binding of said colorectal cancer modulator protein or fragment thereof and said antibody. Wherein there is a change in binding, an agent is identified as an interfering agent. The interfering agent can be an agonist or an antagonist. Preferably, the agent inhibits colorectal cancer.

[17] In a further aspect, a method for inhibiting colorectal cancer is provided. The method can be performed in vitro or in vivo, preferably in vivo to an individual. In a preferred embodiment the method of inhibiting colorectal cancer is provided to an individual with cancer. As described herein, methods of inhibiting colorectal cancer can be performed by administering an inhibitor of the activity of a protein encoded by a nucleic acid of Table 1 or Table 2, including an antisense molecule to the gene or its gene product.

[18] Also provided herein are methods of eliciting an immune response in an individual. In one embodiment a method provided herein comprises administering to an individual a composition comprising a colorectal cancer modulating protein, or a fragment

thereof. In another embodiment, the protein is encoded by a nucleic acid selected from those of Table 1 or Table 2. In another aspect, said composition comprises a nucleic acid comprising a sequence encoding a colorectal cancer modulating protein, or a fragment thereof.

5 [19] Further provided herein are compositions capable of eliciting an immune response in an individual. In one embodiment, a composition provided herein comprises a colorectal cancer modulating protein, preferably encoded by a nucleic acid of Table 1 or Table 2, or a fragment thereof, and a pharmaceutically acceptable carrier. In another embodiment, said composition comprises a nucleic acid comprising a sequence 10 encoding a colorectal cancer modulating protein, preferably selected from the nucleic acids of Table 1 or Table 2 and a pharmaceutically acceptable carrier.

15 [20] Also provided are methods of neutralizing the effect of a colorectal cancer protein, or a fragment thereof, comprising contacting an agent specific for said protein with said protein in an amount sufficient to effect neutralization. In another embodiment, the protein is encoded by a nucleic acid selected from those of Table 1 or Table 2.

20 [21] In another aspect of the invention, a method of treating an individual for colorectal cancer is provided. In one embodiment, the method comprises administering to said individual an inhibitor of a colorectal cancer modulating protein. In another embodiment, the method comprises administering to a patient having colorectal cancer an antibody to a colorectal cancer modulating protein conjugated to a therapeutic moiety. Such a therapeutic moiety can be a cytotoxic agent or a radioisotope.

25 [22] Compounds and compositions are also provided. Other aspects of the invention will become apparent to the skilled artisan by the following description of the invention.

#### BRIEF DESCRIPTION OF THE DRAWINGS

[NOT APPLICABLE]

#### DETAILED DESCRIPTION OF THE INVENTION

30 [23] The present invention provides novel methods for diagnosis and prognosis evaluation for colorectal cancer, as well as methods for screening for compositions which modulate colorectal cancer. The methods herein are related to those of U.S. Patent Application Serial No. 09/525,993 and International Patent Application No. PCT/US00/07044, each of which is incorporated herein in its entirety.

[24] By "colorectal cancer" herein is meant a colon and/or rectal tumor or cancer that is classified as Dukes stage A or B as well as metastatic tumors classified as Dukes stage C or D (see, e.g., Cohen *et al.*, *Cancer of the Colon*, in *Cancer: Principles and Practice of Oncology*, pp. 1144-1197 (Devita *et al.*, eds., 5<sup>th</sup> ed. 1997); see also *Harrison's Principles of Internal Medicine*, pp. 1289-129 (Wilson *et al.*, eds., 12<sup>th</sup> ed., 1991)).

5 "Treatment, monitoring, detection or modulation of colorectal cancer" includes treatment, monitoring, detection, or modulation of colorectal disease in those patients who have colorectal disease (Dukes stage A , B, C or D) in which gene expression from a gene in Table 1 or 2, is increased or decreased, indicating that the subject is more likely to progress to 10 metastatic disease than a patient who does not have an increase or decrease in gene expression of a gene in Table 1 or 2. In Dukes stage A, the tumor has penetrated into, but not through, the bowel wall. In Dukes stage B, the tumor has penetrated through the bowel wall but there is not yet any lymph involvement. In Dukes stage C, the cancer involves regional lymph nodes. In Dukes stage D, there is distant metastasis, e.g., liver, lung, etc.

15 [25] Table 1 provides unigene cluster identification numbers for the nucleotide sequence of genes that exhibit increased expression in colorectal cancer samples. Tables 1 also provides an exemplar accession number that provides a nucleotide sequence that is part of the unigene cluster. Table 2 provides the nucleic acid and protein sequence of the CBF9 gene as well as the Unigene and Exemplar accession numbers for CBF9.

20 [26] In one aspect, the expression levels of genes are determined in different patient samples for which either diagnosis or prognosis information is desired, to provide expression profiles. An expression profile of a particular sample is essentially a "fingerprint" of the state of the sample; while two states may have any particular gene similarly expressed, the evaluation of a number of genes simultaneously allows the 25 generation of a gene expression profile that is unique to the state of the cell. That is, normal tissue may be distinguished from colorectal cancer tissue, and within colorectal cancer tissue, different prognosis states (good or poor long term survival prospects, for example) may be determined. By comparing expression profiles of colon tissue in known different states, information regarding which genes are important (including both up- and down- 30 regulation of genes) in each of these states is obtained. The identification of sequences that are differentially expressed in colorectal cancer versus normal colon tissue, as well as differential expression resulting in different prognostic outcomes, allows the use of this information in a number of ways. For example, the evaluation of a particular treatment regime may be evaluated: does a chemotherapeutic drug act to improve the long-term

prognosis in a particular patient. Similarly, diagnosis may be done or confirmed by comparing patient samples with the known expression profiles. Furthermore, these gene expression profiles (or individual genes) allow screening of drug candidates with an eye to mimicking or altering a particular expression profile; for example, screening can be done for drugs that suppress the colorectal cancer expression profile or convert a poor prognosis profile to a better prognosis profile. This may be done by making biochips comprising sets of the important colorectal cancer genes, which can then be used in these screens. These methods can also be done on the protein basis; that is, protein expression levels of the colorectal cancer proteins can be evaluated for diagnostic and prognostic purposes or to screen candidate agents. In addition, the colorectal cancer nucleic acid sequences can be administered for gene therapy purposes, including the administration of antisense nucleic acids, or the colorectal cancer proteins (including antibodies and other modulators thereof) administered as therapeutic drugs.

[27] Thus the present invention provides nucleic acid and protein sequences that are differentially expressed in colorectal cancer, herein termed "colorectal cancer sequences". As outlined below, colorectal cancer sequences include those that are up-regulated (i.e. expressed at a higher level) in colorectal cancer, as well as those that are down-regulated (i.e. expressed at a lower level) in colorectal cancer. In a preferred embodiment, the colorectal cancer sequences are from humans; however, as will be appreciated by those in the art, colorectal cancer sequences from other organisms may be useful in animal models of disease and drug evaluation; thus, other colorectal cancer sequences are provided, from vertebrates, including mammals, including rodents (rats, mice, hamsters, guinea pigs, etc.), primates, farm animals (including sheep, goats, pigs, cows, horses, etc.). Colorectal cancer sequences from other organisms may be obtained using the techniques outlined below.

[28] Colorectal cancer sequences can include both nucleic acid and amino acid sequences. In a preferred embodiment, the colorectal cancer sequences are recombinant nucleic acids. By the term "recombinant nucleic acid" herein is meant nucleic acid, originally formed in vitro, in general, by the manipulation of nucleic acid by polymerases and endonucleases, in a form not normally found in nature. Thus an isolated nucleic acid, in a linear form, or an expression vector formed in vitro by ligating DNA molecules that are not normally joined, are both considered recombinant for the purposes of this invention. It is understood that once a recombinant nucleic acid is made and reintroduced into a host cell or organism, it will replicate non-recombinantly, i.e. using the in vivo cellular machinery of the

host cell rather than in vitro manipulations; however, such nucleic acids, once produced recombinantly, although subsequently replicated non-recombinantly, are still considered recombinant for the purposes of the invention.

[29] Similarly, a "recombinant protein" is a protein made using recombinant techniques, i.e. through the expression of a recombinant nucleic acid as depicted above. A recombinant protein is distinguished from naturally occurring protein by at least one or more characteristics. For example, the protein may be isolated or purified away from some or all of the proteins and compounds with which it is normally associated in its wild type host, and thus may be substantially pure. For example, an isolated protein is unaccompanied by at least some of the material with which it is normally associated in its natural state, preferably constituting at least about 0.5%, more preferably at least about 5% by weight of the total protein in a given sample. A substantially pure protein comprises at least about 75% by weight of the total protein, with at least about 80% being preferred, and at least about 90% being particularly preferred. The definition includes the production of a colorectal cancer protein from one organism in a different organism or host cell. Alternatively, the protein may be made at a significantly higher concentration than is normally seen, through the use of an inducible promoter or high expression promoter, such that the protein is made at increased concentration levels. Alternatively, the protein may be in a form not normally found in nature, as in the addition of an epitope tag or amino acid substitutions, insertions and deletions, as discussed below.

[30] In a preferred embodiment, the colorectal cancer sequences are nucleic acids. As will be appreciated by those in the art and is more fully outlined below, colorectal cancer sequences are useful in a variety of applications, including diagnostic applications, which will detect naturally occurring nucleic acids, as well as screening applications; for example, biochips comprising nucleic acid probes to the colorectal cancer sequences can be generated. In the broadest sense, then, by "nucleic acid" or "oligonucleotide" or grammatical equivalents herein means at least two nucleotides covalently linked together. A nucleic acid of the present invention will generally contain phosphodiester bonds, although in some cases, as outlined below, nucleic acid analogs are included that may have alternate backbones, comprising, for example, phosphoramidate (Beaucage et al., Tetrahedron 49(10):1925 (1993) and references therein; Letsinger, J. Org. Chem. 35:3800 (1970); Sprinzel et al., Eur. J. Biochem. 81:579 (1977); Letsinger et al., Nucl. Acids Res. 14:3487 (1986); Sawai et al., Chem. Lett. 805 (1984), Letsinger et al., J. Am. Chem. Soc. 110:4470 (1988); and Pauwels et al., Chemica Scripta 26:141 91986)),

phosphorothioate (Mag et al., Nucleic Acids Res. 19:1437 (1991); and U.S. Patent No. 5,644,048), phosphorodithioate (Briu et al., J. Am. Chem. Soc. 111:2321 (1989), O-methylphophoroamidite linkages (see Eckstein, Oligonucleotides and Analogues: A Practical Approach, Oxford University Press), and peptide nucleic acid backbones and linkages (see 5 Egholm, J. Am. Chem. Soc. 114:1895 (1992); Meier et al., Chem. Int. Ed. Engl. 31:1008 (1992); Nielsen, Nature, 365:566 (1993); Carlsson et al., Nature 380:207 (1996), all of which are incorporated by reference). Other analog nucleic acids include those with positive backbones (Denpcy et al., Proc. Natl. Acad. Sci. USA 92:6097 (1995); non-ionic backbones (U.S. Patent Nos. 5,386,023, 5,637,684, 5,602,240, 5,216,141 and 4,469,863; Kiedrowski et 10 al., Angew. Chem. Intl. Ed. English 30:423 (1991); Letsinger et al., J. Am. Chem. Soc. 110:4470 (1988); Letsinger et al., Nucleoside & Nucleotide 13:1597 (1994); Chapters 2 and 3, ASC Symposium Series 580, "Carbohydrate Modifications in Antisense Research", Ed. Y.S. Sanghui and P. Dan Cook; Mesmaeker et al., Bioorganic & Medicinal Chem. Lett. 4:395 (1994); Jeffs et al., J. Biomolecular NMR 34:17 (1994); Tetrahedron Lett. 37:743 (1996)) and 15 non-ribose backbones, including those described in U.S. Patent Nos. 5,235,033 and 5,034,506, and Chapters 6 and 7, ASC Symposium Series 580, "Carbohydrate Modifications in Antisense Research", Ed. Y.S. Sanghui and P. Dan Cook. Nucleic acids containing one or more carbocyclic sugars are also included within one definition of nucleic acids (see Jenkins et al., Chem. Soc. Rev. (1995) pp169-176). Several nucleic acid analogs are described in 20 Rawls, C & E News June 2, 1997 page 35. All of these references are hereby expressly incorporated by reference. These modifications of the ribose-phosphate backbone may be done for a variety of reasons, for example to increase the stability and half-life of such molecules in physiological environments or as probes on a biochip.

[31] As will be appreciated by those in the art, all of these nucleic acid 25 analogs may find use in the present invention. In addition, mixtures of naturally occurring nucleic acids and analogs can be made; alternatively, mixtures of different nucleic acid analogs, and mixtures of naturally occurring nucleic acids and analogs may be made.

[32] Particularly preferred are peptide nucleic acids (PNA) which includes 30 peptide nucleic acid analogs. These backbones are substantially non-ionic under neutral conditions, in contrast to the highly charged phosphodiester backbone of naturally occurring nucleic acids. This results in two advantages. First, the PNA backbone exhibits improved hybridization kinetics. PNAs have larger changes in the melting temperature (T<sub>m</sub>) for mismatched versus perfectly matched basepairs. DNA and RNA typically exhibit a 2-4°C drop in T<sub>m</sub> for an internal mismatch. With the non-ionic PNA backbone, the drop is closer to

7-9°C. Similarly, due to their non-ionic nature, hybridization of the bases attached to these backbones is relatively insensitive to salt concentration. In addition, PNAs are not degraded by cellular enzymes, and thus can be more stable.

[33] The nucleic acids may be single stranded or double stranded, as specified, or contain portions of both double stranded or single stranded sequence. As will be appreciated by those in the art, the depiction of a single strand ("Watson") also defines the sequence of the other strand ("Crick"); thus the sequences described herein also includes the complement of the sequence. The nucleic acid may be DNA, both genomic and cDNA, RNA or a hybrid, where the nucleic acid contains any combination of deoxyribo- and ribo-nucleotides, and any combination of bases, including uracil, adenine, thymine, cytosine, guanine, inosine, xanthine hypoxanthine, isocytosine, isoguanine, etc. As used herein, the term "nucleoside" includes nucleotides and nucleoside and nucleotide analogs, and modified nucleosides such as amino modified nucleosides. In addition, "nucleoside" includes non-naturally occurring analog structures. Thus for example the individual units of a peptide nucleic acid, each containing a base, are referred to herein as a nucleoside.

[34] A colorectal cancer sequence can be initially identified by substantial nucleic acid and/or amino acid sequence homology to the colorectal cancer sequences outlined herein. Such homology can be based upon the overall nucleic acid or amino acid sequence, and is generally determined as outlined below, using either homology programs or hybridization conditions.

[35] The isolation of mRNA comprises isolating total cellular RNA by disrupting a cell and performing differential centrifugation. Once the total RNA is isolated, mRNA is isolated by making use of the adenine nucleotide residues known to those skilled in the art as a poly (A) tail found on virtually every eukaryotic mRNA molecule at the 3'end thereof. Oligonucleotides composed of only deoxythymidine [olgo(dT)] are linked to cellulose and the oligo(dT)-cellulose packed into small columns. When a preparation of total cellular RNA is passed through such a column, the mRNA molecules bind to the oligo(dT) by the poly (A) tails while the rest of the RNA flows through the column. The bound mRNAs are then eluted from the column and collected.

[36] The colorectal cancer sequences of the invention can be identified as follows. Samples of normal and tumor tissue are applied to biochips comprising nucleic acid probes. The samples are first microdissected, if applicable, and treated as described above for the preparation of mRNA. Suitable biochips are commercially available, for example

from Affymetrix. Gene expression profiles as described herein are generated, and the data analyzed.

[37] In a preferred embodiment, the genes showing changes in expression as between normal and disease states are compared to genes expressed in other normal tissues, including, but not limited to lung, heart, brain, liver, breast, kidney, muscle, prostate, small intestine, large intestine, spleen, bone, and placenta. In a preferred embodiment, those genes identified during the colorectal cancer screen that are expressed in any significant amount in other tissues are removed from the profile, although in some embodiments, this is not necessary. That is, when screening for drugs, it is preferable that the target be disease specific, to minimize possible side effects.

[38] In a preferred embodiment, colorectal cancer sequences are those that are up-regulated in colorectal cancer ; that is, the expression of these genes is higher in colorectal carcinoma as compared to normal colon tissue. “Up-regulation” as used herein means at least about a 1.1 fold change, preferably a 1.5 or two fold change, preferably at least about a three fold change, with at least about five-fold or higher being preferred. All accession numbers herein are for the GenBank sequence database and the sequences of the accession numbers are hereby expressly incorporated by reference. GenBank is known in the art, see, e.g., Benson, DA, et al., Nucleic Acids Research 26:1-7 (1998) and <http://www.ncbi.nlm.nih.gov/>. In addition, these genes were found to be expressed in a limited amount or not at all in heart, brain, lung, liver, breast, kidney, prostate, small intestine and spleen.

[39] In a preferred embodiment, colorectal cancer sequences are those that are down-regulated in colorectal cancer ; that is, the expression of these genes is lower in colorectal carcinoma as compared to normal colon tissue. “Down-regulation” as used herein means at least about a two-fold change, preferably at least about a three fold change, with at least about five-fold or higher being preferred.

[40] Colorectal cancer proteins of the present invention may be classified as secreted proteins, transmembrane proteins or intracellular proteins. In a preferred embodiment the colorectal cancer protein is an intracellular protein. Intracellular proteins may be found in the cytoplasm and/or in the nucleus. Intracellular proteins are involved in all aspects of cellular function and replication (including, for example, signaling pathways); aberrant expression of such proteins results in unregulated or deregulated cellular processes. For example, many intracellular proteins have enzymatic activity such as protein kinase activity, protein phosphatase activity, protease activity, nucleotide cyclase activity,

polymerase activity and the like. Intracellular proteins also serve as docking proteins that are involved in organizing complexes of proteins, or targeting proteins to various subcellular localizations, and are involved in maintaining the structural integrity of organelles.

[41] An increasingly appreciated concept in characterizing intracellular proteins is the presence in the proteins of one or more motifs for which defined functions have been attributed. In addition to the highly conserved sequences found in the enzymatic domain of proteins, highly conserved sequences have been identified in proteins that are involved in protein-protein interaction. For example, Src-homology-2 (SH2) domains bind tyrosine-phosphorylated targets in a sequence dependent manner. PTB domains, which are distinct from SH2 domains, also bind tyrosine phosphorylated targets. SH3 domains bind to proline-rich targets. In addition, PH domains, tetratricopeptide repeats and WD domains to name only a few, have been shown to mediate protein-protein interactions. Some of these may also be involved in binding to phospholipids or other second messengers. As will be appreciated by one of ordinary skill in the art, these motifs can be identified on the basis of primary sequence; thus, an analysis of the sequence of proteins may provide insight into both the enzymatic potential of the molecule and/or molecules with which the protein may associate.

[42] In a preferred embodiment, the colorectal cancer sequences are transmembrane proteins. Transmembrane proteins are molecules that span the phospholipid bilayer of a cell. They may have an intracellular domain, an extracellular domain, or both. The intracellular domains of such proteins may have a number of functions including those already described for intracellular proteins. For example, the intracellular domain may have enzymatic activity and/or may serve as a binding site for additional proteins. Frequently the intracellular domain of transmembrane proteins serves both roles. For example certain receptor tyrosine kinases have both protein kinase activity and SH2 domains. In addition, autophosphorylation of tyrosines on the receptor molecule itself, creates binding sites for additional SH2 domain containing proteins.

[43] Transmembrane proteins may contain from one to many transmembrane domains. For example, receptor tyrosine kinases, certain cytokine receptors, receptor guanylyl cyclases and receptor serine/threonine protein kinases contain a single transmembrane domain. However, various other proteins including channels and adenylyl cyclases contain numerous transmembrane domains. Many important cell surface receptors are classified as "seven transmembrane domain" proteins, as they contain 7 membrane spanning regions. Important transmembrane protein receptors include, but are not limited to

insulin receptor, insulin-like growth factor receptor, human growth hormone receptor, glucose transporters, transferrin receptor, epidermal growth factor receptor, low density lipoprotein receptor, epidermal growth factor receptor, leptin receptor, interleukin receptors, e.g. IL-1 receptor, IL-2 receptor, etc.

5 [44] Characteristics of transmembrane domains include approximately 20 consecutive hydrophobic amino acids that may be followed by charged amino acids. Therefore, upon analysis of the amino acid sequence of a particular protein, the localization and number of transmembrane domains within the protein may be predicted.

10 [45] The extracellular domains of transmembrane proteins are diverse; however, conserved motifs are found repeatedly among various extracellular domains. Conserved structure and/or functions have been ascribed to different extracellular motifs. For example, cytokine receptors are characterized by a cluster of cysteines and a WSXWS (W= tryptophan, S= serine, X=any amino acid) motif. Immunoglobulin-like domains are highly conserved. Mucin-like domains may be involved in cell adhesion and leucine-rich repeats 15 participate in protein-protein interactions.

20 [46] Many extracellular domains are involved in binding to other molecules. In one aspect, extracellular domains are receptors. Factors that bind the receptor domain include circulating ligands, which may be peptides, proteins, or small molecules such as adenosine and the like. For example, growth factors such as EGF, FGF and PDGF are 25 circulating growth factors that bind to their cognate receptors to initiate a variety of cellular responses. Other factors include cytokines, mitogenic factors, neurotrophic factors and the like. Extracellular domains also bind to cell-associated molecules. In this respect, they mediate cell-cell interactions. Cell-associated ligands can be tethered to the cell for example via a glycosylphosphatidylinositol (GPI) anchor, or may themselves be transmembrane 30 proteins. Extracellular domains also associate with the extracellular matrix and contribute to the maintenance of the cell structure.

30 [47] Colorectal cancer proteins that are transmembrane are particularly preferred in the present invention as they are good targets for immunotherapeutics, as are described herein. In addition, as outlined below, transmembrane proteins can be also useful in imaging modalities.

[48] It will also be appreciated by those in the art that a transmembrane protein can be made soluble by removing transmembrane sequences, for example through recombinant methods. Furthermore, transmembrane proteins that have been made soluble

can be made to be secreted through recombinant means by adding an appropriate signal sequence.

[49] In a preferred embodiment, the colorectal cancer proteins are secreted proteins; the secretion of which can be either constitutive or regulated. These proteins have a signal peptide or signal sequence that targets the molecule to the secretory pathway. Secreted proteins are involved in numerous physiological events; by virtue of their circulating nature, they serve to transmit signals to various other cell types. The secreted protein may function in an autocrine manner (acting on the cell that secreted the factor), a paracrine manner (acting on cells in close proximity to the cell that secreted the factor) or an endocrine manner (acting on cells at a distance). Thus secreted molecules find use in modulating or altering numerous aspects of physiology. Colorectal cancer proteins that are secreted proteins are particularly preferred in the present invention as they serve as good targets for diagnostic markers, for example for blood tests.

[50] A colorectal cancer sequence is initially identified by substantial nucleic acid and/or amino acid sequence homology to the colorectal cancer sequences outlined herein. Such homology can be based upon the overall nucleic acid or amino acid sequence, and is generally determined as outlined below, using either homology programs or hybridization conditions.

[51] As used herein, the terms "colorectal cancer nucleic acid", "colorectal cancer protein" or "colorectal cancer polynucleotide" or "colorectal cancer-associated transcript" refers to nucleic acid and polypeptide polymorphic variants, alleles, mutants, and interspecies homologs that: (1) have a nucleotide sequence that has greater than about 60% nucleotide sequence identity, 65%, 70%, 75%, 80%, 85%, 90%, preferably 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% or greater nucleotide sequence identity, preferably over a region of over a region of at least about 25, 50, 100, 200, 500, 1000, or more nucleotides, to a nucleotide sequence of or associated with a unigene cluster of Tables 1 or Table 2; (2) bind to antibodies, e.g., polyclonal antibodies, raised against an immunogen comprising an amino acid sequence encoded by a nucleotide sequence of or associated with a unigene cluster of Table 1 or Table 2, and conservatively modified variants thereof; (3) specifically hybridize under stringent hybridization conditions to a nucleic acid sequence, or the complement thereof of Table 1 or Table 2 and conservatively modified variants thereof or (4) have an amino acid sequence that has greater than about 60% amino acid sequence identity, 65%, 70%, 75%, 80%, 85%, 90%, preferably 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% or greater amino sequence identity, preferably over a region of over a region of at least about

25, 50, 100, 200, 500, 1000, or more amino acid, to an amino acid sequence encoded by a nucleotide sequence of or associated with a unigene cluster of Table 1 or Table 2. A polynucleotide or polypeptide sequence is typically from a mammal including, but not limited to, primate, e.g., human; rodent, e.g., rat, mouse, hamster; cow, pig, horse, sheep, or 5 other mammal. A “colorectal cancer polypeptide” and a “colorectal cancer polynucleotide,” include both naturally occurring or recombinant.

[52] Homology in this context means sequence similarity or identity, with identity being preferred. A preferred comparison for homology purposes is to compare the sequence containing sequencing errors to the correct sequence. This homology will be 10 determined using standard techniques known in the art, including, but not limited to, the local homology algorithm of Smith & Waterman, *Adv. Appl. Math.* 2:482 (1981), by the homology alignment algorithm of Needleman & Wunsch, *J. Mol. Biol.* 48:443 (1970), by the search for similarity method of Pearson & Lipman, *PNAS USA* 85:2444 (1988), by computerized implementations of these algorithms (GAP, BESTFIT, FASTA, and TFASTA 15 in the Wisconsin Genetics Software Package, Genetics Computer Group, 575 Science Drive, Madison, WI), the Best Fit sequence program described by Devereux et al., *Nucl. Acid Res.* 12:387-395 (1984), preferably using the default settings, or by inspection.

[53] In a preferred embodiment, the sequences which are used to determine sequence identity or similarity are selected from the sequences set forth in Table 1 or Table 2. 20 In one embodiment the sequences utilized herein are those set forth in Table 1 or Table 2. In another embodiment, the sequences are naturally occurring allelic variants of the sequences set forth in Table 1 or Table 2. In another embodiment, the sequences are sequence variants as further described herein.

[54] The terms “identical” or percent “identity,” in the context of two or 25 more nucleic acids or polypeptide sequences, refer to two or more sequences or subsequences that are the same or have a specified percentage of amino acid residues or nucleotides that are the same (i.e., about 60% identity, preferably 70%, 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or higher identity over a specified region, when compared and aligned for maximum correspondence over a comparison window or designated region) 30 as measured using a BLAST or BLAST 2.0 sequence comparison algorithms with default parameters described below, or by manual alignment and visual inspection (*see, e.g.*, NCBI web site <http://www.ncbi.nlm.nih.gov/BLAST/> or the like). Such sequences are then said to be “substantially identical.” This definition also refers to, or may be applied to, the compliment of a test sequence. The definition also includes sequences that have deletions

and/or additions, as well as those that have substitutions, as well as naturally occurring, e.g., polymorphic or allelic variants, and man-made variants. As described below, the preferred algorithms can account for gaps and the like. Preferably, identity exists over a region that is at least about 25 amino acids or nucleotides in length, or more preferably over a region that is  
5 50-100 amino acids or nucleotides in length.

[55] For sequence comparison, typically one sequence acts as a reference sequence, to which test sequences are compared. When using a sequence comparison algorithm, test and reference sequences are entered into a computer, subsequence coordinates are designated, if necessary, and sequence algorithm program parameters are designated.

10 Preferably, default program parameters can be used, or alternative parameters can be designated. The sequence comparison algorithm then calculates the percent sequence identities for the test sequences relative to the reference sequence, based on the program parameters.

[56] A "comparison window", as used herein, includes reference to a segment of one of the number of contiguous positions selected from the group consisting typically of from 20 to 600, usually about 50 to about 200, more usually about 100 to about 150 in which a sequence may be compared to a reference sequence of the same number of contiguous positions after the two sequences are optimally aligned. Methods of alignment of sequences for comparison are well-known in the art. Optimal alignment of sequences for comparison can be conducted, e.g., by the local homology algorithm of Smith & Waterman, *Adv. Appl. Math.* 2:482 (1981), by the homology alignment algorithm of Needleman & Wunsch, *J. Mol. Biol.* 48:443 (1970), by the search for similarity method of Pearson & Lipman, *Proc. Nat'l. Acad. Sci. USA* 85:2444 (1988), by computerized implementations of these algorithms (GAP, BESTFIT, FASTA, and TFASTA in the Wisconsin Genetics  
20 Software Package, Genetics Computer Group, 575 Science Dr., Madison, WI), or by manual alignment and visual inspection (see, e.g., *Current Protocols in Molecular Biology* (Ausubel et al., eds. 1995 supplement)).

[57] Preferred examples of algorithms that are suitable for determining percent sequence identity and sequence similarity include the BLAST and BLAST 2.0 algorithms, which are described in Altschul et al., *Nuc. Acids Res.* 25:3389-3402 (1977) and Altschul et al., *J. Mol. Biol.* 215:403-410 (1990). BLAST and BLAST 2.0 are used, with the parameters described herein, to determine percent sequence identity for the nucleic acids and proteins of the invention. Software for performing BLAST analyses is publicly available through the National Center for Biotechnology Information (<http://www.ncbi.nlm.nih.gov/>).

This algorithm involves first identifying high scoring sequence pairs (HSPs) by identifying short words of length W in the query sequence, which either match or satisfy some positive-valued threshold score T when aligned with a word of the same length in a database sequence. T is referred to as the neighborhood word score threshold (Altschul *et al.*, *supra*).

- 5 These initial neighborhood word hits act as seeds for initiating searches to find longer HSPs containing them. The word hits are extended in both directions along each sequence for as far as the cumulative alignment score can be increased. Cumulative scores are calculated using, e.g., for nucleotide sequences, the parameters M (reward score for a pair of matching residues; always > 0) and N (penalty score for mismatching residues; always < 0). For amino acid sequences, a scoring matrix is used to calculate the cumulative score. Extension of the word hits in each direction are halted when: the cumulative alignment score falls off by the quantity X from its maximum achieved value; the cumulative score goes to zero or below, due to the accumulation of one or more negative-scoring residue alignments; or the end of either sequence is reached. The BLAST algorithm parameters W, T, and X determine the 10 sensitivity and speed of the alignment. The BLASTN program (for nucleotide sequences) uses as defaults a wordlength (W) of 11, an expectation (E) of 10, M=5, N=-4 and a comparison of both strands. For amino acid sequences, the BLASTP program uses as 15 defaults a wordlength of 3, and expectation (E) of 10, and the BLOSUM62 scoring matrix (*see* Henikoff & Henikoff, *Proc. Natl. Acad. Sci. USA* 89:10915 (1989)) alignments (B) of 20 50, expectation (E) of 10, M=5, N=-4, and a comparison of both strands.

[58] The BLAST algorithm also performs a statistical analysis of the similarity between two sequences (*see, e.g.*, Karlin & Altschul, *Proc. Nat'l. Acad. Sci. USA* 90:5873-5787 (1993)). One measure of similarity provided by the BLAST algorithm is the smallest sum probability (P(N)), which provides an indication of the probability by which a 25 match between two nucleotide or amino acid sequences would occur by chance. For example, a nucleic acid is considered similar to a reference sequence if the smallest sum probability in a comparison of the test nucleic acid to the reference nucleic acid is less than about 0.2, more preferably less than about 0.01, and most preferably less than about 0.001. Log values may be large negative numbers, e.g., 5, 10, 20, 30, 40, 40, 70, 90, 110, 150, 170, 30 etc.

[59] In one embodiment, the nucleic acid homology is determined through hybridization studies. Thus, for example, nucleic acids which hybridize under high stringency to the nucleic acid sequences which encode the peptides identified in Table 1 or Table 2, or their complements, are considered a colorectal cancer sequence. High stringency

conditions are known in the art; see for example Maniatis et al., Molecular Cloning: A Laboratory Manual, 2d Edition, 1989, and Short Protocols in Molecular Biology, ed. Ausubel, et al., both of which are hereby incorporated by reference. Stringent conditions are sequence-dependent and will be different in different circumstances. Longer sequences hybridize specifically at higher temperatures. An extensive guide to the hybridization of nucleic acids is found in Tijssen, Techniques in Biochemistry and Molecular Biology-- Hybridization with Nucleic Acid Probes, "Overview of principles of hybridization and the strategy of nucleic acid assays" (1993). Generally, stringent conditions are selected to be about 5-10°C lower than the thermal melting point (Tm) for the specific sequence at a defined ionic strength pH. The Tm is the temperature (under defined ionic strength, pH and nucleic acid concentration) at which 50% of the probes complementary to the target hybridize to the target sequence at equilibrium (as the target sequences are present in excess, at Tm, 50% of the probes are occupied at equilibrium). Stringent conditions will be those in which the salt concentration is less than about 1.0 M sodium ion, typically about 0.01 to 1.0 M sodium ion concentration (or other salts) at pH 7.0 to 8.3 and the temperature is at least about 30°C for short probes (e.g. 10 to 50 nucleotides) and at least about 60°C for long probes (e.g. greater than 50 nucleotides). Stringent conditions may also be achieved with the addition of destabilizing agents such as formamide.

[60] In another embodiment, less stringent hybridization conditions are used; for example, moderate or low stringency conditions may be used, as are known in the art; see Maniatis and Ausubel, *supra*, and Tijssen, *supra*. For selective or specific hybridization, a positive signal is at least two times background, preferably 10 times background hybridization. Exemplary stringent hybridization conditions can be as following: 50% formamide, 5x SSC, and 1% SDS, incubating at 42°C, or, 5x SSC, 1% SDS, incubating at 65°C, with wash in 0.2x SSC, and 0.1% SDS at 65°C.

[61] Nucleic acids that do not hybridize to each other under stringent conditions are still substantially identical if the polypeptides which they encode are substantially identical. This occurs, for example, when a copy of a nucleic acid is created using the maximum codon degeneracy permitted by the genetic code. In such cases, the nucleic acids typically hybridize under moderately stringent hybridization conditions. Exemplary "moderately stringent hybridization conditions" include a hybridization in a buffer of 40% formamide, 1 M NaCl, 1% SDS at 37°C, and a wash in 1X SSC at 45°C. A positive hybridization is at least twice background. Those of ordinary skill will readily

recognize that alternative hybridization and wash conditions can be utilized to provide conditions of similar stringency. Additional guidelines for determining hybridization parameters are provided in numerous reference, e.g., and *Current Protocols in Molecular Biology*, ed. Ausubel, *et al.*

5 [62] For PCR, a temperature of about 36°C is typical for low stringency amplification, although annealing temperatures may vary between about 32°C and 48°C depending on primer length. For high stringency PCR amplification, a temperature of about 62°C is typical, although high stringency annealing temperatures can range from about 50°C to about 65°C, depending on the primer length and specificity. Typical cycle conditions for  
10 both high and low stringency amplifications include a denaturation phase of 90°C - 95°C for 30 sec - 2 min., an annealing phase lasting 30 sec. - 2 min., and an extension phase of about 72°C for 1 - 2 min. Protocols and guidelines for low and high stringency amplification reactions are provided, e.g., in Innis *et al.*, *PCR Protocols, A Guide to Methods and Applications* (1990).

15 [63] In addition, the colorectal cancer nucleic acid sequences of the invention are fragments of larger genes, i.e. they are nucleic acid segments. "Genes" in this context includes coding regions, non-coding regions, and mixtures of coding and non-coding regions. Accordingly, as will be appreciated by those in the art, using the sequences provided herein, additional sequences of the colorectal cancer genes can be obtained, using techniques  
20 well known in the art for cloning either longer sequences or the full length sequences; see Maniatis *et al.*, and Ausubel, *et al.*, *supra*, hereby expressly incorporated by reference.

25 [64] An indication that two nucleic acid sequences or polypeptides are substantially identical is that the polypeptide encoded by the first nucleic acid is immunologically cross reactive with the antibodies raised against the polypeptide encoded by the second nucleic acid. Thus, a polypeptide is typically substantially identical to a second polypeptide, e.g., where the two peptides differ only by conservative substitutions. Another indication that two nucleic acid sequences are substantially identical is that the two molecules or their complements hybridize to each other under stringent conditions, as described above.  
30 Yet another indication that two nucleic acid sequences are substantially identical is that the same primers can be used to amplify the sequences.

[65] Once the colorectal cancer nucleic acid is identified, it can be cloned and, if necessary, its constituent parts recombined to form the entire colorectal cancer nucleic acid. Once isolated from its natural source, e.g., contained within a plasmid or other vector

or excised therefrom as a linear nucleic acid segment, the recombinant colorectal cancer nucleic acid can be further-used as a probe to identify and isolate other colorectal cancer nucleic acids, for example additional coding regions. It can also be used as a "precursor" nucleic acid to make modified or variant colorectal cancer nucleic acids and proteins.

5 [66] The colorectal cancer nucleic acids of the present invention are used in several ways. In a first embodiment, nucleic acid probes to the colorectal cancer nucleic acids are made and attached to biochips to be used in screening and diagnostic methods, as outlined below, or for administration, for example for gene therapy and/or antisense applications. Alternatively, the colorectal cancer nucleic acids that include coding regions of  
10 colorectal cancer proteins can be put into expression vectors for the expression of colorectal cancer proteins, again either for screening purposes or for administration to a patient.

15 [67] In a preferred embodiment, nucleic acid probes to colorectal cancer nucleic acids (both the nucleic acid sequences encoding peptides outlined in the Table 1 or Table 2 and/or the complements thereof) are made. The nucleic acid probes attached to the biochip are designed to be substantially complementary to the colorectal cancer nucleic acids, i.e. the target sequence (either the target sequence of the sample or to other probe sequences, for example in sandwich assays), such that hybridization of the target sequence and the probes of the present invention occurs. As outlined below, this complementarity need not be perfect; there may be any number of base pair mismatches which will interfere with  
20 hybridization between the target sequence and the single stranded nucleic acids of the present invention. However, if the number of mutations is so great that no hybridization can occur under even the least stringent of hybridization conditions, the sequence is not a complementary target sequence. Thus, by "substantially complementary" herein is meant that the probes are sufficiently complementary to the target sequences to hybridize under  
25 normal reaction conditions, particularly high stringency conditions, as outlined herein.

30 [68] A nucleic acid probe is generally single stranded but can be partially single and partially double stranded. The strandedness of the probe is dictated by the structure, composition, and properties of the target sequence. In general, the nucleic acid probes range from about 8 to about 100 bases long, with from about 10 to about 80 bases being preferred, and from about 30 to about 50 bases being particularly preferred. That is, generally whole genes are not used. In some embodiments, much longer nucleic acids can be used, up to hundreds of bases.

[69] In a preferred embodiment, more than one probe per sequence is used, with either overlapping probes or probes to different sections of the target being used. That

is, two, three, four or more probes, with three being preferred, are used to build in a redundancy for a particular target. The probes can be overlapping (i.e. have some sequence in common), or separate.

[70] As will be appreciated by those in the art, nucleic acids can be attached or immobilized to a solid support in a wide variety of ways. By "immobilized" and grammatical equivalents herein is meant the association or binding between the nucleic acid probe and the solid support is sufficient to be stable under the conditions of binding, washing, analysis, and removal as outlined below. The binding can be covalent or non-covalent. By "non-covalent binding" and grammatical equivalents herein is meant one or more of either electrostatic, hydrophilic, and hydrophobic interactions. Included in non-covalent binding is the covalent attachment of a molecule, such as, streptavidin to the support and the non-covalent binding of the biotinylated probe to the streptavidin. By "covalent binding" and grammatical equivalents herein is meant that the two moieties, the solid support and the probe, are attached by at least one bond, including sigma bonds, pi bonds and coordination bonds. Covalent bonds can be formed directly between the probe and the solid support or can be formed by a cross linker or by inclusion of a specific reactive group on either the solid support or the probe or both molecules. Immobilization may also involve a combination of covalent and non-covalent interactions.

[71] In general, the probes are attached to the biochip in a wide variety of ways, as will be appreciated by those in the art. As described herein, the nucleic acids can either be synthesized first, with subsequent attachment to the biochip, or can be directly synthesized on the biochip.

[72] The biochip comprises a suitable solid substrate. By "substrate" or "solid support" or other grammatical equivalents herein is meant any material that can be modified to contain discrete individual sites appropriate for the attachment or association of the nucleic acid probes and is amenable to at least one detection method. As will be appreciated by those in the art, the number of possible substrates are very large, and include, but are not limited to, glass and modified or functionalized glass, plastics (including acrylics, polystyrene and copolymers of styrene and other materials, polypropylene, polyethylene, polybutylene, polyurethanes, TeflonJ, etc.), polysaccharides, nylon or nitrocellulose, resins, silica or silica-based materials including silicon and modified silicon, carbon, metals, inorganic glasses, plastics, etc. In general, the substrates allow optical detection and do not appreciably fluoresce. A preferred substrate is described in copending application entitled

Reusable Low Fluorescent Plastic Biochip, U.S. Application Serial No. 09/270,214, filed March 15, 1999, herein incorporated by reference in its entirety.

[73] Generally the substrate is planar, although as will be appreciated by those in the art, other configurations of substrates may be used as well. For example, the probes may be placed on the inside surface of a tube, for flow-through sample analysis to minimize sample volume. Similarly, the substrate may be flexible, such as a flexible foam, including closed cell foams made of particular plastics.

[74] In a preferred embodiment, the surface of the biochip and the probe may be derivatized with chemical functional groups for subsequent attachment of the two. Thus, for example, the biochip is derivatized with a chemical functional group including, but not limited to, amino groups, carboxy groups, oxo groups and thiol groups, with amino groups being particularly preferred. Using these functional groups, the probes can be attached using functional groups on the probes. For example, nucleic acids containing amino groups can be attached to surfaces comprising amino groups, for example using linkers as are known in the art; for example, homo- or hetero-bifunctional linkers as are well known (see 1994 Pierce Chemical Company catalog, technical section on cross-linkers, pages 155-200, incorporated herein by reference). In addition, in some cases, additional linkers, such as alkyl groups (including substituted and heteroalkyl groups) may be used.

[75] In this embodiment, the oligonucleotides are synthesized as is known in the art, and then attached to the surface of the solid support. As will be appreciated by those skilled in the art, either the 5' or 3' terminus may be attached to the solid support, or attachment may be via an internal nucleoside.

[76] In an additional embodiment, the immobilization to the solid support may be very strong, yet non-covalent. For example, biotinylated oligonucleotides can be made, which bind to surfaces covalently coated with streptavidin, resulting in attachment.

[77] Alternatively, the oligonucleotides may be synthesized on the surface, as is known in the art. For example, photoactivation techniques utilizing photopolymerization compounds and techniques are used. In a preferred embodiment, the nucleic acids can be synthesized *in situ*, using well known photolithographic techniques, such as those described in WO 95/25116; WO 95/35505; U.S. Patent Nos. 5,700,637 and 5,445,934; and references cited within, all of which are expressly incorporated by reference; these methods of attachment form the basis of the Affymetrix GeneChip™ technology.

[78] In a preferred embodiment, colorectal cancer nucleic acids encoding colorectal cancer proteins are used to make a variety of expression vectors to express colorectal cancer proteins which can then be used in screening assays, as described below. The expression vectors may be either self-replicating extrachromosomal vectors or vectors which integrate into a host genome. Generally, these expression vectors include transcriptional and translational regulatory nucleic acid operably linked to the nucleic acid encoding the colorectal cancer protein. The term "control sequences" refers to DNA sequences necessary for the expression of an operably linked coding sequence in a particular host organism. The control sequences that are suitable for prokaryotes, for example, include a promoter, optionally an operator sequence, and a ribosome binding site. Eukaryotic cells are known to utilize promoters, polyadenylation signals, and enhancers.

[79] Nucleic acid is "operably linked" when it is placed into a functional relationship with another nucleic acid sequence. For example, DNA for a presequence or secretory leader is operably linked to DNA for a polypeptide if it is expressed as a preprotein that participates in the secretion of the polypeptide; a promoter or enhancer is operably linked to a coding sequence if it affects the transcription of the sequence; or a ribosome binding site is operably linked to a coding sequence if it is positioned so as to facilitate translation. Generally, "operably linked" means that the DNA sequences being linked are contiguous, and, in the case of a secretory leader, contiguous and in reading phase. However, enhancers do not have to be contiguous. Linking is accomplished by ligation at convenient restriction sites. If such sites do not exist, the synthetic oligonucleotide adaptors or linkers are used in accordance with conventional practice. The transcriptional and translational regulatory nucleic acid will generally be appropriate to the host cell used to express the colorectal cancer protein; for example, transcriptional and translational regulatory nucleic acid sequences from *Bacillus* are preferably used to express the colorectal cancer protein in *Bacillus*. Numerous types of appropriate expression vectors, and suitable regulatory sequences are known in the art for a variety of host cells.

[80] In general, the transcriptional and translational regulatory sequences may include, but are not limited to, promoter sequences, ribosomal binding sites, transcriptional start and stop sequences, translational start and stop sequences, and enhancer or activator sequences. In a preferred embodiment, the regulatory sequences include a promoter and transcriptional start and stop sequences.

[81] Promoter sequences encode either constitutive or inducible promoters. The promoters may be either naturally occurring promoters or hybrid promoters. Hybrid

promoters, which combine elements of more than one promoter, are also known in the art, and are useful in the present invention.

[82] In addition, the expression vector may comprise additional elements. For example, the expression vector may have two replication systems, thus allowing it to be maintained in two organisms, for example in mammalian or insect cells for expression and in a prokaryotic host for cloning and amplification. Furthermore, for integrating expression vectors, the expression vector contains at least one sequence homologous to the host cell genome, and preferably two homologous sequences which flank the expression construct. The integrating vector may be directed to a specific locus in the host cell by selecting the appropriate homologous sequence for inclusion in the vector. Constructs for integrating vectors are well known in the art.

[83] In addition, in a preferred embodiment, the expression vector contains a selectable marker gene to allow the selection of transformed host cells. Selection genes are well known in the art and will vary with the host cell used.

[84] The colorectal cancer proteins of the present invention are produced by culturing a host cell transformed with an expression vector containing nucleic acid encoding a colorectal cancer protein, under the appropriate conditions to induce or cause expression of the colorectal cancer protein. The conditions appropriate for colorectal cancer protein expression will vary with the choice of the expression vector and the host cell, and will be easily ascertained by one skilled in the art through routine experimentation. For example, the use of constitutive promoters in the expression vector will require optimizing the growth and proliferation of the host cell, while the use of an inducible promoter requires the appropriate growth conditions for induction. In addition, in some embodiments, the timing of the harvest is important. For example, the baculoviral systems used in insect cell expression are lytic viruses, and thus harvest time selection can be crucial for product yield.

[85] Appropriate host cells include yeast, bacteria, archaeabacteria, fungi, and insect and animal cells, including mammalian cells. Of particular interest are Drosophila melanogaster cells, *Saccharomyces cerevisiae* and other yeasts, *E. coli*, *Bacillus subtilis*, Sf9 cells, C129 cells, 293 cells, *Neurospora*, BHK, CHO, COS, HeLa cells, THP1 cell line (a macrophage cell line) and human cells and cell lines.

[86] In a preferred embodiment, the colorectal cancer proteins are expressed in mammalian cells. Mammalian expression systems are also known in the art, and include retroviral systems. A preferred expression vector system is a retroviral vector system such as is generally described in PCT/US97/01019 and PCT/US97/01048, both of which are

hereby expressly incorporated by reference. Of particular use as mammalian promoters are the promoters from mammalian viral genes, since the viral genes are often highly expressed and have a broad host range. Examples include the SV40 early promoter, mouse mammary tumor virus LTR promoter, adenovirus major late promoter, herpes simplex virus promoter, 5 and the CMV promoter. Typically, transcription termination and polyadenylation sequences recognized by mammalian cells are regulatory regions located 3' to the translation stop codon and thus, together with the promoter elements, flank the coding sequence. Examples of transcription terminator and polyadenylation signals include those derived from SV40.

[87] The methods of introducing exogenous nucleic acid into mammalian hosts, as well as other hosts, is well known in the art, and will vary with the host cell used. Techniques include dextran-mediated transfection, calcium phosphate precipitation, polybrene mediated transfection, protoplast fusion, electroporation, viral infection, encapsulation of the polynucleotide(s) in liposomes, and direct microinjection of the DNA into nuclei.

[88] In a preferred embodiment, colorectal cancer proteins are expressed in bacterial systems. Bacterial expression systems are well known in the art. Promoters from bacteriophage may also be used and are known in the art. In addition, synthetic promoters and hybrid promoters are also useful; for example, the tac promoter is a hybrid of the trp and lac promoter sequences. Furthermore, a bacterial promoter can include naturally occurring promoters of non-bacterial origin that have the ability to bind bacterial RNA polymerase and initiate transcription. In addition to a functioning promoter sequence, an efficient ribosome binding site is desirable. The expression vector may also include a signal peptide sequence that provides for secretion of the colorectal cancer protein in bacteria. The protein is either secreted into the growth media (gram-positive bacteria) or into the periplasmic space, located 20 between the inner and outer membrane of the cell (gram-negative bacteria). The bacterial expression vector may also include a selectable marker gene to allow for the selection of bacterial strains that have been transformed. Suitable selection genes include genes which render the bacteria resistant to drugs such as ampicillin, chloramphenicol, erythromycin, kanamycin, neomycin and tetracycline. Selectable markers also include biosynthetic genes, 25 such as those in the histidine, tryptophan and leucine biosynthetic pathways. These components are assembled into expression vectors. Expression vectors for bacteria are well known in the art, and include vectors for *Bacillus subtilis*, *E. coli*, *Streptococcus cremoris*, and *Streptococcus lividans*, among others. The bacterial expression vectors are transformed 30

into bacterial host cells using techniques well known in the art, such as calcium chloride treatment, electroporation, and others.

[89] In one embodiment, colorectal cancer proteins are produced in insect cells. Expression vectors for the transformation of insect cells, and in particular, baculovirus-based expression vectors, are well known in the art.

[90] In a preferred embodiment, colorectal cancer protein is produced in yeast cells. Yeast expression systems are well known in the art, and include expression vectors for *Saccharomyces cerevisiae*, *Candida albicans* and *C. maltosa*, *Hansenula polymorpha*, *Kluyveromyces fragilis* and *K. lactis*, *Pichia guillerimondii* and *P. pastoris*, *Schizosaccharomyces pombe*, and *Yarrowia lipolytica*.

[91] The colorectal cancer protein may also be made as a fusion protein, using techniques well known in the art. Thus, for example, for the creation of monoclonal antibodies, if the desired epitope is small, the colorectal cancer protein may be fused to a carrier protein to form an immunogen. Alternatively, the colorectal cancer protein may be made as a fusion protein to increase expression, or for other reasons. For example, when the colorectal cancer protein is a colorectal cancer peptide, the nucleic acid encoding the peptide may be linked to other nucleic acid for expression purposes.

[92] In one embodiment, the colorectal cancer nucleic acids, proteins and antibodies of the invention are labeled. By "labeled" herein is meant that a compound has at least one element, isotope or chemical compound attached to enable the detection of the compound. In general, labels fall into three classes: a) isotopic labels, which may be radioactive or heavy isotopes; b) immune labels, which may be antibodies or antigens; and c) colored or fluorescent dyes. The labels may be incorporated into the colorectal cancer nucleic acids, proteins and antibodies at any position. For example, the label should be capable of producing, either directly or indirectly, a detectable signal. The detectable moiety may be a radioisotope, such as  $^3\text{H}$ ,  $^{14}\text{C}$ ,  $^{32}\text{P}$ ,  $^{35}\text{S}$ , or  $^{125}\text{I}$ , a fluorescent or chemiluminescent compound, such as fluorescein isothiocyanate, rhodamine, or luciferin, or an enzyme, such as alkaline phosphatase, beta-galactosidase or horseradish peroxidase. Any method known in the art for conjugating the antibody to the label may be employed, including those methods described by Hunter et al., *Nature*, 144:945 (1962); David et al., *Biochemistry*, 13:1014 (1974); Pain et al., *J. Immunol. Meth.*, 40:219 (1981); and Nygren, *J. Histochem. and Cytochem.*, 30:407 (1982).

[93] Accordingly, the present invention also provides colorectal cancer protein sequences. A colorectal cancer protein of the present invention may be identified in

several ways. "Protein" in this sense includes proteins, polypeptides, and peptides terms which are used interchangeably herein to refer to a polymer of amino acid residues. The terms apply to amino acid polymers in which one or more amino acid residue is an artificial chemical mimetic of a corresponding naturally occurring amino acid, as well as to naturally occurring amino acid polymers, those containing modified residues, and non-naturally occurring amino acid polymer.

[94] As will be appreciated by those in the art, the nucleic acid sequences of the invention can be used to generate protein sequences. There are a variety of ways to do this, including cloning the entire gene and verifying its frame and amino acid sequence, or by comparing it to known sequences to search for homology to provide a frame, assuming the colorectal cancer protein has homology to some protein in the database being used. Generally, the nucleic acid sequences are input into a program that will search all three frames for homology. This is done in a preferred embodiment using the following NCBI Advanced BLAST parameters. The program is blastx or blastn. The database is nr. The input data is as "Sequence in FASTA format". The organism list is "none". The "expect" is 10; the filter is default. The "descriptions" is 500, the "alignments" is 500, and the "alignment view" is pairwise. The "Query Genetic Codes" is standard (1). The matrix is BLOSUM62; gap existence cost is 11, per residue gap cost is 1; and the lambda ratio is .85 default. This results in the generation of a putative protein sequence.

[95] Also included within one embodiment of colorectal cancer proteins are amino acid variants of the naturally occurring sequences, as determined herein. Preferably, the variants are preferably greater than about 75% homologous to the wild-type sequence, more preferably greater than about 80%, even more preferably greater than about 85% and most preferably greater than 90%. In some embodiments the homology will be as high as about 93 to 95 or 98%. As for nucleic acids, homology in this context means sequence similarity or identity, with identity being preferred. This homology will be determined using standard techniques known in the art as are outlined above for the nucleic acid homologies.

[96] Colorectal cancer proteins of the present invention may be shorter or longer than the wild type amino acid sequences. Thus, in a preferred embodiment, included within the definition of colorectal cancer proteins are portions or fragments of the wild type sequences. herein. In addition, as outlined above, the colorectal cancer nucleic acids of the invention may be used to obtain additional coding regions, and thus additional protein sequence, using techniques known in the art.

[97] In a preferred embodiment, the colorectal cancer proteins are derivative or variant colorectal cancer proteins as compared to the wild-type sequence. That is, as outlined more fully below, the derivative colorectal cancer peptide will contain at least one amino acid substitution, deletion or insertion, with amino acid substitutions being particularly preferred. The amino acid substitution, insertion or deletion may occur at any residue within the colorectal cancer peptide.

[98] Also included in an embodiment of colorectal cancer proteins of the present invention are amino acid sequence variants. These variants fall into one or more of three classes: substitutional, insertional or deletional variants. These variants ordinarily are prepared by site specific mutagenesis of nucleotides in the DNA encoding the colorectal cancer protein, using cassette or PCR mutagenesis or other techniques well known in the art, to produce DNA encoding the variant, and thereafter expressing the DNA in recombinant cell culture as outlined above. However, variant colorectal cancer protein fragments having up to about 100-150 residues may be prepared by in vitro synthesis using established techniques.

Amino acid sequence variants are characterized by the predetermined nature of the variation, a feature that sets them apart from naturally occurring allelic or interspecies variation of the colorectal cancer protein amino acid sequence. The variants typically exhibit the same qualitative biological activity as the naturally occurring analogue, although variants can also be selected which have modified characteristics as will be more fully outlined below.

[99] While the site or region for introducing an amino acid sequence variation is predetermined, the mutation per se need not be predetermined. For example, in order to optimize the performance of a mutation at a given site, random mutagenesis may be conducted at the target codon or region and the expressed colorectal cancer variants screened for the optimal combination of desired activity. Techniques for making substitution mutations at predetermined sites in DNA having a known sequence are well known, for example, M13 primer mutagenesis and PCR mutagenesis. Screening of the mutants is done using assays of colorectal cancer protein activities.

[100] Amino acid substitutions are typically of single residues; insertions usually will be on the order of from about 1 to 20 amino acids, although considerably larger insertions may be tolerated. Deletions range from about 1 to about 20 residues, although in some cases deletions may be much larger.

[101] Substitutions, deletions, insertions or any combination thereof may be used to arrive at a final derivative. Generally these changes are done on a few amino acids to minimize the alteration of the molecule. However, larger changes may be tolerated in certain

circumstances. When small alterations in the characteristics of the colorectal cancer protein are desired, substitutions are generally made in accordance with the following chart:

**Chart I**

	Original Residue	Exemplary Substitutions
5		
	Ala	Ser
	Arg	Lys
	Asn	Gln, His
	Asp	Glu
10	Cys	Ser
	Gln	Asn
	Glu	Asp
	Gly	Pro
	His	Asn, Gln
15	Ile	Leu, Val
	Leu	Ile, Val
	Lys	Arg, Gln, Glu
	Met	Leu, Ile
	Phe	Met, Leu, Tyr
20	Ser	Thr
	Thr	Ser
	Trp	Tyr
	Tyr	Trp, Phe
	Val	Ile, Leu
25		

[102] Substantial changes in function or immunological identity are made by selecting substitutions that are less conservative than those shown in Chart I. For example, substitutions may be made which more significantly affect: the structure of the polypeptide backbone in the area of the alteration, for example the alpha-helical or beta-sheet structure; the charge or hydrophobicity of the molecule at the target site; or the bulk of the side chain. The substitutions which in general are expected to produce the greatest changes in the polypeptide's properties are those in which (a) a hydrophilic residue, e.g. seryl or threonyl is substituted for (or by) a hydrophobic residue, e.g. leucyl, isoleucyl, phenylalanyl, valyl or alanyl; (b) a cysteine or proline is substituted for (or by) any other residue; (c) a residue

having an electropositive side chain, e.g. lysyl, arginyl, or histidyl, is substituted for (or by) an electronegative residue, e.g. glutamyl or aspartyl; or (d) a residue having a bulky side chain, e.g. phenylalanine, is substituted for (or by) one not having a side chain, e.g. glycine.

[103] The variants typically exhibit the same qualitative biological activity

5 and will elicit the same immune response as the naturally-occurring analogue, although variants also are selected to modify the characteristics of the colorectal cancer proteins as needed. Alternatively, the variant may be designed such that the biological activity of the colorectal cancer protein is altered. For example, glycosylation sites may be altered or removed.

10 [104] Covalent modifications of colorectal cancer polypeptides are included within the scope of this invention. One type of covalent modification includes reacting targeted amino acid residues of a colorectal cancer polypeptide with an organic derivatizing agent that is capable of reacting with selected side chains or the N-or C-terminal residues of a colorectal cancer polypeptide. Derivatization with bifunctional agents is useful, for instance, 15 for crosslinking colorectal cancer to a water-insoluble support matrix or surface for use in the method for purifying anti-colorectal cancer antibodies or screening assays, as is more fully described below. Commonly used crosslinking agents include, e.g., 1,1-bis(diazo-acetyl)-2-phenylethane, glutaraldehyde, N-hydroxy-succinimide esters, for example, esters with 4-azido-salicylic acid, homobifunctional imidoesters, including disuccinimidyl esters 20 such as 3,3'-dithiobis-(succinimidyl-propionate), bifunctional maleimides such as bis-N-maleimido-1,8-octane and agents such as methyl-3-[*(p*-azidophenyl)-dithio]pro-pioimi-date.

25 [105] Other modifications include deamidation of glutaminyl and asparaginyl residues to the corresponding glutamyl and aspartyl residues, respectively, hydroxylation of proline and lysine, phosphorylation of hydroxyl groups of seryl, threonyl or tyrosyl residues, methylation of the  $\alpha$ -amino groups of lysine, arginine, and histidine side chains [T.E. Creighton, Proteins: Structure and Molecular Properties, W.H. Freeman & Co., San Francisco, pp. 79-86 (1983)], acetylation of the N-terminal amine, and amidation of any C-terminal carboxyl group.

30 [106] Another type of covalent modification of the colorectal cancer polypeptide included within the scope of this invention comprises altering the native glycosylation pattern of the polypeptide. "Altering the native glycosylation pattern" is intended for purposes herein to mean deleting one or more carbohydrate moieties found in native sequence colorectal cancer polypeptide, and/or adding one or more glycosylation sites that are not present in the native sequence colorectal cancer polypeptide.

[107] Addition of glycosylation sites to colorectal cancer polypeptides may be accomplished by altering the amino acid sequence thereof. The alteration may be made, for example, by the addition of, or substitution by, one or more serine or threonine residues to the native sequence colorectal cancer polypeptide (for O-linked glycosylation sites). The 5 colorectal cancer amino acid sequence may optionally be altered through changes at the DNA level, particularly by mutating the DNA encoding the colorectal cancer polypeptide at preselected bases such that codons are generated that will translate into the desired amino acids.

[108] Another means of increasing the number of carbohydrate moieties on 10 the colorectal cancer polypeptide is by chemical or enzymatic coupling of glycosides to the polypeptide. Such methods are described in the art, e.g., in WO 87/05330 published 11 September 1987, and in Aplin and Wriston, colorectal cancer Crit. Rev. Biochem., pp. 259-306 (1981).

[109] Removal of carbohydrate moieties present on the colorectal cancer 15 polypeptide may be accomplished chemically or enzymatically or by mutational substitution of codons encoding for amino acid residues that serve as targets for glycosylation. Chemical deglycosylation techniques are known in the art and described, for instance, by Hakimuddin, et al., Arch. Biochem. Biophys., 259:52 (1987) and by Edge et al., Anal. Biochem., 118:131 (1981). Enzymatic cleavage of carbohydrate moieties on polypeptides can be achieved by the 20 use of a variety of endo-and exo-glycosidases as described by Thotakura et al., Meth. Enzymol., 138:350 (1987).

[110] Another type of covalent modification of colorectal cancer comprises linking the colorectal cancer polypeptide to one of a variety of nonproteinaceous polymers, e.g., polyethylene glycol, polypropylene glycol, or polyoxyalkylenes, in the manner set forth 25 in U.S. Patent Nos. 4,640,835; 4,496,689; 4,301,144; 4,670,417; 4,791,192 or 4,179,337.

[111] colorectal cancer polypeptides of the present invention may also be modified in a way to form chimeric molecules comprising a colorectal cancer polypeptide fused to another, heterologous polypeptide or amino acid sequence. In one embodiment, such 30 a chimeric molecule comprises a fusion of a colorectal cancer polypeptide with a tag polypeptide which provides an epitope to which an anti-tag antibody can selectively bind. The epitope tag is generally placed at the amino-or carboxyl-terminus of the colorectal cancer polypeptide. The presence of such epitope-tagged forms of a colorectal cancer polypeptide can be detected using an antibody against the tag polypeptide. Also, provision of the epitope tag enables the colorectal cancer polypeptide to be readily purified by affinity purification

using an anti-tag antibody or another type of affinity matrix that binds to the epitope tag. In an alternative embodiment, the chimeric molecule may comprise a fusion of a colorectal cancer polypeptide with an immunoglobulin or a particular region of an immunoglobulin. For a bivalent form of the chimeric molecule, such a fusion could be to the Fc region of an IgG molecule.

[112] Various tag polypeptides and their respective antibodies are well known in the art. Examples include poly-histidine (poly-his) or poly-histidine-glycine (poly-his-gly) tags; the flu HA tag polypeptide and its antibody 12CA5 [Field et al., Mol. Cell. Biol., 8:2159-2165 (1988)]; the c-myc tag and the 8F9, 3C7, 6E10, G4, B7 and 9E10 antibodies thereto [Evan et al., Molecular and Cellular Biology, 5:3610-3616 (1985)]; and the Herpes Simplex virus glycoprotein D (gD) tag and its antibody [Paborsky et al., Protein Engineering, 3(6):547-553 (1990)]. Other tag polypeptides include the Flag-peptide [Hopp et al., BioTechnology, 6:1204-1210 (1988)]; the KT3 epitope peptide [Martin et al., Science, 255:192-194 (1992)]; tubulin epitope peptide [Skinner et al., J. Biol. Chem., 266:15163-15166 (1991)]; and the T7 gene 10 protein peptide tag [Lutz-Freyermuth et al., Proc. Natl. Acad. Sci. USA, 87:6393-6397 (1990)].

[113] Also included with the definition of colorectal cancer protein in one embodiment are other colorectal cancer proteins of the colorectal cancer family, and colorectal cancer proteins from other organisms, which are cloned and expressed as outlined below. Thus, probe or degenerate polymerase chain reaction (PCR) primer sequences may be used to find other related colorectal cancer proteins from humans or other organisms. As will be appreciated by those in the art, particularly useful probe and/or PCR primer sequences include the unique areas of the colorectal cancer nucleic acid sequence. As is generally known in the art, preferred PCR primers are from about 15 to about 35 nucleotides in length, with from about 20 to about 30 being preferred, and may contain inosine as needed. The conditions for the PCR reaction are well known in the art.

[114] In addition, as is outlined herein, colorectal cancer proteins can be made that are longer than those depicted in the Table 1 or Table 2 for example, by the elucidation of additional sequences, the addition of epitope or purification tags, the addition of other fusion sequences, etc.

[115] Colorectal cancer proteins may also be identified as being encoded by colorectal cancer nucleic acids. Thus, colorectal cancer proteins are encoded by nucleic acids that will hybridize to the sequences of the sequence listings, or their complements, as outlined herein.

[116] In a preferred embodiment, when the colorectal cancer protein is to be used to generate antibodies, for example for immunotherapy, the colorectal cancer protein should share at least one epitope or determinant with the full length protein. By "epitope" or "determinant" herein is meant a portion of a protein which will generate and/or bind an antibody or T-cell receptor in the context of MHC. Thus, in most instances, antibodies made to a smaller colorectal cancer protein will be able to bind to the full length protein. In a preferred embodiment, the epitope is unique; that is, antibodies generated to a unique epitope show little or no cross-reactivity. In a preferred embodiment, the epitope is selected from a peptide encoded by a nucleic acid of Table 1. In another preferred embodiment, the epitope is selected from the CBF9 peptide sequence shown in Table 2.

[117] In one embodiment, the term "antibody" includes antibody fragments, as are known in the art, including Fab, Fab2, single chain antibodies (Fv for example), chimeric antibodies, etc., either produced by the modification of whole antibodies or those synthesized de novo using recombinant DNA technologies.

[118] Methods of preparing polyclonal antibodies are known to the skilled artisan. Polyclonal antibodies can be raised in a mammal, for example, by one or more injections of an immunizing agent and, if desired, an adjuvant. Typically, the immunizing agent and/or adjuvant will be injected in the mammal by multiple subcutaneous or intraperitoneal injections. The immunizing agent may include the CBF9 peptide of Table 2, or a peptide encoded by a nucleic acid of Table 1 or fragment thereof or a fusion protein thereof. It may be useful to conjugate the immunizing agent to a protein known to be immunogenic in the mammal being immunized. Examples of such immunogenic proteins include but are not limited to keyhole limpet hemocyanin, serum albumin, bovine thyroglobulin, and soybean trypsin inhibitor. Examples of adjuvants which may be employed include Freund's complete adjuvant and MPL-TDM adjuvant (monophosphoryl Lipid A, synthetic trehalose dicorynomycolate). The immunization protocol may be selected by one skilled in the art without undue experimentation.

[119] The antibodies may, alternatively, be monoclonal antibodies. Monoclonal antibodies may be prepared using hybridoma methods, such as those described by Kohler and Milstein, *Nature*, 256:495 (1975). In a hybridoma method, a mouse, hamster, or other appropriate host animal, is typically immunized with an immunizing agent to elicit lymphocytes that produce or are capable of producing antibodies that will specifically bind to the immunizing agent. Alternatively, the lymphocytes may be immunized in vitro. The immunizing agent will typically include the CBF9 polypeptide or a peptide encoded by a

nucleic acid of Table 1 or a fragment thereof or a fusion protein thereof. Generally, either peripheral blood lymphocytes ("PBLs") are used if cells of human origin are desired, or spleen cells or lymph node cells are used if non-human mammalian sources are desired. The lymphocytes are then fused with an immortalized cell line using a suitable fusing agent, such 5 as polyethylene glycol, to form a hybridoma cell [Goding, Monoclonal Antibodies: Principles and Practice, Academic Press, (1986) pp. 59-103]. Immortalized cell lines are usually transformed mammalian cells, particularly myeloma cells of rodent, bovine and human origin. Usually, rat or mouse myeloma cell lines are employed. The hybridoma cells may be cultured in a suitable culture medium that preferably contains one or more substances that 10 inhibit the growth or survival of the unfused, immortalized cells. For example, if the parental cells lack the enzyme hypoxanthine guanine phosphoribosyl transferase (HGPRT or HPRT), the culture medium for the hybridomas typically will include hypoxanthine, aminopterin, and thymidine ("HAT medium"), which substances prevent the growth of HGPRT-deficient cells.

[120] In one embodiment, the antibodies are bispecific antibodies.

15 Bispecific antibodies are monoclonal, preferably human or humanized, antibodies that have binding specificities for at least two different antigens. In the present case, one of the binding specificities is for a colorectal cancer protein or a fragment thereof, the other one is for any other antigen, and preferably for a cell-surface protein or receptor or receptor subunit, preferably one that is tumor specific.

20 [121] In a preferred embodiment, the antibodies to colorectal cancer are capable of reducing or eliminating the biological function of colorectal cancer, as is described below. That is, the addition of anti-colorectal cancer antibodies (either polyclonal or preferably monoclonal) to colorectal cancer (or cells containing colorectal cancer) may reduce or eliminate the colorectal cancer activity. Generally, at least a 25% decrease in 25 activity is preferred, with at least about 50% being particularly preferred and about a 95-100% decrease being especially preferred.

[122] In a preferred embodiment the antibodies to the colorectal cancer proteins are humanized antibodies. Humanized forms of non-human (e.g., murine) antibodies are chimeric molecules of immunoglobulins, immunoglobulin chains or fragments thereof 30 (such as Fv, Fab, Fab', F(ab')2 or other antigen-binding subsequences of antibodies) which contain minimal sequence derived from non-human immunoglobulin. Humanized antibodies include human immunoglobulins (recipient antibody) in which residues form a complementary determining region (CDR) of the recipient are replaced by residues from a CDR of a non-human species (donor antibody) such as mouse, rat or rabbit having the desired

specificity, affinity and capacity. In some instances, Fv framework residues of the human immunoglobulin are replaced by corresponding non-human residues. Humanized antibodies may also comprise residues which are found neither in the recipient antibody nor in the imported CDR or framework sequences. In general, the humanized antibody will comprise substantially all of at least one, and typically two, variable domains, in which all or substantially all of the CDR regions correspond to those of a non-human immunoglobulin and all or substantially all of the FR regions are those of a human immunoglobulin consensus sequence. The humanized antibody optimally also will comprise at least a portion of an immunoglobulin constant region (Fc), typically that of a human immunoglobulin [Jones et al., Nature, 321:522-525 (1986); Riechmann et al., Nature, 332:323-329 (1988); and Presta, Curr. Op. Struct. Biol., 2:593-596 (1992)].

[123] Methods for humanizing non-human antibodies are well known in the art. Generally, a humanized antibody has one or more amino acid residues introduced into it from a source which is non-human. These non-human amino acid residues are often referred to as import residues, which are typically taken from an import variable domain. Humanization can be essentially performed following the method of Winter and co-workers [Jones et al., Nature, 321:522-525 (1986); Riechmann et al., Nature, 332:323-327 (1988); Verhoeyen et al., Science, 239:1534-1536 (1988)], by substituting rodent CDRs or CDR sequences for the corresponding sequences of a human antibody. Accordingly, such humanized antibodies are chimeric antibodies (U.S. Patent No. 4,816,567), wherein substantially less than an intact human variable domain has been substituted by the corresponding sequence from a non-human species. In practice, humanized antibodies are typically human antibodies in which some CDR residues and possibly some FR residues are substituted by residues from analogous sites in rodent antibodies.

[124] Human antibodies can also be produced using various techniques known in the art, including phage display libraries [Hoogenboom and Winter, J. Mol. Biol., 227:381 (1991); Marks et al., J. Mol. Biol., 222:581 (1991)]. The techniques of Cole et al. and Boerner et al. are also available for the preparation of human monoclonal antibodies (Cole et al., Monoclonal Antibodies and Cancer Therapy, Alan R. Liss, p. 77 (1985) and Boerner et al., J. Immunol., 147(1):86-95 (1991)]. Similarly, human antibodies can be made by introducing of human immunoglobulin loci into transgenic animals, e.g., mice in which the endogenous immunoglobulin genes have been partially or completely inactivated. Upon challenge, human antibody production is observed, which closely resembles that seen in humans in all respects, including gene rearrangement, assembly, and antibody repertoire.

This approach is described, for example, in U.S. Patent Nos. 5,545,807; 5,545,806; 5,569,825; 5,625,126; 5,633,425; 5,661,016, and in the following scientific publications: Marks et al., Bio/Technology 10, 779-783 (1992); Lonberg et al., Nature 368 856-859 (1994); Morrison, Nature 368, 812-13 (1994); Fishwild et al., Nature Biotechnology 14, 845-51 (1996); Neuberger, Nature Biotechnology 14, 826 (1996); Lonberg and Huszar, Intern. Rev. Immunol. 13 65-93 (1995).

[125] By immunotherapy is meant treatment of colorectal cancer with an antibody raised against colorectal cancer proteins. As used herein, immunotherapy can be passive or active. Passive immunotherapy as defined herein is the passive transfer of antibody to a recipient (patient). Active immunization is the induction of antibody and/or T-cell responses in a recipient (patient). Induction of an immune response is the result of providing the recipient with an antigen to which antibodies are raised. As appreciated by one of ordinary skill in the art, the antigen may be provided by injecting a polypeptide against which antibodies are desired to be raised into a recipient, or contacting the recipient with a nucleic acid capable of expressing the antigen and under conditions for expression of the antigen.

[126] In a preferred embodiment the colorectal cancer proteins against which antibodies are raised are secreted proteins as described above. Without being bound by theory, antibodies used for treatment, bind and prevent the secreted protein from binding to its receptor, thereby inactivating the secreted colorectal cancer protein.

[127] In another preferred embodiment, the colorectal cancer protein to which antibodies are raised is a transmembrane protein. Without being bound by theory, antibodies used for treatment, bind the extracellular domain of the colorectal cancer protein and prevent it from binding to other proteins, such as circulating ligands or cell-associated molecules. The antibody may cause down-regulation of the transmembrane colorectal cancer protein. As will be appreciated by one of ordinary skill in the art, the antibody may be a competitive, non-competitive or uncompetitive inhibitor of protein binding to the extracellular domain of the colorectal cancer protein. The antibody is also an antagonist of the colorectal cancer protein. Further, the antibody prevents activation of the transmembrane colorectal cancer protein. In one aspect, when the antibody prevents the binding of other molecules to the colorectal cancer protein, the antibody prevents growth of the cell. The antibody also sensitizes the cell to cytotoxic agents, including, but not limited to TNF- $\alpha$ , TNF- $\beta$ , IL-1, INF- $\gamma$  and IL-2, or chemotherapeutic agents including 5FU, vinblastine,

actinomycin D, cisplatin, methotrexate, and the like. In some instances the antibody belongs to a sub-type that activates serum complement when complexed with the transmembrane protein thereby mediating cytotoxicity. Thus, colorectal cancer is treated by administering to a patient antibodies directed against the transmembrane colorectal cancer protein.

5 [128] In another preferred embodiment, the antibody is conjugated to a therapeutic moiety. In one aspect the therapeutic moiety is a small molecule that modulates the activity of the colorectal cancer protein. In another aspect the therapeutic moiety modulates the activity of molecules associated with or in close proximity to the colorectal cancer protein. The therapeutic moiety may inhibit enzymatic activity such as protease or  
10 protein kinase activity associated with colorectal cancer .

[129] In a preferred embodiment, the therapeutic moiety may also be a cytotoxic agent. In this method, targeting the cytotoxic agent to tumor tissue or cells, results in a reduction in the number of afflicted cells, thereby reducing symptoms associated with colorectal cancer . Cytotoxic agents are numerous and varied and include, but are not limited  
15 to, cytotoxic drugs or toxins or active fragments of such toxins. Suitable toxins and their corresponding fragments include diphteria A chain, exotoxin A chain, ricin A chain, abrin A chain, curcin, crotin, phenomycin, enomycin and the like. Cytotoxic agents also include radiochemicals made by conjugating radioisotopes to antibodies raised against colorectal cancer proteins, or binding of a radionuclide to a chelating agent that has been covalently  
20 attached to the antibody. Targeting the therapeutic moiety to transmembrane colorectal cancer proteins not only serves to increase the local concentration of therapeutic moiety in the colorectal cancer afflicted area, but also serves to reduce deleterious side effects that may be associated with the therapeutic moiety.

[130] In another preferred embodiment, the colorectal cancer protein against  
25 which the antibodies are raised is an intracellular protein. In this case, the antibody may be conjugated to a protein which facilitates entry into the cell. In one case, the antibody enters the cell by endocytosis. In another embodiment, a nucleic acid encoding the antibody is administered to the individual or cell. Moreover, wherein the colorectal cancer protein can be targeted within a cell, i.e., the nucleus, an antibody thereto contains a signal for that target  
30 localization, i.e., a nuclear localization signal.

[131] The colorectal cancer antibodies of the invention specifically bind to colorectal cancer proteins. By "specifically bind" herein is meant that the antibodies bind to the protein with a binding constant in the range of at least  $10^{-4}$ -  $10^{-6} \text{ M}^{-1}$ , with a preferred range being  $10^{-7}$  -  $10^{-9} \text{ M}^{-1}$ .

[132] In a preferred embodiment, the colorectal cancer protein is purified or isolated after expression. Colorectal cancer proteins may be isolated or purified in a variety of ways known to those skilled in the art depending on what other components are present in the sample. Standard purification methods include electrophoretic, molecular, 5 immunological and chromatographic techniques, including ion exchange, hydrophobic, affinity, and reverse-phase HPLC chromatography, and chromatofocusing. For example, the colorectal cancer protein may be purified using a standard anti-colorectal cancer antibody column. Ultrafiltration and diafiltration techniques, in conjunction with protein concentration, are also useful. For general guidance in suitable purification techniques, see 10 Scopes, R., Protein Purification, Springer-Verlag, NY (1982). The degree of purification necessary will vary depending on the use of the colorectal cancer protein. In some instances no purification will be necessary.

[133] Once expressed and purified if necessary, the colorectal cancer proteins and nucleic acids are useful in a number of applications.

[134] In one aspect, the expression levels of genes are determined for different cellular states in the colorectal cancer phenotype; that is, the expression levels of genes in normal colon tissue and in colorectal cancer tissue (and in some cases, for varying severities of colorectal cancer that relate to prognosis, as outlined below) are evaluated to provide expression profiles. An expression profile of a particular cell state or point of development is essentially a "fingerprint" of the state; while two states may have any particular gene similarly expressed, the evaluation of a number of genes simultaneously allows the generation of a gene expression profile that is unique to the state of the cell. By comparing expression profiles of cells in different states, information regarding which genes are important (including both up- and down-regulation of genes) in each of these states is 20 obtained. Then, diagnosis may be done or confirmed: does tissue from a particular patient 25 have the gene expression profile of normal or colorectal cancer tissue.

[135] "Differential expression," or grammatical equivalents as used herein, refers to both qualitative as well as quantitative differences in the genes' temporal and/or cellular expression patterns within and among the cells. Thus, a differentially expressed gene 30 can qualitatively have its expression altered, including an activation or inactivation, in, for example, normal versus colorectal cancer tissue. That is, genes may be turned on or turned off in a particular state, relative to another state. As is apparent to the skilled artisan, any comparison of two or more states can be made. Such a qualitatively regulated gene will exhibit an expression pattern within a state or cell type which is detectable by standard

techniques in one such state or cell type, but is not detectable in both. Alternatively, the determination is quantitative in that expression is increased or decreased; that is, the expression of the gene is either upregulated, resulting in an increased amount of transcript, or downregulated, resulting in a decreased amount of transcript. The degree to which  
5 expression differs need only be large enough to quantify via standard characterization techniques as outlined below, such as by use of Affymetrix GeneChip™ expression arrays, Lockhart, Nature Biotechnology, 14:1675-1680 (1996), hereby expressly incorporated by reference. Other techniques include, but are not limited to, quantitative reverse transcriptase PCR, Northern analysis and RNase protection. As outlined above, preferably the change in  
10 expression (i.e. upregulation or downregulation) is at least about 50%, more preferably at least about 100%, more preferably at least about 150%, more preferably, at least about 200%, with from 300 to at least 1000% being especially preferred.

[136] As will be appreciated by those in the art, this may be done by evaluation at either the gene transcript, or the protein level; that is, the amount of gene  
15 expression may be monitored using nucleic acid probes to the DNA or RNA equivalent of the gene transcript, and the quantification of gene expression levels, or, alternatively, the final gene product itself (protein) can be monitored, for example through the use of antibodies to the colorectal cancer protein and standard immunoassays (ELISAs, etc.) or other techniques, including mass spectroscopy assays, 2D gel electrophoresis assays, etc. Thus, the proteins  
20 corresponding to colorectal cancer genes, i.e. those identified as being important in a colorectal cancer phenotype, can be evaluated in a colorectal cancer diagnostic test.

[137] In a preferred embodiment, gene expression monitoring is done and a number of genes, i.e. an expression profile, is monitored simultaneously, although multiple protein expression monitoring can be done as well. Similarly, these assays may be done on  
25 an individual basis as well.

[138] In this embodiment, the colorectal cancer nucleic acid probes are attached to biochips as outlined herein for the detection and quantification of colorectal cancer sequences in a particular cell. The assays are further described below in the example.

[139] In a preferred embodiment nucleic acids encoding the colorectal  
30 cancer protein are detected. Although DNA or RNA encoding the colorectal cancer protein may be detected, of particular interest are methods wherein the mRNA encoding a colorectal cancer protein is detected. The presence of mRNA in a sample is an indication that the colorectal cancer gene has been transcribed to form the mRNA, and suggests that the protein

is expressed. Probes to detect the mRNA can be any nucleotide/deoxynucleotide probe that is complementary to and base pairs with the mRNA and includes but is not limited to oligonucleotides, cDNA or RNA. Probes also should contain a detectable label, as defined herein. In one method the mRNA is detected after immobilizing the nucleic acid to be  
5 examined on a solid support such as nylon membranes and hybridizing the probe with the sample. Following washing to remove the non-specifically bound probe, the label is detected. In another method detection of the mRNA is performed *in situ*. In this method permeabilized cells or tissue samples are contacted with a detectably labeled nucleic acid probe for sufficient time to allow the probe to hybridize with the target mRNA. Following  
10 washing to remove the non-specifically bound probe, the label is detected. For example a digoxygenin labeled riboprobe (RNA probe) that is complementary to the mRNA encoding a colorectal cancer protein is detected by binding the digoxygenin with an anti-digoxygenin secondary antibody and developed with nitro blue tetrazolium and 5-bromo-4-chloro-3-indoyl phosphate.

15 [140] In a preferred embodiment, any of the three classes of proteins as described herein (secreted, transmembrane or intracellular proteins) are used in diagnostic assays. The colorectal cancer proteins, antibodies, nucleic acids, modified proteins and cells containing colorectal cancer sequences are used in diagnostic assays. This can be done on an individual gene or corresponding polypeptide level. In a preferred embodiment, the  
20 expression profiles are used, preferably in conjunction with high throughput screening techniques to allow monitoring for expression profile genes and/or corresponding polypeptides.

[141] As described and defined herein, colorectal cancer proteins, including intracellular, transmembrane or secreted proteins, find use as markers of colorectal cancer .  
25 Detection of these proteins in putative colorectal cancer tissue or patients allows for a determination or diagnosis of colorectal cancer . Numerous methods known to those of ordinary skill in the art find use in detecting colorectal cancer . In one embodiment, antibodies are used to detect colorectal cancer proteins. A preferred method separates proteins from a sample or patient by electrophoresis on a gel (typically a denaturing and  
30 reducing protein gel, but may be any other type of gel including isoelectric focusing gels and the like). Following separation of proteins, the colorectal cancer protein is detected by immunoblotting with antibodies raised against the colorectal cancer protein. Methods of immunoblotting are well known to those of ordinary skill in the art.

[142] In another preferred method, antibodies to the colorectal cancer protein find use in in situ imaging techniques. In this method cells are contacted with from one to many antibodies to the colorectal cancer protein(s). Following washing to remove non-specific antibody binding, the presence of the antibody or antibodies is detected. In one embodiment the antibody is detected by incubating with a secondary antibody that contains a detectable label. In another method the primary antibody to the colorectal cancer protein(s) contains a detectable label. In another preferred embodiment each one of multiple primary antibodies contains a distinct and detectable label. This method finds particular use in simultaneous screening for a plurality of colorectal cancer proteins. As will be appreciated by one of ordinary skill in the art, numerous other histological imaging techniques are useful in the invention.

[143] In a preferred embodiment the label is detected in a fluorometer which has the ability to detect and distinguish emissions of different wavelengths. In addition, a fluorescence activated cell sorter (FACS) can be used in the method.

[144] In another preferred embodiment, antibodies find use in diagnosing colorectal cancer from blood samples. As previously described, certain colorectal cancer proteins are secreted/circulating molecules. Blood samples, therefore, are useful as samples to be probed or tested for the presence of secreted colorectal cancer proteins. Antibodies can be used to detect the colorectal cancer by any of the previously described immunoassay techniques including ELISA, immunoblotting (Western blotting), immunoprecipitation, BIACORE technology and the like, as will be appreciated by one of ordinary skill in the art.

[145] In a preferred embodiment, in situ hybridization of labeled colorectal cancer nucleic acid probes to tissue arrays is done. For example, arrays of tissue samples, including colorectal cancer tissue and/or normal tissue, are made. In situ hybridization as is known in the art can then be done.

[146] It is understood that when comparing the fingerprints between an individual and a standard, the skilled artisan can make a diagnosis as well as a prognosis. It is further understood that the genes which indicate the diagnosis may differ from those which indicate the prognosis.

[147] In a preferred embodiment, the colorectal cancer proteins, antibodies, nucleic acids, modified proteins and cells containing colorectal cancer sequences are used in prognosis assays. As above, gene expression profiles can be generated that correlate to colorectal cancer severity, in terms of long term prognosis. Again, this may be done on either a protein or gene level, with the use of genes being preferred. As above, the colorectal

cancer probes are attached to biochips for the detection and quantification of colorectal cancer sequences in a tissue or patient. The assays proceed as outlined for diagnosis.

[148] In a preferred embodiment, any of the three classes of proteins as described herein are used in drug screening assays. The colorectal cancer proteins, antibodies, nucleic acids, modified proteins and cells containing colorectal cancer sequences are used in drug screening assays or by evaluating the effect of drug candidates on a "gene expression profile" or expression profile of polypeptides. In a preferred embodiment, the expression profiles are used, preferably in conjunction with high throughput screening techniques to allow monitoring for expression profile genes after treatment with a candidate agent, Zlokarnik, et al., Science 279, 84-8 (1998), Heid, 1996 #69.

[149] In a preferred embodiment, the colorectal cancer proteins, antibodies, nucleic acids, modified proteins and cells containing the native or modified colorectal cancer proteins are used in screening assays. That is, the present invention provides novel methods for screening for compositions which modulate the colorectal cancer phenotype. As above, this can be done on an individual gene level or by evaluating the effect of drug candidates on a "gene expression profile". In a preferred embodiment, the expression profiles are used, preferably in conjunction with high throughput screening techniques to allow monitoring for expression profile genes after treatment with a candidate agent, see Zlokarnik, *supra*.

Having identified the differentially expressed genes herein, a variety of assays may be executed. In a preferred embodiment, assays may be run on an individual gene or protein level. That is, having identified a particular gene as up regulated in colorectal cancer, candidate bioactive agents may be screened to modulate this gene's response; preferably to down regulate the gene, although in some circumstances to up regulate the gene.

[150] The phrase "functional effects" in the context of assays for testing compounds that modulate activity of a colorectal cancer protein or colorectal cancer nucleic acid includes the determination of a parameter that is indirectly or directly under the influence of a colorectal cancer protein or nucleic acid, *e.g.*, a physical (direct), or phenotypic or chemical effect (indirect), such as the ability to increase or decrease cellular proliferation. It includes cell cycle arrest, the ability of cells to proliferate, and other characteristics of proliferating cells. "Functional effects" include *in vitro*, *in vivo*, and *ex vivo* activities.

[151] By "determining the functional effect" is meant assaying for a compound that increases or decreases a parameter that is indirectly or directly under the influence of a colorectal cancer protein or nucleic acid, *e.g.*, physical, phenotypic and chemical effects. Such functional effects can be measured by any means known to those

skilled in the art, *e.g.*, physical effects such as changes in spectroscopic characteristics (*e.g.*, fluorescence, absorbance, refractive index); hydrodynamic (*e.g.*, shape); chromatographic; or solubility properties for the protein; measuring ligand binding activity or binding assays, *e.g.* binding to antibodies; measuring changes in ligand binding activity; and chemical or  
5 phenotypic effects such as measuring inducible markers or transcriptional activation of the protein; measuring cellular proliferation; measuring cell surface marker expression; measurement of changes in protein levels for colorectal cancer-associated sequences; measurement of RNA stability; phosphorylation or dephosphorylation; signal transduction, *e.g.*, receptor-ligand interactions, second messenger concentrations (*e.g.*, cAMP, IP<sub>3</sub>, or  
10 intracellular Ca<sup>2+</sup>); identification of downstream or reporter gene expression (CAT, luciferase, β-gal, GFP and the like), *e.g.*, via chemiluminescence, fluorescence, colorimetric reactions, antibody binding, and inducible markers.

[152] “Inhibitors”, “activators”, and “modulators” of colorectal cancer polynucleotide and polypeptide sequences are used to refer to activating, inhibitory, or  
15 modulating molecules identified using *in vitro* and *in vivo* assays of colorectal cancer polynucleotide and polypeptide sequences. Inhibitors are compounds that, *e.g.*, bind to, partially or totally block activity, decrease, prevent, delay activation, inactivate, desensitize, or down regulate the activity or expression of colorectal cancer proteins or nucleic acids, *e.g.*, antagonists. “Activators” are compounds that increase, open, activate, facilitate, enhance  
20 activation, sensitize, agonize, or up regulate colorectal cancer protein or nucleic acid activity. Inhibitors, activators, or modulators also include genetically modified versions of colorectal cancer proteins, *e.g.*, versions with altered activity, as well as naturally occurring and synthetic ligands, antagonists, agonists, antibodies, antisense molecules, peptides, ribozymes, small chemical molecules and the like. Such assays for inhibitors and activators include, *e.g.*,  
25 expressing colorectal cancer protein *in vitro*, in cells, or cell membranes, applying putative modulator compounds, and then determining the functional effects on activity, as described above.

[153] Samples or assays comprising colorectal cancer proteins or colorectal cancer nucleic acids that are treated with a potential activator, inhibitor, or modulator are  
30 compared to control samples without the inhibitor, activator, or modulator to examine the extent of inhibition. Control samples (untreated with inhibitors) are assigned a relative activity value of 100%. Inhibition of colorectal cancer is achieved when the activity value relative to the control is about 80%, preferably 50%, more preferably 25-0%. Activation of

colorectal cancer is achieved when the activity value relative to the control (untreated with activators) is 110%, more preferably 150%, more preferably 200-500% (i.e., two to five fold higher relative to the control), more preferably 1000-3000% higher.

[154] As will be appreciated by those in the art, this may be done by evaluation at either the gene or the protein level; that is, the amount of gene expression may be monitored using nucleic acid probes and the quantification of gene expression levels, or, alternatively, the gene product itself can be monitored, for example through the use of antibodies to the colorectal cancer protein and standard immunoassays.

[155] In a preferred embodiment, gene expression monitoring is done and a number of genes, i.e. an expression profile, is monitored simultaneously, although multiple protein expression monitoring can be done as well.

[156] In this embodiment, the colorectal cancer nucleic acid probes are attached to biochips as outlined herein for the detection and quantification of colorectal cancer sequences in a particular cell. The assays are further described below.

[157] Generally, in a preferred embodiment, a candidate bioactive agent is added to the cells prior to analysis. Moreover, screens are provided to identify a candidate bioactive agent which modulates colorectal cancer, modulates colorectal cancer proteins, binds to a colorectal cancer protein, or interferes between the binding of a colorectal cancer protein and an antibody.

[158] The term "candidate bioactive agent" or "test compound" or "drug candidate" or "modulator" or grammatical equivalents as used herein describes any molecule, either naturally occurring or synthetic, e.g., protein, oligopeptide (e.g., from about 5 to about 25 amino acids in length, preferably from about 10 to 20 or 12 to 18 amino acids in length, preferably 12, 15, or 18 amino acids in length), small organic molecule, polysaccharide, lipid, fatty acid, polynucleotide, oligonucleotide, etc., to be tested for the capacity to directly or indirectly modulate colorectal cancer sequences, including both nucleic acid and protein sequences. The test compound can be in the form of a library of test compounds, such as a combinatorial or randomized library that provides a sufficient range of diversity. Test compounds are optionally linked to a fusion partner, e.g., targeting compounds, rescue compounds, dimerization compounds, stabilizing compounds, addressable compounds, and other functional moieties. Conventionally, new chemical entities with useful properties are generated by identifying a test compound (called a "lead compound") with some desirable property or activity, e.g., inhibiting activity, creating variants of the lead compound, and

evaluating the property and activity of those variant compounds. Often, high throughput screening (HTS) methods are employed for such an analysis.

[159] In preferred embodiments, the bioactive agents modulate the expression profiles, or expression profile nucleic acids or proteins provided herein. In a particularly preferred embodiment, the candidate agent suppresses a colorectal cancer phenotype, for example to a normal colon tissue fingerprint. Similarly, the candidate agent preferably suppresses a severe colorectal cancer phenotype. Generally a plurality of assay mixtures are run in parallel with different agent concentrations to obtain a differential response to the various concentrations. Typically, one of these concentrations serves as a negative control, i.e., at zero concentration or below the level of detection.

[160] In one aspect, a candidate agent will neutralize the effect of a colorectal cancer protein. By "neutralize" is meant that activity of a protein is either inhibited or counter acted against so as to have substantially no effect on a cell.

[161] Candidate agents encompass numerous chemical classes, though typically they are organic molecules, preferably small organic compounds having a molecular weight of more than 100 and less than about 2,500 daltons. Preferred small molecules are less than 2000, or less than 1500 or less than 1000 or less than 500 D. Candidate agents comprise functional groups necessary for structural interaction with proteins, particularly hydrogen bonding, and typically include at least an amine, carbonyl, hydroxyl or carboxyl group, preferably at least two of the functional chemical groups. The candidate agents often comprise cyclical carbon or heterocyclic structures and/or aromatic or polyaromatic structures substituted with one or more of the above functional groups. Candidate agents are also found among biomolecules including peptides, saccharides, fatty acids, steroids, purines, pyrimidines, derivatives, structural analogs or combinations thereof. Particularly preferred are peptides.

[162] Candidate agents are obtained from a wide variety of sources including libraries of synthetic or natural compounds. For example, numerous means are available for random and directed synthesis of a wide variety of organic compounds and biomolecules, including expression of randomized oligonucleotides. Alternatively, libraries of natural compounds in the form of bacterial, fungal, plant and animal extracts are available or readily produced. Additionally, natural or synthetically produced libraries and compounds are readily modified through conventional chemical, physical and biochemical means. Known pharmacological agents may be subjected to directed or random chemical modifications, such as acylation, alkylation, esterification, amidification to produce structural analogs.

[163] In a preferred embodiment, the candidate bioactive agents are proteins. By "protein" herein is meant at least two covalently attached amino acids, which includes proteins, polypeptides, oligopeptides and peptides. The protein may be made up of naturally occurring amino acids and peptide bonds, or synthetic peptidomimetic structures.

5 Thus "amino acid", or "peptide residue", as used herein means both naturally occurring and synthetic amino acids. For example, homo-phenylalanine, citrulline and noreleucine are considered amino acids for the purposes of the invention. "Amino acid" also includes imino acid residues such as proline and hydroxyproline. The side chains may be in either the (R) or the (S) configuration. In the preferred embodiment, the amino acids are in the (S) or L-  
10 configuration. If non-naturally occurring side chains are used, non-amino acid substituents may be used, for example to prevent or retard in vivo degradations.

[164] In a preferred embodiment, the candidate bioactive agents are naturally occurring proteins or fragments of naturally occurring proteins. Thus, for example, cellular extracts containing proteins, or random or directed digests of proteinaceous cellular extracts,  
15 may be used. In this way libraries of prokaryotic and eukaryotic proteins may be made for screening in the methods of the invention. Particularly preferred in this embodiment are libraries of bacterial, fungal, viral, and mammalian proteins, with the latter being preferred, and human proteins being especially preferred.

[165] In a preferred embodiment, the candidate bioactive agents are peptides  
20 of from about 5 to about 30 amino acids, with from about 5 to about 20 amino acids being preferred, and from about 7 to about 15 being particularly preferred. The peptides may be digests of naturally occurring proteins as is outlined above, random peptides, or "biased" random peptides. By "randomized" or grammatical equivalents herein is meant that each nucleic acid and peptide consists of essentially random nucleotides and amino acids,  
25 respectively. Since generally these random peptides (or nucleic acids, discussed below) are chemically synthesized, they may incorporate any nucleotide or amino acid at any position. The synthetic process can be designed to generate randomized proteins or nucleic acids, to allow the formation of all or most of the possible combinations over the length of the sequence, thus forming a library of randomized candidate bioactive proteinaceous agents.

30 [166] In one embodiment, the library is fully randomized, with no sequence preferences or constants at any position. In a preferred embodiment, the library is biased. That is, some positions within the sequence are either held constant, or are selected from a limited number of possibilities. For example, in a preferred embodiment, the nucleotides or amino acid residues are randomized within a defined class, for example, of hydrophobic

amino acids, hydrophilic residues, sterically biased (either small or large) residues, towards the creation of nucleic acid binding domains, the creation of cysteines, for cross-linking, prolines for SH-3 domains, serines, threonines, tyrosines or histidines for phosphorylation sites, etc., or to purines, etc.

5 [167] In a preferred embodiment, the candidate bioactive agents are nucleic acids, as defined above.

10 [168] As described above generally for proteins, nucleic acid candidate bioactive agents may be naturally occurring nucleic acids, random nucleic acids, or "biased" random nucleic acids. For example, digests of prokaryotic or eukaryotic genomes may be used as is outlined above for proteins.

[169] In a preferred embodiment, the candidate bioactive agents are organic chemical moieties, a wide variety of which are available in the literature.

15 [170] "Antibody" refers to a polypeptide comprising a framework region from an immunoglobulin gene or fragments thereof that specifically binds and recognizes an antigen. The recognized immunoglobulin genes include the kappa, lambda, alpha, gamma, delta, epsilon, and mu constant region genes, as well as the myriad immunoglobulin variable region genes. Light chains are classified as either kappa or lambda. Heavy chains are classified as gamma, mu, alpha, delta, or epsilon, which in turn define the immunoglobulin classes, IgG, IgM, IgA, IgD and IgE, respectively. Typically, the antigen-binding region of 20 an antibody will be most critical in specificity and affinity of binding.

25 [171] An exemplary immunoglobulin (antibody) structural unit comprises a tetramer. Each tetramer is composed of two identical pairs of polypeptide chains, each pair having one "light" (about 25 kD) and one "heavy" chain (about 50-70 kD). The N-terminus of each chain defines a variable region of about 100 to 110 or more amino acids primarily responsible for antigen recognition. The terms variable light chain ( $V_L$ ) and variable heavy chain ( $V_H$ ) refer to these light and heavy chains respectively.

30 [172] Antibodies exist, e.g., as intact immunoglobulins or as a number of well-characterized fragments produced by digestion with various peptidases. Thus, for example, pepsin digests an antibody below the disulfide linkages in the hinge region to produce  $F(ab')_2$ , a dimer of Fab which itself is a light chain joined to  $V_H$ - $C_H1$  by a disulfide bond. The  $F(ab')_2$  may be reduced under mild conditions to break the disulfide linkage in the hinge region, thereby converting the  $F(ab')_2$  dimer into an Fab' monomer. The Fab' monomer is essentially Fab with part of the hinge region (*see Fundamental Immunology* (Paul ed., 3d ed. 1993). While various antibody fragments are defined in terms of the

digestion of an intact antibody, one of skill will appreciate that such fragments may be synthesized *de novo* either chemically or by using recombinant DNA methodology. Thus, the term antibody, as used herein, also includes antibody fragments either produced by the modification of whole antibodies, or those synthesized *de novo* using recombinant DNA methodologies (e.g., single chain Fv) or those identified using phage display libraries (see, e.g., McCafferty *et al.*, *Nature* 348:552-554 (1990))

[173] For preparation of antibodies, e.g., recombinant, monoclonal, or polyclonal antibodies, many technique known in the art can be used (see, e.g., Kohler & Milstein, *Nature* 256:495-497 (1975); Kozbor *et al.*, *Immunology Today* 4: 72 (1983); Cole *et al.*, pp. 77-96 in *Monoclonal Antibodies and Cancer Therapy*, Alan R. Liss, Inc. (1985); Coligan, *Current Protocols in Immunology* (1991); Harlow & Lane, *Antibodies, A Laboratory Manual* (1988); and Goding, *Monoclonal Antibodies: Principles and Practice* (2d ed. 1986)). The genes encoding the heavy and light chains of an antibody of interest can be cloned from a cell, e.g., the genes encoding a monoclonal antibody can be cloned from a hybridoma and used to produce a recombinant monoclonal antibody. Gene libraries encoding heavy and light chaims of monoclonal antibodies can also be made from hybridoma or plasma cells. Random combinations of the heavy and light chain gene products generate a large pool of antibodies with different antigenic specificity (see, e.g., Kuby, *Immunology* (3<sup>rd</sup> ed. 1997)). Techniques for the production of single chain antibodies or recombinant antibodies (U.S. Patent 4,946,778, U.S. Patent No. 4,816,567) can be adapted to produce antibodies to polypeptides of this invention. Also, transgenic mice, or other organisms such as other mammals, may be used to express humanized or human antibodies (see, e.g., U.S. Patent Nos. 5,545,807; 5,545,806; 5,569,825; 5,625,126; 5,633,425; 5,661,016, Marks *et al.*, *Bio/Technology* 10:779-783 (1992); Lonberg *et al.*, *Nature* 368:856-859 (1994); Morrison, *Nature* 368:812-13 (1994); Fishwild *et al.*, *Nature Biotechnology* 14:845-51 (1996); Neuberger, *Nature Biotechnology* 14:826 (1996); and Lonberg & Huszar, *Intern. Rev. Immunol.* 13:65-93 (1995)). Alternatively, phage display technology can be used to identify antibodies and heteromeric Fab fragments that specifically bind to selected antigens (see, e.g., McCafferty *et al.*, *Nature* 348:552-554 (1990); Marks *et al.*, *Biotechnology* 10:779-783 (1992)). Antibodies can also be made bispecific, i.e., able to recognize two different antigens (see, e.g., WO 93/08829, Traunecker *et al.*, *EMBO J.* 10:3655-3659 (1991); and Suresh *et al.*, *Methods in Enzymology* 121:210 (1986)). Antibodies can also be heteroconjugates, e.g., two covalently joined antibodies, or immunotoxins (see, e.g., U.S. Patent No. 4,676,980 , WO 91/00360; WO 92/200373; and EP 03089).

[174] Methods for humanizing or primatizing non-human antibodies are well known in the art. Generally, a humanized antibody has one or more amino acid residues introduced into it from a source which is non-human. These non-human amino acid residues are often referred to as import residues, which are typically taken from an import variable domain. Humanization can be essentially performed following the method of Winter and co-workers (see, e.g., Jones *et al.*, *Nature* 321:522-525 (1986); Riechmann *et al.*, *Nature* 332:323-327 (1988); Verhoeyen *et al.*, *Science* 239:1534-1536 (1988) and Presta, *Curr. Op. Struct. Biol.* 2:593-596 (1992)), by substituting rodent CDRs or CDR sequences for the corresponding sequences of a human antibody. Accordingly, such humanized antibodies are chimeric antibodies (U.S. Patent No. 4,816,567), wherein substantially less than an intact human variable domain has been substituted by the corresponding sequence from a non-human species. In practice, humanized antibodies are typically human antibodies in which some CDR residues and possibly some FR residues are substituted by residues from analogous sites in rodent antibodies.

[175] A “chimeric antibody” is an antibody molecule in which (a) the constant region, or a portion thereof, is altered, replaced or exchanged so that the antigen binding site (variable region) is linked to a constant region of a different or altered class, effector function and/or species, or an entirely different molecule which confers new properties to the chimeric antibody, e.g., an enzyme, toxin, hormone, growth factor, drug, etc.; or (b) the variable region, or a portion thereof, is altered, replaced or exchanged with a variable region having a different or altered antigen specificity.

[176] In one embodiment, the antibody is conjugated to an “effector” moiety. The effector moiety can be any number of molecules, including labeling moieties such as radioactive labels or fluorescent labels, or can be a therapeutic moiety. In one aspect the antibody modulates the activity of the protein.

[177] After the candidate agent has been added and the cells allowed to incubate for some period of time, the sample containing the target sequences to be analyzed is added to the biochip. If required, the target sequence is prepared using known techniques. For example, the sample may be treated to lyse the cells, using known lysis buffers, electroporation, etc., with purification and/or amplification such as PCR occurring as needed, as will be appreciated by those in the art. For example, an in vitro transcription with labels covalently attached to the nucleosides is done. Generally, the nucleic acids are labeled with biotin-FITC or PE, or with cy3 or cy5.

[178] In a preferred embodiment, the target sequence is labeled with, for example, a fluorescent, a chemiluminescent, a chemical, or a radioactive signal, to provide a means of detecting the target sequence's specific binding to a probe. The label also can be an enzyme, such as, alkaline phosphatase or horseradish peroxidase, which when provided with an appropriate substrate produces a product that can be detected. Alternatively, the label can be a labeled compound or small molecule, such as an enzyme inhibitor, that binds but is not catalyzed or altered by the enzyme. The label also can be a moiety or compound, such as, an epitope tag or biotin which specifically binds to streptavidin. For the example of biotin, the streptavidin is labeled as described above, thereby, providing a detectable signal for the bound target sequence. As known in the art, unbound labeled streptavidin is removed prior to analysis.

[179] As will be appreciated by those in the art, these assays can be direct hybridization assays or can comprise "sandwich assays", which include the use of multiple probes, as is generally outlined in U.S. Patent Nos. 5,681,702, 5,597,909, 5,545,730, 15 5,594,117, 5,591,584, 5,571,670, 5,580,731, 5,571,670, 5,591,584, 5,624,802, 5,635,352, 5,594,118, 5,359,100, 5,124,246 and 5,681,697, all of which are hereby incorporated by reference. In this embodiment, in general, the target nucleic acid is prepared as outlined above, and then added to the biochip comprising a plurality of nucleic acid probes, under conditions that allow the formation of a hybridization complex.

[180] A variety of hybridization conditions may be used in the present invention, including high, moderate and low stringency conditions as outlined above. The assays are generally run under stringency conditions which allows formation of the label probe hybridization complex only in the presence of target. Stringency can be controlled by altering a step parameter that is a thermodynamic variable, including, but not limited to, 25 temperature, formamide concentration, salt concentration, chaotropic salt concentration pH, organic solvent concentration, etc.

[181] These parameters may also be used to control non-specific binding, as is generally outlined in U.S. Patent No. 5,681,697. Thus it may be desirable to perform certain steps at higher stringency conditions to reduce non-specific binding.

[182] The reactions outlined herein may be accomplished in a variety of ways, as will be appreciated by those in the art. Components of the reaction may be added simultaneously, or sequentially, in any order, with preferred embodiments outlined below. In addition, the reaction may include a variety of other reagents may be included in the assays. These include reagents like salts, buffers, neutral proteins, e.g. albumin, detergents, etc which

may be used to facilitate optimal hybridization and detection, and/or reduce non-specific or background interactions. Also reagents that otherwise improve the efficiency of the assay, such as protease inhibitors, nuclease inhibitors, anti-microbial agents, etc., may be used, depending on the sample preparation methods and purity of the target.

5 [183] Once the assay is run, the data is analyzed to determine the expression levels, and changes in expression levels as between states, of individual genes, forming a gene expression profile.

10 [184] The screens are done to identify drugs or bioactive agents that modulate the colorectal cancer phenotype. Specifically, there are several types of screens that can be run. A preferred embodiment is in the screening of candidate agents that can induce or suppress a particular expression profile, thus preferably generating the associated phenotype. That is, candidate agents that can mimic or produce an expression profile in colorectal cancer similar to the expression profile of normal colon tissue is expected to result in a suppression of the colorectal cancer phenotype. Thus, in this embodiment, mimicking an 15 expression profile, or changing one profile to another, is the goal.

20 [185] In a preferred embodiment, as for the diagnosis and prognosis applications, having identified the differentially expressed genes important in any one state, screens can be run to alter the expression of the genes individually. That is, screening for modulation of regulation of expression of a single gene can be done; that is, rather than try to mimic all or part of an expression profile, screening for regulation of individual genes can be done. Thus, for example, particularly in the case of target genes whose presence or absence 25 is unique between two states, screening is done for modulators of the target gene expression.

25 [186] In a preferred embodiment, screening is done to alter the biological function of the expression product of the differentially expressed gene. Again, having identified the importance of a gene in a particular state, screening for agents that bind and/or modulate the biological activity of the gene product can be run as is more fully outlined below.

30 [187] Thus, screening of candidate agents that modulate the colorectal cancer phenotype either at the gene expression level or the protein level can be done.

[188] In addition screens can be done for novel genes that are induced in response to a candidate agent. After identifying a candidate agent based upon its ability to suppress a colorectal cancer expression pattern leading to a normal expression pattern, or modulate a single colorectal cancer gene expression profile so as to mimic the expression of the gene from normal tissue, a screen as described above can be performed to identify genes

that are specifically modulated in response to the agent. Comparing expression profiles between normal tissue and agent treated colorectal cancer tissue reveals genes that are not expressed in normal tissue or colorectal cancer tissue, but are expressed in agent treated tissue. These agent specific sequences can be identified and used by any of the methods 5 described herein for colorectal cancer genes or proteins. In particular these sequences and the proteins they encode find use in marking or identifying agent treated cells. In addition, antibodies can be raised against the agent induced proteins and used to target novel therapeutics to the treated colorectal cancer tissue sample.

[189] Thus, in one embodiment, a candidate agent is administered to a 10 population of colorectal cancer cells, that thus has an associated colorectal cancer expression profile. By "administration" or "contacting" herein is meant that the candidate agent is added to the cells in such a manner as to allow the agent to act upon the cell, whether by uptake and intracellular action, or by action at the cell surface. In some embodiments, nucleic acid encoding a proteinaceous candidate agent (i.e. a peptide) may be put into a viral 15 construct such as a retroviral construct and added to the cell, such that expression of the peptide agent is accomplished; see PCT US97/01019, hereby expressly incorporated by reference.

[190] Once the candidate agent has been administered to the cells, the cells 20 can be washed if desired and are allowed to incubate under preferably physiological conditions for some period of time. The cells are then harvested and a new gene expression profile is generated, as outlined herein.

[191] Thus, for example, colorectal cancer tissue may be screened for 25 agents that reduce or suppress the colorectal cancer phenotype. A change in at least one gene of the expression profile indicates that the agent has an effect on colorectal cancer activity. By defining such a signature for the colorectal cancer phenotype, screens for new drugs that alter the phenotype can be devised. With this approach, the drug target need not be known and need not be represented in the original expression screening platform, nor does the level of transcript for the target protein need to change.

[192] In a preferred embodiment, as outlined above, screens may be done on 30 individual genes and gene products (proteins). That is, having identified a particular differentially expressed gene as important in a particular state, screening of modulators of either the expression of the gene or the gene product itself can be done. The gene products of differentially expressed genes are sometimes referred to herein as "colorectal cancer modulator proteins". The colorectal cancer modulator protein may be a fragment, or

alternatively, be the full length protein to a fragment shown herein. Preferably, the colorectal cancer modulator protein is a fragment of approximately 14 to 24 amino acids long. More preferably the fragment is a soluble fragment.

[193] In a preferred embodiment, the fragment is charged and from the c-terminus. In one embodiment, the c-terminus of the fragment is kept as a free acid and the n-terminus is a free amine to aid in coupling, i.e., to cysteine. In another embodiment, the fragment is an internal peptide overlapping hydrophilic stretch the protein. In a preferred embodiment, the termini is blocked. In another preferred embodiment, the fragment is a novel fragment from the N-terminal. In one embodiment, the fragment excludes sequence outside of the N-terminal, in another embodiment, the fragment includes at least a portion of the N-terminal. "N-terminal" is used interchangeably herein with "N-terminus" which is further described above.

[194] In one embodiment the colorectal cancer proteins are conjugated to an immunogenic agent as discussed herein. In one embodiment the colorectal cancer protein is conjugated to BSA.

[195] Thus, in a preferred embodiment, screening for modulators of expression of specific genes can be done. This will be done as outlined above, but in general the expression of only one or a few genes are evaluated.

[196] In a preferred embodiment, screens are designed to first find candidate agents that can bind to differentially expressed proteins, and then these agents may be used in assays that evaluate the ability of the candidate agent to modulate differentially expressed activity. Thus, as will be appreciated by those in the art, there are a number of different assays which may be run; binding assays and activity assays.

[197] In a preferred embodiment, binding assays are done. In general, purified or isolated gene product is used; that is, the gene products of one or more differentially expressed nucleic acids are made. In general, this is done as is known in the art. For example, antibodies are generated to the protein gene products, and standard immunoassays are run to determine the amount of protein present. Alternatively, cells comprising the colorectal cancer proteins can be used in the assays.

[198] Thus, in a preferred embodiment, the methods comprise combining a colorectal cancer protein and a candidate bioactive agent, and determining the binding of the candidate agent to the colorectal cancer protein. Preferred embodiments utilize the human colorectal cancer protein, although other mammalian proteins may also be used, for example

for the development of animal models of human disease. In some embodiments, as outlined herein, variant or derivative colorectal cancer proteins may be used.

[199] Generally, in a preferred embodiment of the methods herein, the colorectal cancer protein or the candidate agent is non-diffusably bound to an insoluble support having isolated sample receiving areas (e.g. a microtiter plate, an array, etc.). The insoluble supports may be made of any composition to which the compositions can be bound, is readily separated from soluble material, and is otherwise compatible with the overall method of screening. The surface of such supports may be solid or porous and of any convenient shape. Examples of suitable insoluble supports include microtiter plates, arrays, membranes and beads. These are typically made of glass, plastic (e.g., polystyrene), polysaccharides, nylon or nitrocellulose, teflon, etc. Microtiter plates and arrays are especially convenient because a large number of assays can be carried out simultaneously, using small amounts of reagents and samples. The particular manner of binding of the composition is not crucial so long as it is compatible with the reagents and overall methods of the invention, maintains the activity of the composition and is nondiffusible. Preferred methods of binding include the use of antibodies (which do not sterically block either the ligand binding site or activation sequence when the protein is bound to the support), direct binding to "sticky" or ionic supports, chemical crosslinking, the synthesis of the protein or agent on the surface, etc. Following binding of the protein or agent, excess unbound material is removed by washing. The sample receiving areas may then be blocked through incubation with bovine serum albumin (BSA), casein or other innocuous protein or other moiety.

[200] In a preferred embodiment, the colorectal cancer protein is bound to the support, and a candidate bioactive agent is added to the assay. Alternatively, the candidate agent is bound to the support and the colorectal cancer protein is added. Novel binding agents include specific antibodies, non-natural binding agents identified in screens of chemical libraries, peptide analogs, etc. Of particular interest are screening assays for agents that have a low toxicity for human cells. A wide variety of assays may be used for this purpose, including labeled in vitro protein-protein binding assays, electrophoretic mobility shift assays, immunoassays for protein binding, functional assays (phosphorylation assays, etc.) and the like.

[201] The determination of the binding of the candidate bioactive agent to the colorectal cancer protein may be done in a number of ways. In a preferred embodiment, the candidate bioactive agent is labeled, and binding determined directly. For example, this may be done by attaching all or a portion of the colorectal cancer protein to a solid support,

adding a labeled candidate agent (for example a fluorescent label), washing off excess reagent, and determining whether the label is present on the solid support. Various blocking and washing steps may be utilized as is known in the art.

[202] By "labeled" herein is meant that the compound is either directly or indirectly labeled with a label which provides a detectable signal, e.g. radioisotope, fluorescers, enzyme, antibodies, particles such as magnetic particles, chemiluminescers, or specific binding molecules, etc. Specific binding molecules include pairs, such as biotin and streptavidin, digoxin and antidigoxin etc. For the specific binding members, the complementary member would normally be labeled with a molecule which provides for detection, in accordance with known procedures, as outlined above. The label can directly or indirectly provide a detectable signal.

[203] In some embodiments, only one of the components is labeled. For example, the proteins (or proteinaceous candidate agents) may be labeled at tyrosine positions using <sup>125</sup>I, or with fluorophores. Alternatively, more than one component may be labeled with different labels; using <sup>125</sup>I for the proteins, for example, and a fluorophor for the candidate agents.

[204] In a preferred embodiment, the binding of the candidate bioactive agent is determined through the use of competitive binding assays. In this embodiment, the competitor is a binding moiety known to bind to the target molecule (i.e. colorectal cancer ), such as an antibody, peptide, binding partner, ligand, etc. Under certain circumstances, there may be competitive binding as between the bioactive agent and the binding moiety, with the binding moiety displacing the bioactive agent.

[205] In one embodiment, the candidate bioactive agent is labeled. Either the candidate bioactive agent, or the competitor, or both, is added first to the protein for a time sufficient to allow binding, if present. Incubations may be performed at any temperature which facilitates optimal activity, typically between 4 and 40°C. Incubation periods are selected for optimum activity, but may also be optimized to facilitate rapid high through put screening. Typically between 0.1 and 1 hour will be sufficient. Excess reagent is generally removed or washed away. The second component is then added, and the presence or absence of the labeled component is followed, to indicate binding.

[206] In a preferred embodiment, the competitor is added first, followed by the candidate bioactive agent. Displacement of the competitor is an indication that the candidate bioactive agent is binding to the colorectal cancer protein and thus is capable of binding to, and potentially modulating, the activity of the colorectal cancer protein. In this

embodiment, either component can be labeled. Thus, for example, if the competitor is labeled, the presence of label in the wash solution indicates displacement by the agent. Alternatively, if the candidate bioactive agent is labeled, the presence of the label on the support indicates displacement.

5 [207] In an alternative embodiment, the candidate bioactive agent is added first, with incubation and washing, followed by the competitor. The absence of binding by the competitor may indicate that the bioactive agent is bound to the colorectal cancer protein with a higher affinity. Thus, if the candidate bioactive agent is labeled, the presence of the label on the support, coupled with a lack of competitor binding, may indicate that the  
10 candidate agent is capable of binding to the colorectal cancer protein.

[208] In a preferred embodiment, the methods comprise differential screening to identify bioactive agents that are capable of modulating the activity of the colorectal cancer proteins. In this embodiment, the methods comprise combining a colorectal cancer protein and a competitor in a first sample. A second sample comprises a  
15 candidate bioactive agent, a colorectal cancer protein and a competitor. The binding of the competitor is determined for both samples, and a change, or difference in binding between the two samples indicates the presence of an agent capable of binding to the colorectal cancer protein and potentially modulating its activity. That is, if the binding of the competitor is different in the second sample relative to the first sample, the agent is capable  
20 of binding to the colorectal cancer protein.

[209] Alternatively, a preferred embodiment utilizes differential screening to identify drug candidates that bind to the native colorectal cancer protein, but cannot bind to modified colorectal cancer proteins. The structure of the colorectal cancer protein may be modeled, and used in rational drug design to synthesize agents that interact with that site.  
25 Drug candidates that affect colorectal cancer bioactivity are also identified by screening drugs for the ability to either enhance or reduce the activity of the protein.

[210] Positive controls and negative controls may be used in the assays. Preferably all control and test samples are performed in at least triplicate to obtain statistically significant results. Incubation of all samples is for a time sufficient for the  
30 binding of the agent to the protein. Following incubation, all samples are washed free of non-specifically bound material and the amount of bound, generally labeled agent determined. For example, where a radiolabel is employed, the samples may be counted in a scintillation counter to determine the amount of bound compound.

[211] A variety of other reagents may be included in the screening assays.

These include reagents like salts, neutral proteins, e.g. albumin, detergents, etc which may be used to facilitate optimal protein-protein binding and/or reduce non-specific or background interactions. Also reagents that otherwise improve the efficiency of the assay, such as

5 protease inhibitors, nuclease inhibitors, anti-microbial agents, etc., may be used. The mixture of components may be added in any order that provides for the requisite binding.

[212] Screening for agents that modulate the activity of colorectal cancer proteins may also be done. In a preferred embodiment, methods for screening for a bioactive agent capable of modulating the activity of colorectal cancer proteins comprise the steps of  
10 adding a candidate bioactive agent to a sample of colorectal cancer proteins, as above, and determining an alteration in the biological activity of colorectal cancer proteins.

“Modulating the activity of colorectal cancer ” includes an increase in activity, a decrease in activity, or a change in the type or kind of activity present. Thus, in this embodiment, the candidate agent should both bind to colorectal cancer proteins (although this may not be necessary), and alter its biological or biochemical activity as defined herein. The methods include both in vitro screening methods, as are generally outlined above, and in vivo screening of cells for alterations in the presence, distribution, activity or amount of colorectal cancer proteins.

[213] Thus, in this embodiment, the methods comprise combining a  
20 colorectal cancer sample and a candidate bioactive agent, and evaluating the effect on colorectal cancer activity. By “colorectal cancer activity” or grammatical equivalents herein is meant one of the colorectal cancer 's biological activities, including, but not limited to, cell division, preferably in colon tissue, cell proliferation, tumor growth, transformation of cells. In one embodiment, colorectal cancer activity includes activation of a gene identified by a  
25 nucleic acid of Table 1. An inhibitor of colorectal cancer activity is the inhibition of any one or more colorectal cancer activities.

[214] In a preferred embodiment, the activity of the colorectal cancer protein is increased; in another preferred embodiment, the activity of the colorectal cancer protein is decreased. Thus, bioactive agents that are antagonists are preferred in some embodiments,  
30 and bioactive agents that are agonists may be preferred in other embodiments.

[215] In a preferred embodiment, the invention provides methods for screening for bioactive agents capable of modulating the activity of a colorectal cancer protein. The methods comprise adding a candidate bioactive agent, as defined above, to a cell comprising colorectal cancer proteins. Preferred cell types include almost any cell. The

cells contain a recombinant nucleic acid that encodes a colorectal cancer protein. In a preferred embodiment, a library of candidate agents are tested on a plurality of cells.

[216] In one aspect, the assays are evaluated in the presence or absence or previous or subsequent exposure of physiological signals, for example hormones, antibodies, peptides, antigens, cytokines, growth factors, action potentials, pharmacological agents including chemotherapeutics, radiation, carcinogenics, or other cells (i.e. cell-cell contacts). In another example, the determinations are determined at different stages of the cell cycle process.

[217] In this way, bioactive agents are identified. Compounds with pharmacological activity are able to enhance or interfere with the activity of the colorectal cancer protein. In one embodiment, "colorectal cancer protein activity" as used herein includes at least one of the following: colorectal cancer activity, binding to the colorectal cancer protein, activation of the colorectal cancer protein or activation of substrates of the colorectal cancer protein by the colorectal cancer protein. In one embodiment, colorectal cancer activity is defined as the unregulated proliferation of colon tissue, or the growth of cancer in colon tissue. In one aspect, colorectal cancer activity as defined herein is related to the activity of the colorectal cancer protein in the upregulation of the colorectal cancer protein in colon cancer tissue.

[218] In another embodiment, colorectal cancer protein activity includes at least one of the following: colorectal cancer activity, binding to the CBF9 nucleic acid or poly peptide of Table 2 or binding to a nucleic acid of Table 1, or a peptide encoded by a nucleic acid of Table 1 or activation of substrates of the gene products identified by a nucleic acid of Table 1 or substrates of CBF9, which is shown in Table 2. In one aspect, colorectal cancer activity as defined herein is related to the activity of genes defined by the nucleic acids of Table 1 or of CBF9 as defined in Table 2, in colon cancer tissue.

[219] In one embodiment, a method of inhibiting colon cancer cell division is provided. The method comprises administration of a colorectal cancer inhibitor.

[220] In another embodiment, a method of inhibiting tumor growth is provided. The method comprises administration of a colorectal cancer inhibitor.

[221] In a further embodiment, methods of treating cells or individuals with cancer are provided. The method comprises administration of a colorectal cancer inhibitor.

[222] In one embodiment, a colorectal cancer inhibitor is an antibody as discussed above. In another embodiment, the colorectal cancer inhibitor is an antisense molecule. Antisense molecules as used herein include antisense or sense oligonucleotides

comprising a single-stranded nucleic acid sequence (either RNA or DNA) capable of binding to target mRNA (sense) or DNA (antisense) sequences for colorectal cancer molecules. A preferred antisense molecule is for the colorectal cancer sequences referenced in Table 1 or Table 2, or for a ligand or activator thereof. Antisense or sense oligonucleotides, according  
5 to the present invention, comprise a fragment generally at least about 14 nucleotides, preferably from about 14 to 30 nucleotides. The ability to derive an antisense or a sense oligonucleotide, based upon a cDNA sequence encoding a given protein is described in, for example, Stein and Cohen (Cancer Res. 48:2659, 1988) and van der Krol et al.  
(BioTechniques 6:958, 1988).

10 [223] Antisense molecules may be introduced into a cell containing the target nucleotide sequence by formation of a conjugate with a ligand binding molecule, as described in WO 91/04753. Suitable ligand binding molecules include, but are not limited to, cell surface receptors, growth factors, other cytokines, or other ligands that bind to cell surface receptors. Preferably, conjugation of the ligand binding molecule does not substantially  
15 interfere with the ability of the ligand binding molecule to bind to its corresponding molecule or receptor, or block entry of the sense or antisense oligonucleotide or its conjugated version into the cell. Alternatively, a sense or an antisense oligonucleotide may be introduced into a cell containing the target nucleic acid sequence by formation of an oligonucleotide-lipid complex, as described in WO 90/10448. It is understood that the use of antisense molecules  
20 or knock out and knock in models may also be used in screening assays as discussed above, in addition to methods of treatment.

[224] The compounds having the desired pharmacological activity may be administered in a physiologically acceptable carrier to a host, as previously described. The agents may be administered in a variety of ways, orally, parenterally e.g., subcutaneously,  
25 intraperitoneally, intravascularly, etc. Depending upon the manner of introduction, the compounds may be formulated in a variety of ways. The concentration of therapeutically active compound in the formulation may vary from about 0.1-100 wt.%. The agents may be administered alone or in combination with other treatments, i.e., radiation.

[225] The pharmaceutical compositions can be prepared in various forms, such as granules, tablets, pills, suppositories, capsules, suspensions, salves, lotions and the like. Pharmaceutical grade organic or inorganic carriers and/or diluents suitable for oral and topical use can be used to make up compositions containing the therapeutically-active compounds. Diluents known to the art include aqueous media, vegetable and animal oils and fats. Stabilizing agents, wetting and emulsifying agents, salts for varying the osmotic

pressure or buffers for securing an adequate pH value, and skin penetration enhancers can be used as auxiliary agents.

[226] Without being bound by theory, it appears that the various colorectal cancer sequences are important in colorectal cancer. Accordingly, disorders based on 5 mutant or variant colorectal cancer genes may be determined. In one embodiment, the invention provides methods for identifying cells containing variant colorectal cancer genes comprising determining all or part of the sequence of at least one endogenous colorectal cancer genes in a cell. As will be appreciated by those in the art, this may be done using any number of sequencing techniques. In a preferred embodiment, the invention provides 10 methods of identifying the colorectal cancer genotype of an individual comprising determining all or part of the sequence of at least one colorectal cancer gene of the individual. This is generally done in at least one tissue of the individual, and may include the evaluation of a number of tissues or different samples of the same tissue. The method may include comparing the sequence of the sequenced colorectal cancer gene to a known 15 colorectal cancer gene, i.e. a wild-type gene.

[227] The sequence of all or part of the colorectal cancer gene can then be compared to the sequence of a known colorectal cancer gene to determine if any differences exist. This can be done using any number of known homology programs, such as Bestfit, etc. In a preferred embodiment, the presence of a difference in the sequence between the 20 colorectal cancer gene of the patient and the known colorectal cancer gene is indicative of a disease state or a propensity for a disease state, as outlined herein.

[228] In a preferred embodiment, the colorectal cancer genes are used as probes to determine the number of copies of the colorectal cancer gene in the genome.

[229] In another preferred embodiment colorectal cancer genes are used as 25 probed to determine the chromosomal localization of the colorectal cancer genes. Information such as chromosomal localization finds use in providing a diagnosis or prognosis in particular when chromosomal abnormalities such as translocations, and the like are identified in colorectal cancer gene loci.

[230] Thus, in one embodiment, methods of modulating colorectal cancer in 30 cells or organisms are provided. In one embodiment, the methods comprise administering to a cell an anti-colorectal cancer antibody that reduces or eliminates the biological activity of an endogenous colorectal cancer protein. Alternatively, the methods comprise administering to a cell or organism a recombinant nucleic acid encoding a colorectal cancer protein. As will be appreciated by those in the art, this may be accomplished in any number

of ways. In a preferred embodiment, for example when the colorectal cancer sequence is down-regulated in colorectal cancer, the activity of the colorectal cancer gene is increased by increasing the amount of colorectal cancer in the cell, for example by overexpressing the endogenous colorectal cancer or by administering a gene encoding the colorectal cancer sequence, using known gene-therapy techniques, for example. In a preferred embodiment, the gene therapy techniques include the incorporation of the endogenous gene using enhanced homologous recombination (EHR), for example as described in PCT/US93/03868, hereby incorporated by reference in its entirety. Alternatively, for example when the colorectal cancer sequence is up-regulated in colorectal cancer, the activity of the endogenous colorectal cancer gene is decreased, for example by the administration of a colorectal cancer antisense nucleic acid.

[231] In one embodiment, the colorectal cancer proteins of the present invention may be used to generate polyclonal and monoclonal antibodies to colorectal cancer proteins, which are useful as described herein. Similarly, the colorectal cancer proteins can be coupled, using standard technology, to affinity chromatography columns. These columns may then be used to purify colorectal cancer antibodies. In a preferred embodiment, the antibodies are generated to epitopes unique to a colorectal cancer protein; that is, the antibodies show little or no cross-reactivity to other proteins. These antibodies find use in a number of applications. For example, the colorectal cancer antibodies may be coupled to standard affinity chromatography columns and used to purify colorectal cancer proteins. The antibodies may also be used as blocking polypeptides, as outlined above, since they will specifically bind to the colorectal cancer protein.

[232] In one embodiment, a therapeutically effective dose of a colorectal cancer or modulator thereof is administered to a patient. By "therapeutically effective dose" herein is meant a dose that produces the effects for which it is administered. The exact dose will depend on the purpose of the treatment, and will be ascertainable by one skilled in the art using known techniques. As is known in the art, adjustments for colorectal cancer degradation, systemic versus localized delivery, and rate of new protease synthesis, as well as the age, body weight, general health, sex, diet, time of administration, drug interaction and the severity of the condition may be necessary, and will be ascertainable with routine experimentation by those skilled in the art.

[233] A "patient" for the purposes of the present invention includes both humans and other animals, particularly mammals, and organisms. Thus the methods are

applicable to both human therapy and veterinary applications. In the preferred embodiment the patient is a mammal, and in the most preferred embodiment the patient is human.

[234] The administration of the colorectal cancer proteins and modulators of the present invention can be done in a variety of ways as discussed above, including, but 5 not limited to, orally, subcutaneously, intravenously, intranasally, transdermally, intraperitoneally, intramuscularly, intrapulmonary, vaginally, rectally, or intraocularly. In some instances, for example, in the treatment of wounds and inflammation, the colorectal cancer proteins and modulators may be directly applied as a solution or spray.

[235] The pharmaceutical compositions of the present invention comprise a 10 colorectal cancer protein in a form suitable for administration to a patient. In the preferred embodiment, the pharmaceutical compositions are in a water soluble form, such as being present as pharmaceutically acceptable salts, which is meant to include both acid and base addition salts. "Pharmaceutically acceptable acid addition salt" refers to those salts that retain the biological effectiveness of the free bases and that are not biologically or otherwise 15 undesirable, formed with inorganic acids such as hydrochloric acid, hydrobromic acid, sulfuric acid, nitric acid, phosphoric acid and the like, and organic acids such as acetic acid, propionic acid, glycolic acid, pyruvic acid, oxalic acid, maleic acid, malonic acid, succinic acid, fumaric acid, tartaric acid, citric acid, benzoic acid, cinnamic acid, mandelic acid, methanesulfonic acid, ethanesulfonic acid, p-toluenesulfonic acid, salicylic acid and the like. 20 "Pharmaceutically acceptable base addition salts" include those derived from inorganic bases such as sodium, potassium, lithium, ammonium, calcium, magnesium, iron, zinc, copper, manganese, aluminum salts and the like. Particularly preferred are the ammonium, potassium, sodium, calcium, and magnesium salts. Salts derived from pharmaceutically acceptable organic non-toxic bases include salts of primary, secondary, and tertiary amines, 25 substituted amines including naturally occurring substituted amines, cyclic amines and basic ion exchange resins, such as isopropylamine, trimethylamine, diethylamine, triethylamine, tripropylamine, and ethanolamine.

[236] The pharmaceutical compositions may also include one or more of the following: carrier proteins such as serum albumin; buffers; fillers such as microcrystalline 30 cellulose, lactose, corn and other starches; binding agents; sweeteners and other flavoring agents; coloring agents; and polyethylene glycol. Additives are well known in the art, and are used in a variety of formulations.

[237] In a preferred embodiment, colorectal cancer proteins and modulators are administered as therapeutic agents, and can be formulated as outlined above. Similarly,

colorectal cancer genes (including both the full-length sequence, partial sequences, or regulatory sequences of the colorectal cancer coding regions) can be administered in gene therapy applications, as is known in the art. These colorectal cancer genes can include antisense applications, either as gene therapy (i.e. for incorporation into the genome) or as antisense compositions, as will be appreciated by those in the art.

[238] In a preferred embodiment, colorectal cancer genes are administered as DNA vaccines, either single genes or combinations of colorectal cancer genes. Naked DNA vaccines are generally known in the art. Brower, Nature Biotechnology, 16:1304-1305 (1998).

[239] In one embodiment, colorectal cancer genes of the present invention are used as DNA vaccines. Methods for the use of genes as DNA vaccines are well known to one of ordinary skill in the art, and include placing a colorectal cancer gene or portion of a colorectal cancer gene under the control of a promoter for expression in a colorectal cancer patient. The colorectal cancer gene used for DNA vaccines can encode full-length colorectal cancer proteins, but more preferably encodes portions of the colorectal cancer proteins including peptides derived from the colorectal cancer protein. In a preferred embodiment a patient is immunized with a DNA vaccine comprising a plurality of nucleotide sequences derived from a colorectal cancer gene. Similarly, it is possible to immunize a patient with a plurality of colorectal cancer genes or portions thereof as defined herein. Without being bound by theory, expression of the polypeptide encoded by the DNA vaccine, cytotoxic T-cells, helper T-cells and antibodies are induced which recognize and destroy or eliminate cells expressing colorectal cancer proteins.

[240] In a preferred embodiment, the DNA vaccines include a gene encoding an adjuvant molecule with the DNA vaccine. Such adjuvant molecules include cytokines that increase the immunogenic response to the colorectal cancer polypeptide encoded by the DNA vaccine. Additional or alternative adjuvants are known to those of ordinary skill in the art and find use in the invention.

[241] In another preferred embodiment colorectal cancer genes find use in generating animal models of colorectal cancer. As is appreciated by one of ordinary skill in the art, when the colorectal cancer gene identified is repressed or diminished in colorectal cancer tissue, gene therapy technology wherein antisense RNA directed to the colorectal cancer gene will also diminish or repress expression of the gene. An animal generated as such serves as an animal model of colorectal cancer that finds use in screening bioactive drug candidates. Similarly, gene knockout technology, for example as a result of

homologous recombination with an appropriate gene targeting vector, will result in the absence of the colorectal cancer protein. When desired, tissue-specific expression or knockout of the colorectal cancer protein may be necessary.

[242] It is also possible that the colorectal cancer protein is overexpressed in colorectal cancer. As such, transgenic animals can be generated that overexpress the colorectal cancer protein. Depending on the desired expression level, promoters of various strengths can be employed to express the transgene. Also, the number of copies of the integrated transgene can be determined and compared for a determination of the expression level of the transgene. Animals generated by such methods find use as animal models of colorectal cancer and are additionally useful in screening for bioactive molecules to treat colorectal cancer .

## EXAMPLES

[243] It is understood that the examples described herein in no way serve to limit the true scope of this invention, but rather are presented for illustrative purposes. All references and sequences of accession numbers cited herein are incorporated by reference in their entirety.

[244] Example 1

Tissue Preparation, Labeling Chips, and Fingerprints

[245] Purify total RNA from tissue using TRIzol Reagent

[246] Estimate tissue weight. Homogenize tissue samples in 1ml of TRIzol per 50mg of tissue using a Polytron 3100 homogenizer. The generator/probe used depends upon the tissue size. A generator that is too large for the amount of tissue to be homogenized will cause a loss of sample and lower RNA yield. Use the 20mm generator for tissue weighing more than 0.6g. If the working volume is greater than 2ml, then homogenize tissue in a 15ml polypropylene tube (Falcon 2059). Fill tube no greater than 10ml.

## HOMOGENIZATION

[247] Before using generator, it should have been cleaned after last usage by running it through soapy H<sub>2</sub>O and rinsing thoroughly. Run through with EtOH to sterilize. Keep tissue frozen until ready. Add TRIzol directly to frozen tissue then homogenize.

[248] Following homogenization, remove insoluble material from the homogenate by centrifugation at 7500 x g for 15 min. in a Sorvall superspeed or 12,000 x g for 10 min. in an Eppendorf centrifuge at 4oC. Transfer the cleared homogenate to a new tube(s). The samples may be frozen now at -60 to -70oC (and kept for at least one month) or  
5 you may continue with the purification.

### PHASE SEPARATION

[249] Incubate the homogenized samples for 5 minutes at room temperature.

[250] Add 0.2ml of chloroform per 1ml of TRIzol reagent used in the  
10 original homogenization.

[251] Cap tubes securely and shake tubes vigorously by hand (do not vortex) for 15 seconds.

[252] Incubate samples at room temp. for 2-3 minutes. Centrifuge samples at 6500rpm in a Sorvall superspeed for 30 min. at 4oC. (You may spin at up to 12,000 x g  
15 for 10 min. but you risk breaking your tubes in the centrifuge.)

### RNA PRECIPITATION

[253] Transfer the aqueous phase to a fresh tube. Save the organic phase if isolation of DNA or protein is desired. Add 0.5ml of isopropyl alcohol per 1ml of TRIzol reagent used in the original homogenization. Cap tubes securely and invert to mix. Incubate samples at room temp. for 10 minutes. Centrifuge samples at 6500rpm in Sorvall for 20min. at 4oC.  
20

### RNA WASH

[254] Pour off the supernate. Wash pellet with cold 75% ethanol. Use 1ml of 75% ethanol per 1ml of TRIzol reagent used in the initial homogenization. Cap tubes securely and invert several times to loosen pellet. (Do not vortex). Centrifuge at <8000rpm (<7500 x g) for 5 minutes at 4oC.  
25

[255] Pour off the wash. Carefully transfer pellet to an eppendorf tube (let it slide down the tube into the new tube and use a pipet tip to help guide it in if necessary). Depending on the volumes you are working with, you can decide what size tube(s) you want to precipitate the RNA in. When I tried leaving the RNA in the large 15ml tube, it took so long to dry (i.e. it did not dry) that I eventually had to transfer it to a smaller tube. Let pellet  
30

dry in hood. Resuspend RNA in an appropriate volume of DEPC H<sub>2</sub>O. Try for 2-5ug/ul. Take absorbance readings.

[256] Purify poly A+ mRNA from total RNA or clean up total RNA with

5 Qiagen's RNeasy kit

[257] Purification of poly A+ mRNA from total RNA. Heat oligotex suspension to 37oC and mix immediately before adding to RNA. Incubate Elution Buffer at 70oC. Warm up 2 x Binding Buffer at 65oC if there is precipitate in the buffer. Mix total RNA with DEPC-treated water, 2 x Binding Buffer, and Oligotex according to Table 2 on 10 page 16 of the Oligotex Handbook. Incubate for 3 minutes at 65oC. Incubate for 10 minutes at room temperature.

[258] Centrifuge for 2 minutes at 14,000 to 18,000 g. If centrifuge has a "soft setting," then use it. Remove supernatant without disturbing Oligotex pellet. A little bit 15 of solution can be left behind to reduce the loss of Oligotex. Save sup until certain that satisfactory binding and elution of poly A+ mRNA has occurred.

[259] Gently resuspend in Wash Buffer OW2 and pipet onto spin column. Centrifuge the spin column at full speed (soft setting if possible) for 1 minute.

20

[260] Transfer spin column to a new collection tube and gently resuspend in Wash Buffer OW2 and centrifuge as describe herein.

[261] Transfer spin column to a new tube and elute with 20 to 100 ul of 25 preheated (70oC) Elution Buffer. Gently resuspend Oligotex resin by pipetting up and down. Centrifuge as above. Repeat elution with fresh elution buffer or use first eluate to keep the elution volume low.

[262] Read absorbance, using diluted Elution Buffer as the blank.

30

[263] Before proceeding with cDNA synthesis, the mRNA must be precipitated. Some component leftover or in the Elution Buffer from the Oligotex purification procedure will inhibit downstream enzymatic reactions of the mRNA.

Ethanol Precipitation

- [264] Add 0.4 vol. of 7.5 M NH<sub>4</sub>OAc + 2.5 vol. of cold 100% ethanol. Precipitate at -20°C 1 hour to overnight (or 20-30 min. at -70°C). Centrifuge at 14,000-16,000 x g for 30 minutes at 4°C. Wash pellet with 0.5ml of 80%ethanol (-20°C) then 5 centrifuge at 14,000-16,000 x g for 5 minutes at room temperature. Repeat 80% ethanol wash. Dry the last bit of ethanol from the pellet in the hood. (Do not speed vacuum). Suspend pellet in DEPC H<sub>2</sub>O at 1ug/ul concentration.

Clean up total RNA using Qiagen's RNeasy kit

- 10 [265] Add no more than 100ug to an RNeasy column. Adjust sample to a volume of 100ul with RNase-free water. Add 350ul Buffer RLT then 250ul ethanol (100%) to the sample. Mix by pipetting (do not centrifuge) then apply sample to an RNeasy mini spin column. Centrifuge for 15 sec at >10,000rpm. If concerned about yield, re-apply flowthrough to column and centrifuge again.
- 15 [266] Transfer column to a new 2-ml collection tube. Add 500ul Buffer RPE and centrifuge for 15 sec at >10,000rpm. Discard flowthrough. Add 500ul Buffer RPE and centrifuge for 15 sec at >10,000rpm. Discard flowthrough then centrifuge for 2 min at maximum speed to dry column membrane. Transfer column to a new 1.5-ml collection tube and apply 30-50ul of RNase-free water directly onto column membrane. Centrifuge 1 min at 20 >10,000rpm. Repeat elution.

[267] Take absorbance reading. If necessary, ethanol precipitate with ammonium acetate and 2.5X volume 100% ethanol.

- 25 [268] Make cDNA using Gibco's "SuperScript Choice System for cDNA Synthesis" kit

First Strand cDNA Synthesis

- [269] Use 5ug of total RNA or 1ug of polyA+ mRNA as starting material. For total RNA, use 2ul of SuperScript RT. For polyA+ mRNA, use 1ul of SuperScript RT. Final volume of first strand synthesis mix is 20ul. RNA must be in a volume no greater than 30 10ul. Incubate RNA with 1ul of 100pmol T7-T24 oligo for 10 min at 70C. On ice, add 7 ul of: 4ul 5X 1st Strand Buffer, 2ul of 0.1M DTT, and 1 ul of 10mM dNTP mix. Incubate at 37C for 2 min then add SuperScript RT

Incubate at 37C for 1 hour.

Second Strand Synthesis

Place 1st strand reactions on ice.

Add: 91ul DEPC H2O

30ul 5X 2nd Strand Buffer

3ul 10mM dNTP mix

- 5  
1ul 10U/ul E.coli DNA Ligase  
4ul 10U/ul E.coli DNA Polymerase  
1ul 2U/ul RNase H

[270] Make the above into a mix if there are more than 2 samples. Mix and  
10 incubate 2 hours at 16C.

[271] Add 2ul T4 DNA Polymerase. Incubate 5 min at 16C. Add 10ul of  
0.5M EDTA

[272] Clean up cDNA

15 [273] Phenol:Chloroform:Isoamyl Alcohol (25:24:1) purification using  
Phase-Lock gel tubes:

20 [274] Centrifuge PLG tubes for 30 sec at maximum speed. Transfer cDNA  
mix to PLG tube. Add equal volume of phenol:chloroform:isamyl alcohol and shake  
vigorously (do not vortex). Centrifuge 5 minutes at maximum speed. Transfer top aqueous  
solution to a new tube. Ethanol precipitate: add 7.5X 5M NH4Oac and 2.5X volume of  
100% ethanol. Centrifuge immediately at room temp. for 20 min, maximum speed. Remove  
sup then wash pellet 2X with cold 80% ethanol. Remove as much ethanol wash as possible  
then let pellet air dry. Resuspend pellet in 3ul RNase-free water.

25 In vitro Transcription (IVT) and labeling with biotin  
Pipet 1.5ul of cDNA into a thin-wall PCR tube.

Make NTP labeling mix:

30 Combine at room temperature: 2ul T7 10xATP (75mM) (Ambion)  
2ul T7 10xGTP (75mM) (Ambion)  
1.5ul T7 10xCTP (75mM) (Ambion)  
1.5ul T7 10xUTP (75mM) (Ambion)  
3.75ul 10mM Bio-11-UTP (Boehringer-Mannheim/Roche or Enzo)  
3.75ul 10mM Bio-16-CTP (Enzo)

2ul 10x T7 transcription buffer (Ambion)  
2ul 10x T7 enzyme mix (Ambion)

- 5 [275] Final volume of total reaction is 20ul. Incubate 6 hours at 37C in a PCR machine.

#### RNeasy clean-up of IVT product

[276] Follow previous instructions for RNeasy columns or refer to Qiagen's RNeasy protocol handbook.

10

[277] cRNA will most likely need to be ethanol precipitated. Resuspend in a volume compatible with the fragmentation step.

#### Fragmentation

15 [278] 15 ug of labeled RNA is usually fragmented. Try to minimize the fragmentation reaction volume; a 10 ul volume is recommended but 20 ul is all right. Do not go higher than 20 ul because the magnesium in the fragmentation buffer contributes to precipitation in the hybridization buffer.

20 [279] Fragment RNA by incubation at 94 C for 35 minutes in 1 x Fragmentation buffer.

5 x Fragmentation buffer:

200 mM Tris-acetate, pH 8.1

500 mM KOAc

25 150 mM MgOAc

30 [280] The labeled RNA transcript can be analyzed before and after fragmentation. Samples can be heated to 65C for 15 minutes and electrophoresed on 1% agarose/TBE gels to get an approximate idea of the transcript size range

#### Hybridization

[281] 200 ul (10ug cRNA) of a hybridization mix is put on the chip. If multiple hybridizations are to be done (such as cycling through a 5 chip set), then it is recommended that an initial hybridization mix of 300 ul or more be made.

Hybridization Mix: fragment labeled RNA (50ng/ $\mu$ l final conc.)  
50 pM 948-b control oligo  
1.5 pM BioB  
5 pM BioC  
25 pM BioD  
100 pM CRE  
0.1mg/ml herring sperm DNA  
0.5mg/ml acetylated BSA  
10 to 300  $\mu$ l with 1xMES hyb. buffer

[282] The instruction manuals for the products used herein are incorporated herein in their entirety.

15 Labeling Protocol Provided Herein

Hybridization reaction:

Start with non-biotinylated IVT (purified by RNeasy columns)  
(see example 1 for steps from tissue to IVT)

IVT antisense RNA; 4  $\mu$ g:  $\mu$ l

20 Random Hexamers (1  $\mu$ g/ $\mu$ l): 4  $\mu$ l

H<sub>2</sub>O:  $\mu$ l

---

14  $\mu$ l

25 - Incubate 70°C, 10 min. Put on ice.

Reverse transcription:

5X First Strand (BRL) buffer: 6  $\mu$ l

0.1 M DTT: 3  $\mu$ l

30 50X dNTP mix: 0.6  $\mu$ l

H<sub>2</sub>O: 2.4  $\mu$ l

Cy3 or Cy5 dUTP (1mM): 3  $\mu$ l

SS RT II (BRL): 1  $\mu$ l

16  $\mu$ l

- Add to hybridization reaction.
- Incubate 30 min., 42°C.
- Add 1 µl SSII and let go for another hour.

Put on ice.

5 - 50X dNTP mix (25mM of cold dATP, dCTP, and dGTP, 10mM of dTTP: 25 µl each of 100mM dATP, dCTP, and dGTP; 10 µl of 100mM dTTP to 15 µl H2O. dNTPs from Pharmacia)

#### RNA degradation:

10 86 µl H2O  
- Add 1.5 µl 1M NaOH/ 2mM EDTA, incubate at 65°C, 10 min.  
10 µl 10N NaOH  
4 µl 50mM EDTA  
U-Con 30  
15 500 µl TE/sample spin at 7000g for 10 min, save flow through for purification

#### Qiagen purification:

20 -suspend u-con recovered material in 500µl buffer PB  
-proceed w/ normal Qiagen protocol  
DNAse digest:  
- Add 1 µl of 1/100 dil of DNase/30µl Rx and incubate at 37°C for 15 min.  
-5 min 95°C to denature enzyme

#### Sample preparation:

25 - Add:  
Cot-1 DNA: 10 µl  
50X dNTPs: 1 µl  
Na pyro phosphate: 7.5 µl  
10mg/ml Herring sperm DNA 1ul of 1/10 dilution  
30 21.8 final vol.  
- Dry down in speed vac.  
- Resuspend in 15 µl H2O.  
- Add 0.38 µl 10% SDS.  
- Heat 95°C, 2 min.

- Slow cool at room temp. for 20 min.
- Put on slide and hybridize overnight at 64°C.

5

**Washing after the hybridization:**

3X SSC/0.03% SDS:            2 min. 37.5 ml 20X SSC+0.75ml 10% SDS in  
250ml H<sub>2</sub>O

1X SSC: 5 min.                12.5 ml 20X SSC in 250ml H<sub>2</sub>O

0.2X SSC: 5 min.                2.5 ml 20X SSC in 250ml H<sub>2</sub>O

10

Dry slides in centrifuge, 1000 RPM, 1min.

[283] Scan using appropriate Photomultiplier tube (PMT) and fluorescent excitation and emission channels.

[284] The results are shown in Table 1 and Table 2. The lists of genes come from colorectal tumors from a variety of stages of the disease. The genes that are up regulated in the tumors (overall) were also found to be expressed at a limited amount or not at all in the body map. The body map consists of at least 28 tissue types, including Adrenal Gland, Bladder, Bone Marrow, Brain, Breast, Cervix, Colon, Diaphragm, Heart, Kidney, Liver, Lung, Lymph Node, Muscle, Pancreas, Prostate, Rectum, Salivary Gland, Skin, Small Intestine, Spinal Cord, Spleen, Stomach, Testis, Thymus, Thyroid Trachea and Uterus. As indicated, some of the Accession numbers include expression sequence tags (ESTs). Thus, in one embodiment herein, genes within an expression profile, also termed expression profile genes, include ESTs and are not necessarily full length.

[285] Table 1 shows Accession numbers for 1747 genes upregulated in colon tumor tissue. The table provides the exemplar accession numbers, Unigene ID numbers, unique Eos codes, descriptions of the genes encoded, and relative amount of expression as compared with expression in other normal body tissue.

**TABLE 1. GENES INVOLVED IN COLORECTAL CANCER**

	<b>Pkey</b>	Primekey(unique probeset identifier)				
	<b>Ex. Accn.</b>	Exemplar accession number				
	<b>Probeset</b>	Eos Code number				
	<b>Unigene#</b>	Unigene number				
5						
10	<b>Pkey</b>	<b>Probeset</b>	<b>Ex_Accn</b>	<b>UniG_ID</b>	<b>UniGene Title</b>	<b>Ratio TumMet/Body</b>
15	332264	EOS32195	N72849	Hs.115263	epiregulin	17.6
	332716	EOS32647	L00058	Hs.79070	v-myc avian myelocytomatisis viral oncogene homolog	15.0
20	312845	EOS12776	AI911215	Hs.186555	ESTs	14.3
	310257	EOS10188	AW389247	Hs.148826	ESTs	11.6
25	322567	EOS22498	AF155108		EST cluster (not in UniGene)	11.5
	331060	EOS30991	N75081	Hs.21648	ESTs	10.3
30	322303	EOS22234	W07459		EST cluster (not in UniGene)	9.6
	301891	EOS01822	AF131855	Hs.106127	Homo sapiens clone 25056 mRNA sequence	9.5
35	318524	EOS18455	AW291511	Hs.253687	ESTs	8.9
	314001	EOS13932	AW168495	Hs.8750	ESTs	7.8
40	331183	EOS31114	T40769	Hs.8469	EST	7.3
	315429	EOS15360	AW009951	Hs.206892	ESTs	7.3
45	303344	EOS03275	AA255977	Hs.250646	ESTs; Highly similar to ubiquitin-conjugating enzyme [M.musculus]	6.7
	313625	EOS13556	AW468402	Hs.254020	ESTs	6.7
50	307084	EOS07015	AI160527		EST singleton (not in UniGene) with exon hit	6.1
	314943	EOS14874	AI476797	Hs.184572	cell division cycle 2; G1 to S and G2 to M	6.1
55	303753	EOS03684	AW503733	Hs.170315	ESTs	5.7
	315593	EOS15524	AW198103	Hs.158154	ESTs	5.3
60	313604	EOS13535	AI745325	Hs.182286	ESTs; Moderately similar to !!! ALU SUBFAMILY SB2 WARNING ENTRY !!! [H.sapiens]	5.1
	312319	EOS12250	AA216698	Hs.180780	Homo sapiens agrin precursor mRNA; partial cds	5.1
65	312614	EOS12545	AI766732	Hs.201194	ESTs	4.8
	323176	EOS23107	AW071648	Hs.123199	ESTs	4.8
70	317916	EOS17847	AI565071	Hs.159983	ESTs	4.7
	301846	EOS01777	R20002	Hs.6823	ESTs; Weakly similar to intrinsic factor-B12 receptor precursor [H.sapiens]	4.6
75	311157	EOS11088	AI990122	Hs.196988	ESTs	4.6
	332640	EOS32571	AA417152	Hs.5101	protein regulator of cytokinesis 1	4.6
80	311728	EOS11659	AW083000	Hs.184776	ribosomal protein L23a	4.5
	313774	EOS13705	AW136836	Hs.144583	ESTs	4.5
85	312339	EOS12270	AA524394		EST cluster (not in UniGene)	4.4
	315369	EOS15300	AA764918	Hs.256531	ESTs	4.3
90	303756	EOS03687	AI738488	Hs.115838	ESTs	4.3
	301050	EOS00981	AW136973	Hs.144475	ESTs; Weakly similar to mitogen inducible gene mig-2 [H.sapiens]	4.3
95	300319	EOS00250	AW157646	Hs.153506	ESTs; Weakly similar to microtubule-actin crosslinking factor [M.musculus]	4.3
	300664	EOS0595	AI44628	Hs.256809	ESTs	4.3
100	302655	EOS02586	AJ227892		EST cluster (not in UniGene) with exon hit	4.1
	316175	EOS15106	AI025842	Hs.152530	ESTs	4.1
105	330786	EOS30717	D60374	Hs.258712	EST	4.1
	310875	EOS10806	T47764	Hs.132917	ESTs	4.1
110	313425	EOS13356	AA745689	Hs.186838	ESTs; Weakly similar to similar to zinc finger 5 protein from Gallus gallus; U51640 [H.sapiens]	4.0
	301804	EOS01735	AA581004		EST cluster (not in UniGene) with exon hit	4.0
115	332203	EOS32134	H49388	Hs.102082	EST	3.9
	322968	EOS22899	AI905228		EST cluster (not in UniGene)	3.8
120	321524	EOS21455	N79126		EST cluster (not in UniGene)	3.8
	302476	EOS02407	AF182294		EST cluster (not in UniGene) with exon hit	3.8
125	303295	EOS03226	AA205625	Hs.208067	ESTs	3.8
	310016	EOS09947	AW449612	Hs.152475	ESTs	3.7
130	324871	EOS24802	AW297755	Hs.148832	ESTs	3.7
	322887	EOS22818	AI986306	Hs.233460	ESTs; Weakly similar to KIAA0969 protein [H.sapiens]	3.7
135	313171	EOS13102	NG7879	Hs.157695	ESTs	3.7
	321638	EOS21569	AI356352	Hs.108932	ESTs	3.7
140	320445	EOS20376	R33916		EST cluster (not in UniGene)	3.6
	302149	EOS02080	AI383794	Hs.205237	protein arginine N-methyltransferase 3(hnRNP methyltransferase S. cerevisiae)-like 3	3.6
145	316905	EOS16836	AW138241	Hs.210846	ESTs	3.6
	313166	EOS13097	AI801098	Hs.151500	ESTs	3.6
150	323338	EOS23269	R74219	Hs.23348	S-phase kinase-associated protein 2 (p45)	3.5
	311434	EOS11365	AW016607	Hs.201582	ESTs	3.5
155	312742	EOS12673	AI650363	Hs.116462	ESTs	3.4
	323587	EOS23518	AI905527	Hs.141901	ESTs; Moderately similar to !!! ALU SUBFAMILY SP WARNING ENTRY !!! [H.sapiens]	3.4
160	317390	EOS17321	AW136551	Hs.181245	ESTs	3.4
	315282	EOS15213	AI222165	Hs.144923	ESTs	3.4
165	318565	EOS18496	AI440137	Hs.164989	ESTs	3.4
	307586	EOS07517	AI285499		EST singleton (not in UniGene) with exon hit	3.4
170	321052	EOS20983	AW372884	Hs.240770	nuclear cap binding protein subunit 2; 20kD	3.3
	324338	EOS24269	AL138357	Hs.247514	ESTs	3.3
175	307517	EOS07448	AI275055	Hs.164989	ESTs	3.3
	314852	EOS14783	AI903735	Hs.137527	ESTs; Weakly similar to X-linked retinopathy protein [H.sapiens]	3.3
180	324657	EOS24588	AW451142	Hs.255628	ESTs	3.2
	314912	EOS14843	AI431345	Hs.161784	ESTs	3.2
185	324790	EOS24721	AI334367	Hs.159337	ESTs	3.2
	315498	EOS15429	AA628539	Hs.116252	ESTs; Moderately similar to !!! ALU SUBFAMILY J WARNING ENTRY !!! [H.sapiens]	3.2
190	312857	EOS12788	AA772279	Hs.126914	ESTs	3.2

300762	EOS00693	AI497778	Hs.168053	ESTs		3.2
325587	EOS25518	c12_hs_gi 6682462 ref	gn 1 + 126724 126967 ex 7 7 CDSI 2.44 244 3099 CH.12_hs_gi 6682462			3.2
5	320654	EOS20585	AW263086	Hs.118112	ESTs	3.2
316715	EOS16646	AI440266	Hs.170673	ESTs		3.1
333279	EOS33210	CH22_522FG_126_1_LINK_EM:AC005500.GENSCAN.8-1				
			CH22_FGENES.126_1			3.1
309689	EOS09620	AW236171	Hs.181357	laminin receptor 1 (67kD; ribosomal protein SA)		3.1
323846	EOS23777	AA337621	Hs.137635	ESTs		3.1
10	324678	EOS24609	AI990739	Hs.236511	ESTs; Moderately similar to RNA splicing-related protein [R.norvegicus]	3.1
308362	EOS08293	AI613519		EST singleton (not in UniGene) with exon hit		3.1
308615	EOS08546	AI738593		EST singleton (not in UniGene) with exon hit		3.0
315397	EOS15328	AA218940	Hs.137516	ESTs		3.0
302236	EOS02167	AI128606	Hs.167558	zinc finger protein 161		3.0
321693	EOS21624	AA700017	Hs.173737	ras-related C3 botulinum toxin substrate 1 (rho family; small GTP binding protein Rac1)		3.0
15	330814	EOS30745	AA015730	Hs.247277	ESTs; Weakly similar to transformation-related protein [H.sapiens]	3.0
302977	EOS02908	AW263124		EST cluster (not in UniGene) with exon hit		3.0
327516	EOS27447	c_2_hs_gi 6117815 ref	gn 6 + 199078 199216 ex 4 4 CDSI 9.15 139 1551			
			CH.02_hs_gi 6117815			2.9
20	333278	EOS33209	CH22_521FG_125_2_LINK_EM:AC005500.GENSCAN.7-2			
			CH22_FGENES.125_2			2.9
302088	EOS02019	U77629	Hs.135639	achaete-scute complex (Drosophila) homolog-like 2		2.9
322718	EOS22649	AF150270	Hs.233322	ESTs; Weakly similar to cDNA EST EMBL:T01156 comes from this gene [C.elegans]		2.9
25	329154	EOS29085	c_x_hs_gi 5868686 ref	gn 2 - 200851 201356 ex 1 3 CDSI 30.28 506 1812		
			CH.X_hs_gi 5868686			2.9
315978	EOS15909	AA830893	Hs.119769	ESTs		2.9
302677	EOS02608	H63227	Hs.132880	ESTs; Highly similar to ubiquitin-conjugating enzyme [M.musculus]		2.9
315007	EOS14938	AI806583	Hs.125291	ESTs		2.9
303780	EOS03711	AI424014	Hs.243450	ESTs; Moderately similar to KIAA0456 protein [H.sapiens]		2.9
30	331362	EOS31293	AA417956	Hs.40782	ESTs	2.9
335815	EOS35746	CH22_3187FG_618_3_LINK_EM:AC005500.GENSCAN.510-3				2.8
			CH22_FGENES.618_3			
332070	EOS32001	AA598545	Hs.228138	EST		2.8
315720	EOS15651	AW291875	Hs.163900	ESTs		2.8
311913	EOS11844	AI358522	Hs.221417	ESTs		2.8
331014	EOS30945	H98597	Hs.30340	ESTs		2.8
35	322035	EOS21966	AL137517	EST cluster (not in UniGene)		2.8
338057	EOS37988	CH22_6558FG__LINK_EM:AC005500.GENSCAN.160-1				2.8
			CH22_EM:AC005500.GENSCAN.160-1			
40	335829	EOS35760	CH22_3202FG_620_3_LINK_EM:AC005500.GENSCAN.512-3			
			CH22_FGENES.620_3			2.8
312136	EOS12067	AW451469	Hs.209990	ESTs		2.8
303132	EOS03063	AI929819	Hs.193330	ESTs		2.8
45	317548	EOS17479	AI654187	Hs.195704	ESTs	2.8
325585	EOS25516	c12_hs_gi 6682462 ref	gn 1 + 73476 73574 ex 5 7 CDSI 8.52 99 309			
			7 CH.12_hs_gi 6682462			2.7
334631	EOS34562	CH22_1939FG_416_7_LINK_EM:AC005500.GENSCAN.277-7				2.7
			CH22_FGENES.416_7			
50	329156	EOS29087	c_x_hs_gi 5868686 ref	gn 2 - 202013 202341 ex 3 3 CDSf 10.23 329 1814		
			CH.X_hs_gi 5868686			2.7
318615	EOS18546	AI133617	Hs.191088	ESTs		2.7
300734	EOS00665	AW205197	Hs.240951	ESTs		2.7
55	324430	EOS24361	AA464018	EST cluster (not in UniGene)		2.7
322296	EOS22227	W76326	Hs.251937	ESTs		2.7
303842	EOS03773	AI337304	Hs.126268	ESTs; Weakly similar to PDZ domain [C.elegans]		2.7
320909	EOS20840	D62269		EST cluster (not in UniGene)		2.7
325195	EOS25126	T20258	Hs.171443	ESTs; Weakly similar to actin binding protein MAYVEN [H.sapiens]		2.7
324959	EOS24890	AW367745	Hs.143137	ESTs		2.7
60	309997	EOS09928	AI291621	Hs.145199	ESTs	2.7
329367	EOS29298	c_x_hs_gi 5868842 ref	gn 1 - 87201 87587 ex 1 4 CDSI 8.13 387 3908			
			CH.X_hs_gi 5868842			2.7
316697	EOS16628	AW293174	Hs.252627	ESTs		2.7
313600	EOS13531	AA429564	Hs.185802	ESTs		2.7
65	301471	EOS01402	AA995014	Hs.129544	ESTs; Weakly similar to ORF YLL027w [S.cerevisiae]	2.6
300810	EOS00741	AI076890	Hs.186949	ESTs		2.6
319976	EOS19907	N48809	Hs.250824	ESTs		2.6
313434	EOS13365	W92070	Hs.231902	ESTs		2.6
70	333849	EOS33780	CH22_1118FG_290_8_LINK_EM:AC005500.GENSCAN.146-7			
			CH22_FGENES.290_8			2.6
330744	EOS30675	AA406142	Hs.12393	dTDP-D-glucose 4,6-dehydratase		2.6
309398	EOS09329	AW081820		EST singleton (not in UniGene) with exon hit		2.6
338727	EOS38658	CH22_7523FG__LINK_EM:AC005500.GENSCAN.500-2				2.6
			CH22_EM:AC005500.GENSCAN.500-2			
75	324620	EOS24551	AA448021	EST cluster (not in UniGene)		2.6
335755	EOS35686	CH22_3122FG_604_4_LINK_EM:AC005500.GENSCAN.493-9				2.6
			CH22_FGENES.604_4			
315858	EOS15789	AA737345		EST cluster (not in UniGene)		2.6
307288	EOS07219	AI205169		EST singleton (not in UniGene) with exon hit		2.5
80	330542	EOS30473	U23942	Hs.226213	cytochrome P450; 51 (lanosterol 14-alpha-demethylase)	2.5
335896	EOS35827	CH22_3273FG_635_4_LINK_EM:AC005500.GENSCAN.525-6				2.5
			CH22_FGENES.635_4			
316578	EOS16509	AA775623	Hs.211683	ESTs		2.5
329193	EOS29124	c_x_hs_gi 5868716 ref	gn 3 + 168095 168181 ex 9 9 CDSI -1.11 87 2064			
			CH.X_hs_gi 5868716			2.5
85	315193	EOS15124	AI241331	Hs.131765	ESTs	2.5
319478	EOS19409	R06841		EST cluster (not in UniGene)		2.5

334727	EOS34658	CH22_2038FG_424_1_LINK_EM:AC005500.GENSCAN.285-3 CH22_FGENES.424_1		2.5
328113	EOS28044	c_6_hs_gi 5868024 refl gn 2 - 80378 80491 ex 2 3 CDSi 3.89 114 3247 CH.06_hs_gi 5868024		2.5
5	315214	EOS15145	AI915927 Hs.34771 ESTs	2.5
	324718	EOS24649	AI557019 Hs.116467 ESTs	2.5
	313326	EOS13257	AI088120 Hs.122329 ESTs	2.5
	319480	EOS19411	R06933 Hs.184221 ESTs	2.5
10	317902	EOS17833	AI828602 Hs.211265 ESTs	2.5
	323341	EOS23272	AL134875 Hs.192386 ESTs	2.5
	336003	EOS35934	CH22_3385FG_664_4_LINK_DJ32I10.GENSCAN.5-4 CH22_FGENES.664_4	2.5
	322992	EOS22923	AA142891 Hs.193165 ESTs	2.5
	314911	EOS14842	AW292329 Hs.163481 ESTs	2.5
15	313603	EOS13534	AW468119 EST cluster (not in UniGene)	2.5
	306469	EOS06400	AA983792 EST singleton (not in UniGene) with exon hit	2.5
	324715	EOS24646	AI739168 EST cluster (not in UniGene)	2.5
	302455	EOS02386	AA356923 Hs.240770 nuclear cap binding protein subunit 2; 20kD	2.4
20	321023	EOS20954	H25135 Hs.125608 ESTs	2.4
	302099	EOS02030	AL021397 Hs.137576 ribosomal protein L34 pseudogene 1	2.4
	314092	EOS14023	AI984040 Hs.226946 ESTs	2.4
	318587	EOS18518	AA779704 Hs.168830 ESTs	2.4
	303702	EOS03633	AW500748 Hs.224961 ESTs; Weakly similar to 73 kDa subunit of cleavage and polyadenylation specificity factor [H.sapiens]	2.4
	301822	EOS01753	X17033 Hs.1142 integrin; alpha 2 (CD49B); alpha 2 subunit of VLA-2 receptor	2.4
25	322694	EOS22625	AI110872 EST cluster (not in UniGene)	2.4
	323333	EOS23264	AA228883 EST cluster (not in UniGene)	2.4
	301954	EOS01885	AJ099936 Hs.118138 nuclear receptor subfamily 1; group 1; member 2	2.4
	331363	EOS31294	AA421562 Hs.91011 anterior gradient 2 ( <i>Xenopus laevis</i> ) homolog	2.4
	303811	EOS03742	AW182340 Hs.246155 ESTs; Weakly similar to DNA TOPOISOMERASE I [H.sapiens]	2.4
30	308243	EOS08174	AI560037 EST singleton (not in UniGene) with exon hit	2.4
	336021	EOS35952	CH22_3404FG_669_10_LINK_DJ32I10.GENSCAN.9-15 CH22_FGENES.669_10	2.4
	334789	EOS34720	CH22_2101FG_432_14_LINK_EM:AC005500.GENSCAN.293-17 CH22_FGENES.432_14	2.4
35	320807	EOS20738	AA086110 Hs.188536 Homo sapiens clone 24838 mRNA sequence	2.4
	328903	EOS28834	c_8_hs_gi 5868514 refl gn 1 + 23625 24468 ex 3 5 CDSi 91.18 844 219 CH.08_hs_gi 5868514	2.4
	338759	EOS38690	CH22_7581FG__LINK_EM:AC005500.GENSCAN.517-6 CH22_EM:AC005500.GENSCAN.517-6	2.3
40	333769	EOS33700	CH22_1036FG_271_8_LINK_EM:AC005500.GENSCAN.127-8 CH22_FGENES.271_8	2.3
	303597	EOS03528	AI792141 Hs.143560 ESTs; Weakly similar to brain mitochondrial carrier protein-1 [H.sapiens]	2.3
	305898	EOS05829	AA872838 Hs.242463 keratin 8	2.3
45	304439	EOS04370	AA398882 EST singleton (not in UniGene) with exon hit	2.3
	301604	EOS01535	AA373124 Hs.105837 ESTs; Weakly similar to C17G10.1 [C.elegans]	2.3
	315071	EOS15002	AA552690 Hs.152423 ESTs	2.3
	330565	EOS30496	U51095 Hs.1545 caudal type homeo box transcription factor 1	2.3
	331589	EOS31520	N71027 Hs.41856 ESTs	2.3
50	303216	EOS03147	AA581439 Hs.152328 ESTs	2.3
	324988	EOS24919	T06997 EST cluster (not in UniGene)	2.3
	312996	EOS12927	AA249018 EST cluster (not in UniGene)	2.3
	332314	EOS32245	T25862 Hs.101774 ESTs	2.3
	313325	EOS13256	AI420611 Hs.127832 ESTs	2.3
	322991	EOS22922	C18965 Hs.159473 ESTs	2.3
55	335496	EOS35427	CH22_2848FG_571_4_LINK_EM:AC005500.GENSCAN.460-25 CH22_FGENES.571_4	2.3
	315135	EOS15066	AA627561 Hs.192446 ESTs	2.3
	319488	EOS19419	AW250340 EST cluster (not in UniGene)	2.3
60	323571	EOS23502	AA984133 Hs.153260 c-Cbl-interacting protein	2.3
	322826	EOS22757	AI807883 Hs.156932 ESTs	2.3
	322221	EOS22152	AI890619 Hs.179562 nucleosome assembly protein 1-like 1	2.3
	312242	EOS12173	AI380207 Hs.125276 ESTs	2.3
	315238	EOS15169	AA593867 Hs.170890 ESTs	2.3
	315168	EOS15099	AA622130 Hs.152524 ESTs	2.3
65	300504	EOS00435	AW204624 Hs.192927 ESTs; Weakly similar to Lim kinase [H.sapiens]	2.3
	323243	EOS23174	W44372 EST cluster (not in UniGene)	2.3
	331628	EOS31559	R80965 Hs.204079 ESTs	2.3
	320746	EOS20677	AA128302 Hs.156704 EST cluster (not in UniGene)	2.3
70	324598	EOS24529	AA502659 Hs.163986 ESTs	2.3
	308667	EOS08598	AI758754 Hs.152524 EST singleton (not in UniGene) with exon hit	2.2
	302944	EOS20875	AA340708 Hs.256204 ESTs; Weakly similar to cyclic nucleotide-gated channel beta subunit [R.norvegicus]	2.2
	316291	EOS16222	AW375974 Hs.156704 ESTs	2.2
	315296	EOS15227	AA876905 Hs.125286 ESTs	2.2
75	334150	EOS34081	CH22_1429FG_339_1_LINK_EM:AC005500.GENSCAN.189-1 CH22_FGENES.339_1	2.2
	331380	EOS31311	AA453266 Hs.246131 ESTs	2.2
	321795	EOS21726	AI796896 Hs.222446 ESTs	2.2
	331493	EOS31424	N34357 Hs.44571 ESTs	2.2
80	32890	EOS12821	AI813654 Hs.127478 ESTs	2.2
	315583	EOS15514	AW003622 Hs.126555 ESTs	2.2
	314306	EOS14237	AI697901 Hs.192425 ESTs	2.2
	314138	EOS14069	AA740616 EST cluster (not in UniGene)	2.2
	302656	EOS02587	AW293005 Hs.220905 ESTs	2.2
	313564	EOS13495	AA810141 Hs.192182 ESTs	2.2
85	332792	EOS32723	CH22_8FG_3_2_LINK_C4G1.GENSCAN.3-2 CH22_FGENES.3_2	2.2

	332020	EOS31951	AA488895	Hs.105219	ESTs	2.2
5	315143	EOS15074	AA878324	Hs.192734	ESTs	2.2
	313385	EOS13316	AI032087	Hs.176711	ESTs	2.2
	323935	EOS23766	AL042005	EST cluster (not in UniGene)		2.2
	314014	EOS13945	AW291847	Hs.121715	ESTs; Weakly similar to HP protein [H.sapiens]	2.2
	336016	EOS35947	CH22_3399FG_669_5_LINK_DJ3210.GENSCAN.9-10	CH22_FGENES.669_5		2.2
10	323218	EOS23149	AF131846	Hs.13396	Homo sapiens clone 25028 mRNA sequence	2.2
	338059	EOS37990	CH22_6561FG_LINK_EM:AC005500.GENSCAN.160-4	CH22_EM:AC005500.GENSCAN.160-4		2.2
	302613	EOS02544	AA371059	Hs.251636	ubiquitin specific protease 3	2.2
	304852	EOS04783	AA588595	EST singleton (not in UniGene) with exon hit		2.2
	308457	EOS08388	AI669859	EST singleton (not in UniGene) with exon hit		2.2
15	311736	EOS11667	AA765897	EST cluster (not in UniGene)		2.2
	334183	EOS34114	CH22_1464FG_350_13_LINK_EM:AC005500.GENSCAN.209-16	CH22_FGENES.350_13		2.2
	315021	EOS14952	AA533447	EST cluster (not in UniGene)		2.2
	303013	EOS02944	F07898	Hs.214190	interleukin enhancer binding factor 1	2.2
20	315006	EOS14937	AI538613	Hs.135657	ESTs	2.2
	337534	EOS37465	CH22_5803FG_828_3	CH22_FGENES.828-3		2.2
	303276	EOS03207	AA431599	Hs.132799	ESTs	2.1
	318617	EOS18548	AW247252	Hs.75514	nucleoside phosphorylase	2.1
	330760	EOS30691	AA448663	Hs.30469	ESTs	2.1
25	319545	EOS19476	R83716	Hs.14355	ESTs	2.1
	312252	EOS12183	AI128388	Hs.143655	ESTs	2.1
	322882	EOS22813	AW248508	Hs.2491	DiGeorge syndrome critical region gene 2	2.1
	312684	EOS12615	AW294020	Hs.117721	ESTs	2.1
	315782	EOS15713	AW515455	Hs.115558	ESTs; Weakly similar to !!!! ALU SUBFAMILY J WARNING ENTRY !!!! [H.sapiens]	2.1
30	320076	EOS20007	AI653733	Hs.204079	ESTs	2.1
	300566	EOS00497	H86709	Hs.21371	son of sevenless (Drosophila) homolog 1	2.1
	300908	EOS00839	AA618335	Hs.146137	ESTs; Weakly similar to putative [C.elegans]	2.1
	314778	EOS14709	AW079559	Hs.152258	ESTs	2.1
	319233	EOS19164	R21054	Hs.211522	ESTs	2.1
35	335488	EOS35419	CH22_2840FG_570_20_LINK_EM:AC005500.GENSCAN.460-15	CH22_FGENES.570_20		2.1
	334616	EOS34547	CH22_1923FG_411_15_LINK_EM:AC005500.GENSCAN.274-22	CH22_FGENES.411_15		2.1
40	306792	EOS06723	AI042426	EST singleton (not in UniGene) with exon hit		2.1
	301661	EOS01592	AI815558	EST cluster (not in UniGene) with exon hit		2.1
	311332	EOS11263	AW292247	Hs.255052	ESTs	2.1
	314785	EOS14716	AI538226	Hs.135184	ESTs	2.1
	301460	EOS1391	AA196758	Hs.165998	DKFZP564M2423 protein	2.1
	332015	EOS31946	AA487910	Hs.208800	ESTs; Weakly similar to !!! ALU CLASS B WARNING ENTRY !!!! [H.sapiens]	2.1
45	321529	EOS21460	AI269506	Hs.146066	ESTs	2.1
	323740	EOS23671	AA324643	Hs.246106	ESTs	2.1
	336019	EOS35950	CH22_3402FG_669_8_LINK_EM:DJ3210.GENSCAN.9-13	CH22_FGENES.669_8		2.1
	314954	EOS14885	AA521381	Hs.187726	ESTs	2.1
50	303037	EOS02986	AA118395	EST cluster (not in UniGene) with exon hit		2.1
	302056	EOS01987	AI457532	Hs.126082	ESTs; Moderately similar to ROSA26AS [M.musculus]	2.1
	315178	EOS15109	AW362945	Hs.162459	ESTs	2.1
	332246	EOS32177	N57927	Hs.120777	ESTs; Weakly similar to RNA POLYMERASE II ELONGATION FACTOR ELL2 [H.sapiens]	2.0
	334288	EOS34219	CH22_1577FG_369_18_LINK_EM:AC005500.GENSCAN.229-18	CH22_FGENES.369_18		2.0
55	324690	EOS24621	N88286	Hs.132808	ESTs; Weakly similar to Similar to S.pombe -rad4+/cut5+product [H.sapiens]	2.0
	305257	EOS05188	AA679005	EST singleton (not in UniGene) with exon hit		2.0
	311315	EOS11246	AW450536	Hs.209260	ESTs	2.0
	311988	EOS11919	AW016096	Hs.13801	ESTs	2.0
60	302638	EOS02569	AA463798	Hs.102696	ESTs; Weakly similar to C11D2.4 [C.elegans]	2.0
	320531	EOS20462	W03691	Hs.24884	ESTs; Moderately similar to RNA polymerase I associated factor [M.musculus]	2.0
	323604	EOS23535	AI751438	Hs.182827	ESTs; Weakly similar to !!! ALU SUBFAMILY SQ WARNING ENTRY !!!! [H.sapiens]	2.0
	308852	EOS08783	AI829848	Hs.182937	peptidyl/prolyl Isomerase A (cyclophilin A)	2.0
	320521	EOS20452	N31464	Hs.24743	ESTs	2.0
	331306	EOS13237	AA262079	Hs.63931	dachshund (Drosophila) homolog	2.0
65	314941	EOS14872	AA515902	Hs.130650	ESTs	2.0
	336684	EOS36615	CH22_4167FG_46_1	CH22_FGENES.46-1		2.0
	301137	EOS01088	AF049569	Hs.137096	ESTs	2.0
	338454	EOS38385	CH22_7128FG_LINK_EM:AC005500.GENSCAN.360-4	CH22_EM:AC005500.GENSCAN.360-4		2.0
70	309700	EOS09631	AW241170	Hs.179661	Homo sapiens clone 24703 beta-tubulin mRNA; complete cds	2.0
	330262	EOS30193	c_5_p2 gl 6671884 gb A gn 1 + 67913 68053 ex 3 3 CDSI 5.41 141 597	CH.05_p2 gl 6671884		2.0
	324163	EOS24094	AL046827	Hs.134651	ESTs	2.0
75	316493	EOS16424	AA766142	Hs.131810	ESTs; Weakly similar to !!! ALU SUBFAMILY J WARNING ENTRY !!!! [H.sapiens]	2.0
	311873	EOS11804	AA730045	Hs.187866	ESTs	2.0
	326757	EOS26688	c20_hs gl 6249610 ref  gn 3 + 74531 74597 ex 1 3 CDSf 9.52 67 1416	CH.20_hs gl 6249610		2.0
	319167	EOS19098	F05984	Hs.250138	protein phosphatase 2C; magnesium-dependent; catalytic subunit	2.0
80	316011	EOS15942	AW516953	Hs.201372	ESTs	2.0
	313635	EOS13566	AA507227	Hs.6390	ESTs	2.0
	310027	EOS09958	AW449009	Hs.126647	ESTs	2.0
	336662	EOS36593	CH22_4138FG_41_1	CH22_FGENES.41-1		2.0
	334648	EOS34579	CH22_1956FG_417_15_LINK_EM:AC005500.GENSCAN.278-15	CH22_FGENES.417_15		2.0
85	308676	EOS08607	AI761036	EST singleton (not in UniGene) with exon hit		2.0
	312047	EOS11978	AA588275	Hs.14258	ESTs	2.0

	324826	EOS24757	AA704806	Hs.143842	ESTs	2.0
	322889	EOS22820	AA081924	Hs.211417	ESTs	2.0
5	316345	EOS16276	AW139408	Hs.152940	ESTs	2.0
	313922	EOS13853	AI702038	Hs.100057	ESTs	2.0
	319423	EOS19354	T83024	Hs.15119	ESTs	2.0
	320244	EOS20175	AA296922	Hs.129778	gastrointestinal peptide EST singleton (not in UniGene) with exon hit	2.0
	308957	EOS08888	AI869642		EST singleton (not in UniGene) with exon hit	2.0
	334223	EOS34154	CH22_1507FG_360_4_LINK_EM:AC005500.GENSCAN.218-4			
10	302980	EOS02911	W93435		CH22_FGENES.360_4	1.9
	312153	EOS12084	AA759250	Hs.153028	EST cluster (not in UniGene) with exon hit	1.9
	326460	EOS26391	c19_hs_gi 5867400 ref  gn 3 - 142633 142935 ex 1 2 CDSI 19.03 303 1731	cytochrome b-561		1.9
	319962	EOS19893	H06350	Hs.135056	ESTs	1.9
15	307064	EOS06995	AI149335		EST singleton (not in UniGene) with exon hit	1.9
	331608	EOS31539	N89861	Hs.44162	ESTs; Weakly similar to cDNA EST yk342h12.5 comes from this gene [C.elegans]	1.9
	328142	EOS28073	c_6_hs_gi 5868050 ref  gn 1 - 9656 9778 ex 2 6 CDSI 11.11 123 3339			
	319622	EOS19910	AB018281	Hs.107479	CH.06_hs_gi 5868050	1.9
20	312527	EOS12458	AI695522	Hs.191271	ESTs	1.9
	318581	EOS18512	AA769058		EST cluster (not in UniGene)	1.9
	319979	EOS19910	AB018281	Hs.107479	KIAA0738 gene product	1.9
	336107	EOS36038	CH22_3496FG_696_3_LINK_DA59H18.GENSCAN.4-3			
	305232	EOS05163	AA670052	Hs.195188	CH22_FGENES.696_3	1.9
25	315043	EOS14974	AA806538	Hs.130732	glyceraldehyde-3-phosphate dehydrogenase	1.9
	323377	EOS23308	AA133260	Hs.8454	ESTs	1.9
	338260	EOS38191	CH22_6863FG__LINK_EM:AC005500.GENSCAN.279-10	protein kinase; cAMP-dependent; regulatory; type II; alpha		1.9
	334891	EOS34822	CH22_2208FG_452_5_LINK_EM:AC005500.GENSCAN.341-8			
30	316055	EOS15986	AA693880		CH22_FGENES.452_5	1.9
	312414	EOS12345	AI915014	Hs.164235	EST cluster (not in UniGene)	1.9
	300225	EOS00156	AI989963	Hs.197505	ESTs; Weakly similar to !!!! ALU SUBFAMILY J WARNING ENTRY !!!! [H.sapiens]	1.9
35	332607	EOS32538	R41791	Hs.36566	ESTs	1.9
	312405	EOS12336	AI523875		LIM domain kinase 1	1.9
	313605	EOS13536	AI761786	Hs.204674	EST cluster (not in UniGene)	1.9
	337755	EOS37686	CH22_6105FG__LINK_EM:AC000097.GENSCAN.109-2			1.9
40	323216	EOS23147	AA332145		CH22_EM:AC000097.GENSCAN.109-2	1.9
	334872	EOS34803	CH22_2186FG_450_2_LINK_EM:AC005500.GENSCAN.339-2		EST cluster (not in UniGene)	1.9
	332034	EOS31965	AA489847	Hs.112019	CH22_FGENES.450_2	1.9
	332103	EOS32034	AA609161	Hs.112657	ESTs; Moderately similar to !!!! ALU SUBFAMILY J WARNING ENTRY !!!! [H.sapiens]	1.9
45	318196	EOS18127	AI056776	Hs.133397	ESTs; Weakly similar to ORF YOR243c [S.cerevisiae]	1.9
	329141	EOS29072	c_x_hs_gi 6017060 ref  gn 1 + 343924 343997 ex 2 3 CDSI 8.53 74 1715		ESTs; Weakly similar to ORF YOR243c [S.cerevisiae]	1.9
	321539	EOS21470	N98619	Hs.62461	CH.X_hs_gi 6017060	1.9
50	313881	EOS13812	AA535580	Hs.16331	ARP2 (actin-related protein 2; yeast) homolog	1.9
	314046	EOS13977	AW021917	Hs.181878	ESTs	1.9
	336045	EOS35976	CH22_3430FG_679_7_LINK_EM:AC005500.GENSCAN.18-8			1.9
	324799	EOS24730	AW272262	Hs.250468	CH22_FGENES.679_7	1.9
55	312656	EOS12587	AW152449	Hs.226469	ESTs	1.9
	324662	EOS24593	AW504689		EST cluster (not in UniGene)	1.9
	323930	EOS23861	AA570698	Hs.193203	ESTs	1.9
	314465	EOS14396	AA602917	Hs.156974	ESTs	1.9
	335897	EOS35828	CH22_3274FG_635_5_LINK_EM:AC005500.GENSCAN.525-7			1.9
60	321746	EOS21677	AI806500	Hs.102652	CH22_FGENES.635_5	1.9
	335687	EOS35618	CH22_3048FG_596_2_LINK_EM:AC005500.GENSCAN.488-2		ESTs; Weakly similar to KIAA0437 [H.sapiens]	1.9
	330731	EOS30662	AA278816	Hs.177204	CH22_FGENES.596_2	1.9
65	315542	EOS15473	AA079476	Hs.109857	ESTs; Highly similar to CGI-89 protein [H.sapiens]	1.9
	336379	EOS36310	CH22_3791FG_821_7_LINK_EM:BA232E17.GENSCAN.4-19			1.9
	305691	EOS05622	AA813590	Hs.119500	CH22_FGENES.821_7	1.9
	310639	EOS10570	AW269082	Hs.175162	karyopherin alpha 4 (importin alpha 3)	1.9
	327481	EOS27412	c_2_hs_gi 5867783 ref  gn 3 + 104472 104673 ex 1 4 CDSf 14.33 202 1308		ESTs	1.9
70	301910	EOS1841	T84852	Hs.98370	CH22_FGENES.821_7	1.9
	335478	EOS35409	CH22_2830FG_569_1_LINK_EM:AC005500.GENSCAN.456-1		cytochrome P540 family member predicted from ESTs	1.9
	331135	EOS31066	R61398	Hs.4197	CH22_FGENES.569_1	1.9
75	335690	EOS35621	CH22_3051FG_596_5_LINK_EM:AC005500.GENSCAN.488-5		ESTs	1.9
	308047	EOS07978	AI459633		EST singleton (not in UniGene) with exon hit	1.9
	334500	EOS34431	CH22_1800FG_397_16_LINK_EM:AC005500.GENSCAN.260-18			1.9
80	338250	EOS38181	CH22_6848FG__LINK_EM:AC005500.GENSCAN.269-2		CH22_FGENES.397_16	1.9
	320618	EOS20549	AI220276	Hs.235228	CH22_EM:AC005500.GENSCAN.269-2	1.8
	335044	EOS34975	CH22_2367FG_480_1_LINK_EM:AC005500.GENSCAN.374-1		EST	1.8
	313789	EOS13720	AI167810	Hs.217743	CH22_FGENES.480_1	1.8
85	311911	EOS11842	AI087123	Hs.114434	ESTs; Weakly similar to !!!! ALU SUBFAMILY J WARNING ENTRY !!!! [H.sapiens]	1.8
	320180	EOS20111	AA846203	Hs.193974	ESTs; Weakly similar to alternatively spliced product using exon 13A [H.sapiens]	1.8

5	311036	EOS10967	AI539227	Hs.214039	ESTs	1.8
	323903	EOS23834	AA773580	Hs.193598	ESTs	1.8
	318676	EOS18607	T57448	Hs.15467	ESTs; Moderately similar to putative phosphoinositide 5-phosphatase type II [M.musculus]	1.8
	303007	EOS02938	AA478876	Hs.7037	ESTs; pallid (mouse) homolog; pallidin	1.8
	334806	EOS34737	CH22_2119FG_435_7_LINK_EM:AC005500.GENSCAN.296-6			1.8
				CH22_FGENES.435_7		
	311767	EOS11698	AI076686	Hs.190066	ESTs	1.8
10	331750	EOS31681	AA284372	Hs.111471	ESTs	1.8
	314872	EOS14803	AI144254	Hs.239726	ESTs	1.8
	314071	EOS14002	AA192455	Hs.188690	ESTs	1.8
	328450	EOS23831	c_7_hs_gi 5868425 ref	gn 2 - 209192 209321 ex 2 3 CDSi 10.41 130 1407		1.8
				CH.07_hs_gi 5868425		
	328857	EOS28788	c_7_hs_gi 6381927 ref	gn 3 - 80557 81051 ex 1 1 CDSi 41.51 495 6090		1.8
				CH.07_hs_gi 6381927		
15	313781	EOS13712	AA078836		EST cluster (not in UniGene)	1.8
	336953	EOS36884	CH22_4746FG_361_22_		CH22_FGENES.361-22	1.8
	300233	EOS00164	AI380777	Hs.189402	ESTs	1.8
	326862	EOS26793	c20_hs_gi 6552465 ref	gn 2 + 107702 107782 ex 12 13 CDSi 3.62 81 2149		1.8
				CH.20_hs_gi 6552465		
20	312364	EOS12295	R40111	Hs.187618	ESTs	1.8
	321541	EOS21472	AI220292	Hs.254467	ESTs	1.8
	307432	EOS07363	AI244269	Hs.181165	eukaryotic translation elongation factor 1 alpha 1	1.8
	320921	EOS20852	R94038	Hs.199538	inhibin; beta C	1.8
	333110	EOS33041	CH22_338FG_79_16_LINK_EM:AC00097.GENSCAN.59-15			1.8
				CH22_FGENES.79_16		
	324914	EOS24845	AA847510	Hs.161292	ESTs	1.8
	312681	EOS12612	AI028149	Hs.193124	pyruvate dehydrogenase kinase; isoenzyme 3	1.8
	335697	EOS35628	CH22_3058FG_596_12_LINK_EM:AC005500.GENSCAN.488-13			1.8
				CH22_FGENES.596_12		
30	308462	EOS08393	AI671311		EST singleton (not in UniGene) with exon hit	1.8
	312138	EOS12069	T89405	Hs.218851	ESTs; Weakly similar to !!! ALU SUBFAMILY J WARNING ENTRY !!! [H.sapiens]	1.8
	309116	EOS09047	AI927149	Hs.29797	ribosomal protein L10	1.8
	320730	EOS20651	AA534539	Hs.151072	ESTs	1.8
	300844	EOS00775	AL042759	Hs.191762	ESTs	1.8
	337570	EOS37501	CH22_5856FG__LINK_C65E1.GENSCAN.4-2			1.8
				CH22_C65E1.GENSCAN.4-2		
	332756	EOS32687	D63479	Hs.115907	diacylglycerol kinase; delta (130kD)	1.8
	332161	EOS23092	AA621523	Hs.165464	ESTs	1.8
40	300942	EOS00873	AW275006	Hs.195969	ESTs	1.8
	300680	EOS00611	AW468066	Hs.257712	ESTs; Weakly similar to KIAA0986 protein [H.sapiens]	1.8
	328783	EOS28714	c_7_hs_gi 5868309 ref	gn 5 - 73658 73822 ex 2 5 CDSi 0.78 165 5371		1.8
				CH.07_hs_gi 5868309		
	307542	EOS07473	AI280859		EST singleton (not in UniGene) with exon hit	1.8
45	331975	EOS31906	AA464972	Hs.99624	ESTs	1.8
	321532	EOS21463	T77886	Hs.83428	nuclear factor of kappa light polypeptide gene enhancer in B-cells 1 (p105)	1.8
	318721	EOS18652	Z28504		EST cluster (not in UniGene)	1.8
	302124	EOS02055	AB023967	Hs.145078	regulator of differentiation (in S. pombe) 1	1.8
	323541	EOS23472	AI185116	Hs.104613	ESTs; Weakly similar to Similar to S.cerevisiae hypothetical protein L3111 [H.sapiens]	1.8
	331057	EOS30988	N71399	Hs.28143	ESTs	1.8
50	316860	EOS16791	AW139099	Hs.127489	ESTs	1.8
	330601	EOS30532	U90916	Hs.82845	Human clone 23815 mRNA sequence	1.8
	307334	EOS07265	AI214811	Hs.220615	ESTs; Weakly similar to TFI-I protein [H.sapiens]	1.8
	323195	EOS23126	AI064982	Hs.117950	multifunctional polypeptide similar to SAICAR synthetase and AIR carboxylase	1.8
	303856	EOS03787	AA968589	Hs.944	glucose phosphate isomerase	1.8
	321553	EOS21484	H92449	Hs.116406	ESTs	1.8
	332705	EOS32636	T59161	Hs.76293	thymosin; beta 10	1.8
	333139	EOS33070	CH22_368FG_83_16_LINK_EM:AC00097.GENSCAN.67-19			1.8
				CH22_FGENES.83_16		
60	338997	EOS38928	CH22_7881FG__LINK_DA59H18.GENSCAN.8-22			1.8
				CH22_DA59H18.GENSCAN.8-22		
	301509	EOS01440	AI025435	Hs.117532	ESTs	1.8
	314522	EOS14453	AI732331	Hs.187750	ESTs; Moderately similar to !!! ALU CLASS C WARNING ENTRY !!! [H.sapiens]	1.8
	303072	EOS03003	AF157833		EST cluster (not in UniGene) with exon hit	1.8
65	305271	EOS05202	AA679895		EST singleton (not in UniGene) with exon hit	1.8
	335287	EOS35218	CH22_2629FG_526_11_LINK_EM:AC005500.GENSCAN.420-4			1.8
				CH22_FGENES.526_11		
	321286	EOS21217	AI380940		EST cluster (not in UniGene)	1.8
	318740	EOS18671	NM_002543		EST cluster (not in UniGene)	1.8
70	323465	EOS23396	AA287406		EST cluster (not in UniGene)	1.8
	300611	EOS00542	N75450		EST cluster (not in UniGene) with exon hit	1.8
	306235	EOS06166	AA932299		EST cluster (not in UniGene) with exon hit	1.8
	336721	EOS36652	CH22_4244FG_83_17_		EST singleton (not in UniGene) with exon hit	1.8
	311291	EOS11222	A782601	Hs.122684	CH22_FGENES.83-17	1.8
	310247	EOS10178	A124982	Hs.211454	ESTs	1.8
	316556	EOS16495	AI743571	Hs.168799	ESTs; Weakly similar to !!! ALU SUBFAMILY J WARNING ENTRY !!! [H.sapiens]	1.8
	328170	EOS28101	c_6_hs_gi 5868071 ref	gn 1 + 93170 93295 ex 9 9 CDSi 13.31 126 3591		1.8
				CH.06_hs_gi 5868071		
	300909	EOS00840	AW295479	Hs.154903	ESTs; Weakly similar to Abl substrate ena [D.melanogaster]	1.8
80	330869	EOS30800	AA115197	Hs.183702	ESTs	1.8
	311048	EOS10979	AA506952	Hs.210508	ESTs	1.8
	333764	EOS33695	CH22_1031FG_271_3_LINK_EM:AC005500.GENSCAN.127-3			1.8
				CH22_FGENES.271_3		
	338862	EOS38793	CH22_7715FG__LINK_DJ3210.GENSCAN.1-6			1.8
				CH22_DJ3210.GENSCAN.1-6		
85	331467	EOS31398	N22206	Hs.43112	ESTs	1.8
	327742	EOS27673	c_5_hs_gi 5867944 ref	gn 3 - 143307 143512 ex 1 3 CDSi 11.07 206 172		1.8

				CH.05_hs gj 5867944	1.8
5	320955	EOS20886	AL049415	Hs.204290 Homo sapiens mRNA; cDNA DKFZp586N2119 (from clone DKFZp586N2119)	1.8
	323589	EOS23520	AW390054	Hs.192843 ESTs	1.8
	319951	EOS19882	AA307665	Hs.14559 ESTs	1.8
	333763	EOS33694	CH22_1030FG_271_2_LINK_EM:AC005500.GENSCAN.127-2	CH22_FGENES.271_2	1.7
10	331046	EOS30977	N66563	Hs.191358 ESTs	1.7
	320001	EOS19932	AA873350	EST cluster (not in UniGene)	1.7
	316869	EOS16800	AI954880	Hs.134604 ESTs	1.7
	310774	EOS10705	AW134483	Hs.164371 ESTs	1.7
	319379	EOS19310	T91443	Hs.193963 ESTs	1.7
	321549	EOS21480	AA470984	Hs.161947 ESTs	1.7
	308023	EOS00754	AI863068	Hs.222665 ESTs; Weakly similar to putative zinc finger protein NY-REN-34 antigen [H.sapiens]	1.7
	324228	EOS24159	A1798146	Hs.207780 ESTs	1.7
	313902	EOS13833	AI308165	Hs.156242 ESTs	1.7
	308928	EOS08859	A1863908	EST singleton (not in UniGene) with exon hit	1.7
	333770	EOS33701	CH22_1037FG_272_1_LINK_EM:AC005500.GENSCAN.127-10	CH22_FGENES.272_1	1.7
20	316934	EOS16865	AI571647	Hs.146170 ESTs	1.7
	313219	EOS13150	N74924	Hs.182099 ESTs	1.7
	317360	EOS17291	AI125252	Hs.126419 ESTs	1.7
	303530	EOS03461	AI274851	Hs.258744 ESTs	1.7
	334739	EOS34670	CH22_2051FG_424_14_LINK_EM:AC005500.GENSCAN.285-16	CH22_FGENES.424_14	1.7
25	337670	EOS37601	CH22_5996FG__LINK_EM:AC000097.GENSCAN.57-2	CH22_EM:AC000097.GENSCAN.57-2	1.7
	312079	EOS12010	T79745	Hs.189717 ESTs	1.7
	320211	EOS20142	AL039402	Hs.125783 DEME-6 protein	1.7
	316218	EOS16149	AW207642	Hs.174021 ESTs	1.7
	335682	EOS35613	CH22_3043FG_595_2_LINK_EM:AC005500.GENSCAN.487-11	CH22_FGENES.595_2	1.7
30	330696	EOS30627	AA022632	Hs.15825 ESTs	1.7
	314449	EOS14380	AL042667	Hs.225539 ESTs	1.7
	311972	EOS11903	N51511	Hs.188449 ESTs	1.7
	307691	EOS07622	AI318285	Hs.182371 prothymosin; alpha (gene sequence 28)	1.7
	338249	EOS318180	CH22_6847FG__LINK_EM:AC005500.GENSCAN.269-1	CH22_EM:AC005500.GENSCAN.269-1	1.7
	326399	EOS26330	c19_hs gj 5867353 ref	gn 1 + 6385 6536 ex 6 6 CDSI 10.69 152 684 CH.19_hs gj 5867353	1.7
40	313290	EOS13221	AI753247	Hs.206454 ESTs	1.7
	301615	EOS01546	W39477	EST cluster (not in UniGene) with exon hit	1.7
	307034	EOS06965	AI142526	EST singleton (not in UniGene) with exon hit	1.7
	313577	EOS13508	AA565051	Hs.155029 ESTs	1.7
	324703	EOS24634	AB009282	Hs.31086 Homo sapiens mRNA for cytochrome b5; partial cds	1.7
	321317	EOS21248	A1937060	Hs.202040 ESTs; Weakly similar to KIAA0938 protein [H.sapiens]	1.7
	312278	EOS12209	AW205234	Hs.201587 ESTs	1.7
	333358	EOS33289	CH22_604FG_141_9_LINK_EM:AC005500.GENSCAN.21-9	CH22_FGENES.141_9	1.7
50	322735	EOS22666	AA086123	EST cluster (not in UniGene)	1.7
	326752	EOS26683	c20_hs gj 5867615 ref	gn 1 - 1214 1562 ex 2 2 CDSI 33.07 349 1366 CH.20_hs gj 5867615	1.7
	314733	EOS14664	AW452355	Hs.256037 ESTs	1.7
	312902	EOS12833	AW292797	Hs.130316 ESTs	1.7
	322653	EOS22584	AI828654	Hs.171891 ESTs	1.7
	336015	EOS35946	CH22_3398FG_669_4_LINK_EM:J32 10.GENSCAN.9-9	CH22_FGENES.669_4	1.7
	324500	EOS24431	AW269819	Hs.169905 ESTs	1.7
	310900	EOS10831	AI922728	Hs.165803 ESTs; Weakly similar to !!! ALU SUBFAMILY SB WARNING ENTRY !!! [H.sapiens]	1.7
	337908	EOS37839	CH22_6323FG__LINK_EM:AC005500.GENSCAN.57-1	CH22_EM:AC005500.GENSCAN.57-1	1.7
60	304084	EOS04015	T91986	EST singleton (not in UniGene) with exon hit	1.7
	332539	EOS32470	AA412528	Hs.20183 ESTs; Weakly similar to cDNA EST EMBL:T01421 comes from this gene [C.elegans]	1.7
	314332	EOS14263	AL037551	Hs.95612 ESTs	1.7
	321412	EOS21343	AW366305	EST cluster (not in UniGene)	1.7
	321287	EOS12118	AI700439	Hs.188490 ESTs	1.7
	314147	EOS14078	A1656135	Hs.129805 ESTs	1.7
	303131	EOS03062	AW081061	Hs.103180 actin-like 6	1.7
	331341	EOS31272	AA303125	Hs.119009 ESTs; Weakly similar to !!! ALU SUBFAMILY SB2 WARNING ENTRY !!! [H.sapiens]	1.7
	313615	EOS13546	AW295194	Hs.25264 DKFZP434N126 protein	1.7
	329598	EOS29529	c10_p2 gj 3962482 gb	A gn 4 + 39924 40220 ex 2 3 CDSI 8.71 297 420 CH.10_p2 gj 3962482	1.7
	303579	EOS03510	AA381124	Hs.193353 ESTs; Weakly similar to !!! ALU SUBFAMILY J WARNING ENTRY !!! [H.sapiens]	1.7
	331692	EOS31623	W93592	Hs.47343 ESTs	1.7
	323977	EOS23908	AW328177	Hs.234713 ESTs	1.7
	332930	EOS32861	CH22_151FG_38_4_LINK_EM:20H12.GENSCAN.29-4	CH22_FGENES.38_4	1.7
	326596	EOS26527	c19_hs gj 6138928 ref	gn 4 + 133386 133563 ex 7 9 CDSI -1.32 178 3520 CH.19_hs gj 6138928	1.7
80					
	314946	EOS14877	AI097229	Hs.217484 ESTs; Weakly similar to !!! ALU SUBFAMILY J WARNING ENTRY !!! [H.sapiens]	1.7
	315357	EOS15288	AA608684	Hs.121705 ESTs; Moderately similar to !!! ALU CLASS C WARNING ENTRY !!! [H.sapiens]	1.7
	324728	EOS24659	AA303024	EST cluster (not in UniGene)	1.7
	317501	EOS17432	AA931245	Hs.137097 ESTs	1.7
	332219	EOS32150	N22508	Hs.139315 ESTs	1.7
	335369	EOS35300	CH22_2718FG_543_7_LINK_EM:AC005500.GENSCAN.432-9	CH22_FGENES.543_7	1.7
	322417	EOS22348	W36286	Hs.171873 ESTs; Weakly similar to PUTATIVE STEROID DEHYDROGENASE KIK-I [M.musculus]	1.7

5	316100	EOS16031	AW203986	Hs.213003	ESTs	1.7	
	314866	EOS14797	AW305124	Hs.191682	ESTs	1.7	
	300328	EOS00259	AW015600	Hs.224623	ESTs	1.7	
	315676	EOS15607	AW002565	Hs.136590	ESTs	1.7	
	314183	EOS14114	AA748600		EST cluster (not in UniGene)	1.7	
	321354	EOS21285	AA078493		EST cluster (not in UniGene)	1.7	
	311904	EOS11835	T86907	Hs.119371	ESTs	1.7	
	322890	EOS22821	AA082030		EST cluster (not in UniGene)	1.7	
10	302759	EOS02690	AI885815	Hs.184727	ESTs	1.7	
	324600	EOS24531	AA503297	Hs.117108	ESTs	1.7	
	314973	EOS14904	AW273128	Hs.254669	EST	1.7	
	324432	EOS24363	AA464510		EST cluster (not in UniGene)	1.7	
	331520	EOS31451	N49068	Hs.93966	ESTs	1.7	
	308380	EOS08311	A1623988		EST singleton (not in UniGene) with exon hit	1.7	
	331010	EOS30941	H95039	Hs.32168	KIAA0442 protein	1.7	
	325363	EOS25294	c12_hs_gi 5866920 ref  gn 7 + 700446 700516 ex 6 8 CDSi -6.58 71 113 CH.12_hs_gi 5866920			1.7	
	310470	EOS10401	AI281848	Hs.165547	ESTs	1.7	
	330711	EOS30642	AA164687	Hs.177576	mannosyl (alpha-1;3)-glycoprotein beta-1;4-N-acetylglucosaminyltransferase; isoenzyme A	1.7	
20	332074	EOS32005	AA599012	Hs.22826	ESTs	1.7	
	309732	EOS09663	AW262211	Hs.5662	guanine nucleotide binding protein (G protein); beta polypeptide 2-like 1	1.6	
	306337	EOS05268	AA954221	Hs.73742	ribosomal protein; large; P0	1.6	
	335189	EOS35120	CH22_2525FG_507_4_LINK_EM:AC005500.GENSCAN.400-4			1.6	
	316253	EOS16184	AI919537	Hs.118056	ESTs	1.6	
	332908	EOS32839	CH22_129FG_36_12_LINK_C20H12.GENSCAN.28-9			1.6	
	310002	EOS09933	AI439096	Hs.25832	CH22_FGENES.36_12	1.6	
	332258	EOS32189	N68670	Hs.103808	ESTs; Weakly similar to RanBPM [H.sapiens]	1.6	
30	336182	EOS36113	CH22_3576FG_715_2_LINK_DA59H18.GENSCAN.19-3			1.6	
	328987	EOS28918	c_9_hs_gi 5868535 ref  gn 1 - 25705 25764 ex 3 10 CDSi 9.90 60 438 CH.09_hs_gi 5868535			1.6	
	324481	EOS24412	AI916284	Hs.199671	ESTs	1.6	
	331406	EOS31337	AA610064	Hs.23440	KIAA1105 protein	1.6	
	332280	EOS32211	R38100	Hs.106294	ESTs	1.6	
	332173	EOS32104	F09281	Hs.90424	ESTs	1.6	
	335739	EOS35670	CH22_3102FG_601_10_LINK_EM:AC005500.GENSCAN.491-10			1.6	
40	332104	EOS32035	AA609177	Hs.109363	CH22_FGENES.601_10	1.6	
	315033	EOS14964	AI493046	Hs.146133	ESTs	1.6	
	334740	EOS34671	CH22_2052FG_424_15_LINK_EM:AC005500.GENSCAN.285-17			1.6	
	334783	EOS34714	CH22_2095FG_432_8_LINK_EM:AC005500.GENSCAN.293-11			1.6	
	308010	EOS07941	AI439190	Hs.181165	CH22_FGENES.432_8	1.6	
	304521	EOS04452	AA464716		eukaryotic translation elongation factor 1 alpha 1	1.6	
	318719	EOS18650	Z25900	Hs.18724	EST singleton (not in UniGene) with exon hit	1.6	
	321920	EOS21851	N63915		Homo sapiens mRNA; cDNA DKFZp564F093 (from clone DKFZp564F093)	1.6	
	315019	EOS14950	AA532807	Hs.105822	EST cluster (not in UniGene)	1.6	
	320793	EOS20724	AL049980	Hs.184216	ESTs	1.6	
	305371	EOS05302	AA714180		DKFZP564C152 protein	1.6	
	308054	EOS04985	AA634127	Hs.182426	EST singleton (not in UniGene) with exon hit	1.6	
	314643	EOS14574	AI587502	Hs.192088	ribosomal protein S2	1.6	
	308186	EOS08117	AI537940		ESTs	1.6	
	319371	EOS19302	R00321	Hs.174928	EST singleton (not in UniGene) with exon hit	1.6	
	331700	EOS31631	Z40011	Hs.180582	ESTs	1.6	
	316965	EOS16886	AW203959	Hs.149532	ESTs	1.6	
60	314961	EOS14892	AW008061	Hs.231994	ESTs	1.6	
	336676	EOS36607	CH22_4154FG_43_4		CH22_FGENES.43-4	1.6	
	322801	EOS22732	A1831910	Hs.163734	ESTs	1.6	
	303363	EOS03294	AI964095	Hs.226801	ESTs; Weakly similar to DIA-156 protein [H.sapiens]	1.6	
	328105	EOS28036	c_6_hs_gi 5868020 ref  gn 11 - 301705 301784 ex 4 7 CDSi 5.30 80 3147			1.6	
	325481	EOS25412	c12_hs_gi 5866957 ref  gn 3 + 47590 47672 ex 4 7 CDSi 2.69 83 1895 CH.12_hs_gi 5866957			1.6	
	315361	EOS15292	AI335229	Hs.122031	ESTs	1.6	
	324902	EOS24833	D31323	Hs.211188	ESTs	1.6	
	336018	EOS35949	CH22_3401FG_669_7_LINK_DJ32 10.GENSCAN.9-12			1.6	
	308747	EOS08678	AI804500	Hs.181165	CH22_FGENES.669_7	1.6	
	328251	EOS28182	c_6_hs_gi 5868191 ref  gn 4 + 124444 124557 ex 2 3 CDSi 0.40 114 4554 CH.06_hs_gi 5868191			1.6	
	303153	EOS03084	U09759	Hs.8325	mitogen-activated protein kinase 9	1.6	
	327809	EOS27740	c_5_hs_gi 5867968 ref  gn 3 + 54610 54761 ex 4 4 CDSi 0.78 152 993 CH.05_hs_gi 5867968			1.6	
	314107	EOS14038	AA806113	Hs.189025	ESTs	1.6	
	300304	EOS00235	A1637934	Hs.224978	ESTs	1.6	
	313009	EOS12940	W62010	Hs.191379	ESTs	1.6	
80	331074	EOS31005	R08440		yf19f.s1 Soares fetal liver spleen 1NFLS Homo sapiens cDNA clone IMAGE:127337 3' similar to contains Alu repetitive element; mRNA sequence	1.6	
	335773	EOS35704	CH22_3142FG_607_9_LINK_EM:AC005500.GENSCAN.496-4			1.6	
	334991	EOS34922	CH22_2312FG_469_11_LINK_EM:AC005500.GENSCAN.365-11			1.6	
	322959	EOS22890	AI267606		CH22_FGENES.469_11 EST cluster (not in UniGene)	1.6	

	323731	EOS23662	AA323414	EST cluster (not in UniGene)	1.6
	331073	EOS31004	R07998	Hs.18628 ESTs; Weakly similar to !!! ALU SUBFAMILY J WARNING ENTRY !!! [H.sapiens]	1.6
	313573	EOS13504	AI076259	Hs.190337 ESTs	1.6
	316949	EOS16880	AA856749	Hs.124620 ESTs	1.6
5	328084	EOS28015	c_6_hs_gi 6469819 ref  gn 3 - 155366 155459 ex 1 4 CDSI 1.23 94 2982 CH.06_hs_gi 6469819		1.6
	331526	EOS31457	N49967	Hs.46624 ESTs	1.6
	317987	EOS17918	AW138174	Hs.130651 ESTs	1.6
10	325594	EOS25525	c13_hs_gi 5866992 ref  gn 4 - 470474 470566 ex 2 3 CDSI 8.09 93 68 CH.13_hs_gi 5866992		1.6
	310848	EOS10779	AI459554	Hs.161286 ESTs	1.6
	309268	EOS09199	AI985821	Hs.62954 ferritin; heavy polypeptide 1	1.6
	304518	EOS04449	AA461438	EST singleton (not in UniGene) with exon hit	1.6
15	331065	EOS30996	N09584	Hs.9167 Homo sapiens clone 25085 mRNA sequence	1.6
	306501	EOS06432	AA987294	EST singleton (not in UniGene) with exon hit	1.6
	323289	EOS23220	AL134235	Hs.222442 ESTs	1.6
	334630	EOS34561	CH22_1938FG_416_6_LINK_EM:AC005500.GENSCAN.277-6 CH22_FGENES.416_6		1.6
20	302025	EOS01956	AI091466	Hs.127241 DKFZP564F052 protein	1.6
	328998	EOS28929	c_9_hs_gi 5868538 ref  gn 1 + 40996 41104 ex 1 3 CDSI 11.00 109 480 CH.09_hs_gi 5868538		1.6
	313197	EOS13128	AI738851	Hs.222487 ESTs	1.6
	338763	EOS38694	CH22_7585FG__LINK_EM:AC005500.GENSCAN.517-16 CH22_EM:AC005500.GENSCAN.517-16		1.6
25	332247	EOS32178	N58172	Hs.109370 ESTs	1.6
	316724	EOS16655	AA810788	Hs.123337 ESTs	1.6
	303306	EOS03237	AA215297	EST cluster (not in UniGene) with exon hit	1.6
	306336	EOS06267	AA954198	EST singleton (not in UniGene) with exon hit	1.6
	308256	EOS08187	AI565498	EST singleton (not in UniGene) with exon hit	1.6
30	307056	EOS06987	AI148675	EST singleton (not in UniGene) with exon hit	1.6
	321370	EOS21301	AJ227900	EST cluster (not in UniGene)	1.6
	336262	EOS36193	CH22_3661FG_754_9_LINK_DA59H18.GENSCAN.57-11 CH22_FGENES.754_9		1.6
35	335497	EOS35428	CH22_2849FG_571_5_LINK_EM:AC005500.GENSCAN.460-26 CH22_FGENES.571_5		1.6
	309582	EOS09513	AW169657	EST singleton (not in UniGene) with exon hit	1.6
	329563	EOS29494	c10_p2_gi 3962490 gb A gn 1 - 410 635 ex 2 2 CDSI 13.80 226 267 CH.10_p2_gi 3962490		1.6
40	332504	EOS32435	AA053917	Hs.15106 chromosome 14 open reading frame 1	1.6
	308090	EOS08021	AI474601	Hs.2186 eukaryotic translation elongation factor 1 gamma	1.6
	331752	EOS31683	AA287312	Hs.191648 ESTs	1.6
	330881	EOS30812	AA132986	Hs.69321 ESTs; Weakly similar to Similar to mucin and several other Ser-Thr-rich proteins [S.cerevisiae]	1.6
	315647	EOS15578	AA648983	Hs.212911 ESTs	1.6
45	336766	EOS36697	CH22_4341FG_143_20	CH22_FGENES.143-20	1.6
	302592	EOS02523	AA294921	Hs.250811 v-ral simian leukemia viral oncogene homolog B (ras related; GTP binding protein)	1.6
	315076	EOS15007	AI623817	Hs.168457 ESTs	1.6
	337056	EOS36987	CH22_4946FG_441_4	CH22_FGENES.441-4	1.6
	322175	EOS22106	AF085975	EST cluster (not in UniGene)	1.6
	336833	EOS36764	CH22_4504FG_242_2	CH22_FGENES.242-2	1.6
50	334902	EOS34833	CH22_2219FG_452_16_LINK_EM:AC005500.GENSCAN.341-19 CH22_FGENES.452_16		1.6
	318671	EOS18602	AA188823	Hs.212621 ESTs	1.6
	308064	EOS07995	AI469273	Hs.181165 eukaryotic translation elongation factor 1 alpha 1	1.6
	320559	EOS20490	AB021981	Hs.159322 solute carrier family 35 (UDP-N-acetylglucosamine (UDP-GlcNAc) transporter); member 3	1.6
55	317881	EOS17812	AI827248	Hs.224398 ESTs	1.6
	313078	EOS13009	N49730	EST cluster (not in UniGene)	1.6
	338689	EOS38620	CH22_7464FG_527_3	CH22_EM:AC005500.GENSCAN.475-3	1.6
60	311804	EOS11735	AA135159	Hs.203349 ESTs	1.6
	316359	EOS16290	AI472213	Hs.123415 ESTs	1.6
	330182	EOS30113	c_4_p2_gi 5123954 emb  gn 4 + 120156 120245 ex 2 2 CDSI 4.69 90 11 CH.04_p2_gi 5123954		1.6
	334718	EOS34649	CH22_2028FG_421_29_LINK_EM:AC005500.GENSCAN.282-29 CH22_FGENES.421_29		1.6
65	324196	EOS24127	AA405524	Hs.178000 ESTs	1.6
	305350	EOS05281	AA706676	EST singleton (not in UniGene) with exon hit	1.6
	331469	EOS31400	N22273	Hs.39140 ESTs	1.6
	305715	EOS05646	AA826884	EST singleton (not in UniGene) with exon hit	1.6
70	314460	EOS14391	AI263231	Hs.145607 ESTs	1.6
	317634	EOS17565	AA953088	Hs.127550 ESTs	1.6
	335293	EOS35224	CH22_2635FG_527_6_LINK_EM:AC005500.GENSCAN.421-9 CH22_FGENES.527_6		1.6
	305611	EOS05542	AA782331	EST singleton (not in UniGene) with exon hit	1.6
	310430	EOS10361	AI670843	Hs.200257 ESTs	1.6
	323696	EOS23627	AA641201	Hs.222051 ESTs	1.6
	300610	EOS00541	N72596	Hs.99120 DEAD/H (Asp-Glu-Ala-Asp/His) box polypeptide; Y chromosome	1.6
75	327364	EOS27295	c_1_hs_gi 6552412 ref  gn 2 - 115235 115396 ex 1 9 CDSI 2.77 162 3007 CH.01_hs_gi 6552412		1.6
	324848	EOS24779	AW021857	EST cluster (not in UniGene)	1.6
	321491	EOS21422	H70665	Hs.183960 ESTs	1.6
	336367	EOS36298	CH22_3779FG_818_11_LINK_BA232E17.GENSCAN.3-17 CH22_FGENES.818_11		1.6
	331549	EOS31480	N56866	Hs.237507 EST	1.6
80	328332	EOS28263	c_7_hs_gi 5868375 ref  gn 6 + 280154 280289 ex 3 5 CDSI -1.04 136 516 CH.07_hs_gi 5868375		1.5
	322817	EOS22748	C02420	EST cluster (not in UniGene)	1.5

303983	EOS03914	AW514111	Hs.181165	eukaryotic translation elongation factor 1 alpha 1		1.5	
329434	EOS29365	c_y_hs	gi 5868883 ref	gn 1 - 31124 31263 ex 3 20 CDSi 6.38 140 241 CH.Y_hs gi 5868883		1.5	
5	338196	EOS38127	CH22_6763FG	_LINK_EM:AC005500.GENSCAN.235-16	CH22_EM:AC005500.GENSCAN.235-16	1.5	
308488	EOS08419	AI682148	Hs.179661	Homo sapiens clone 24703 beta-tubulin mRNA; complete cds		1.5	
314883	EOS14814	AW178807	Hs.246182	ESTs		1.5	
307095	EOS07026	AI167910		EST singleton (not in UniGene) with exon hit		1.5	
306953	EOS05884	AI124971		EST singleton (not in UniGene) with exon hit		1.5	
331788	EOS31717	AA398539	Hs.97369	EST		1.5	
303509	EOS03440	AW378236	Hs.256050	ESTs		1.5	
324515	EOS24446	AW501686	Hs.163539	ESTs		1.5	
339323	EOS39254	CH22_8284FG	_LINK_EM:AC005500.GENSCAN.23-2	CH22_BA354I12.GENSCAN.23-2	1.5		
15	306563	EOS06494	AA995296	EST singleton (not in UniGene) with exon hit		1.5	
316076	EOS16007	AW297895	Hs.116424	ESTs		1.5	
325622	EOS25553	c14_hs	gi 5867000 ref	gn 2 + 69994 70075 ex 6 8 CDSi 9.40 82 194 CH.14_hs gi 5867000		1.5	
20	309632	EOS09563	AW193261	Hs.156110	Immunoglobulin kappa variable 1D-8	1.5	
314926	EOS14857	AI380838	Hs.124835	ESTs		1.5	
314458	EOS14389	AI217440	Hs.143873	ESTs		1.5	
335219	EOS35150	CH22_2558FG	513_2_LINK_EM:AC005500.GENSCAN.406-2	CH22_FGENES.513_2		1.5	
25	301079	EOS01010	AA305047	Hs.183654	ESTs; Weakly similar to unknown [S.cerevisiae]	1.5	
334122	EOS34053	CH22_1400FG	333_3_LINK_EM:AC005500.GENSCAN.185-27	CH22_FGENES.333_3		1.5	
30	308139	EOS08070	AI494477	EST singleton (not in UniGene) with exon hit		1.5	
317412	EOS17343	AI301528	Hs.132604	ESTs		1.5	
315073	EOS15004	AW452948	Hs.257631	ESTs		1.5	
313139	EOS13070	AA362113		EST cluster (not in UniGene)		1.5	
307012	EOS05943	AI140212		EST singleton (not in UniGene) with exon hit		1.5	
322895	EOS22826	AW470295	Hs.192152	ESTs		1.5	
303779	EOS03710	AA897296	Hs.221266	ESTs		1.5	
35	312344	EOS12275	AI742618	Hs.181733	ESTs; Weakly similar to nitrilase homolog 1 [H.sapiens]	1.5	
323632	EOS23563	AL039950		EST cluster (not in UniGene)		1.5	
332336	EOS32267	T96130	Hs.137551	ESTs		1.5	
304547	EOS04478	AA486189		EST singleton (not in UniGene) with exon hit		1.5	
335692	EOS35623	CH22_3053FG	596_7_LINK_EM:AC005500.GENSCAN.488-7	CH22_FGENES.596_7		1.5	
40	328333	EOS28264	c_7_hs	gi 5868375 ref	gn 6 + 282506 282664 ex 4 5 CDSi 7.71 159 517 CH.07_hs gi 5868375	1.5	
304143	EOS04074	R88737		EST singleton (not in UniGene) with exon hit		1.5	
329625	EOS29556	c11_p2	gi 4567169 gb	A gn 2 - 85893 85984 ex 3 5 CDSi 2.24 92 29 CH.11_p2 gi 4567169		1.5	
45	329960	EOS29891	c16_p2	gi 5091594 gb	A gn 1 - 1031 1162 ex 1 3 CDSi 10.75 132 415 CH.16_p2 gi 5091594	1.5	
318975	EOS18906	Z44110		EST cluster (not in UniGene)		1.5	
321875	EOS21806	N49122		EST cluster (not in UniGene)		1.5	
50	320451	EOS20382	R26944	Hs.180777	Homo sapiens mRNA; cDNA DKFZp564M0264 (from clone DKFZp564M0264)	1.5	
336020	EOS35951	CH22_3403FG	669_9_LINK_EM:DJ32110.GENSCAN.9-14	CH22_FGENES.669_9		1.5	
55	332581	EOS32512	T28799	Hs.2913	EphB3	1.5	
336622	EOS38553	CH22_7384FG	_LINK_EM:AC005500.GENSCAN.451-1	CH22_EM:AC005500.GENSCAN.451-1		1.5	
330397	EOS30328	D14659	Hs.154387	KIAA0103 gene product		1.5	
314359	EOS14290	AA205569	Hs.194193	ESTs		1.5	
313456	EOS13387	AW380579	Hs.209657	ESTs		1.5	
318486	EOS18417	H09123	Hs.139258	ESTs		1.5	
60	318175	EOS18106	AA644624		EST cluster (not in UniGene)	1.5	
335684	EOS35615	CH22_3045FG	595_4_LINK_EM:AC005500.GENSCAN.487-13	CH22_FGENES.595_4		1.5	
65	327814	EOS27745	c_5_hs	gi 5867968 ref	gn 6 + 69377 70566 ex 1 2 CDSi 86.15 1190 999 CH.05_hs gi 5867968	1.5	
322120	EOS22051	W84351	Hs.213846	ESTs		1.5	
311749	EOS11680	R06249	Hs.13911	ESTs		1.5	
329797	EOS29728	c14_p2	gi 6523160 emb	gn 1 - 10616 10894 ex 3 6 CDSi 5.86 279 1549 CH.14_p2 gi 6523160		1.5	
70	330630	EOS30561	X78669	Hs.79088	reticulocalbin 2; EF-hand calcium binding domain	1.5	
303777	EOS03708	AA348491		EST cluster (not in UniGene) with exon hit		1.5	
309656	EOS09587	AW197060	Hs.195188	glyceraldehyde-3-phosphate dehydrogenase		1.5	
326165	EOS26096	c17_hs	gi 5867208 ref	gn 2 - 62787 62929 ex 1 10 CDSi 0.87 143 2037 CH.17_hs gi 5867208		1.5	
75	308328	EOS08259	AI590571	Hs.186412	EST	1.5	
300601	EOS00532	A1762130	Hs.165619	ESTs		1.5	
303610	EOS03541	AA323288		EST cluster (not in UniGene) with exon hit		1.5	
307856	EOS07787	A1366158		EST singleton (not in UniGene) with exon hit		1.5	
319920	EOS19851	R54575	Hs.13337	ESTs; Weakly similar to similar to Phosphoglucomutase and phosphomannomutase phosphoserine [C.elegans]		1.5	
80	332167	EOS32098	D57389	Hs.75447	ralA binding protein 1	1.5	
316427	EOS16358	AI241019	Hs.145644	ESTs		1.5	
303866	EOS03817	AW365963		EST cluster (not in UniGene) with exon hit		1.5	
314292	EOS14223	AA732590	Hs.134740	ESTs		1.5	
315408	EOS15339	AW273261	Hs.216292	ESTs		1.5	
85	335698	EOS35629	CH22_3059FG	597_1_LINK_EM:AC005500.GENSCAN.489-1	CH22_FGENES.597_1		1.5
	315084	EOS15015	AI821085	Hs.187796	ESTs	1.5	

302299	EOS02230	R64632	Hs.182167	hemoglobin; gamma A	1.5	
306803	EOS06734	AI055860	Hs.193717	interleukin 10	1.5	
315802	EOS15733	AA677540	Hs.117064	ESTs	1.5	
326257	EOS26188	c17_hs_gi 5867264 ref	gn 6 + 222712 222819 ex 2 2 CDSI	4.46 108 3597		
			CH.17_hs_gi 5867264		1.5	
5	319599	EOS19530	H56112	EST cluster (not in UniGene)	1.5	
	321891	EOS21822	AW157424	Hs.165954	ESTs	1.5
	335164	EOS35095	CH22_2500FG_502_8_LINK_EM:AC005500.GENSCAN.396-23	CH22_FGENES.502_8	1.5	
10	327133	EOS27064	c21_hs_gi 6682522 ref	gn 1 + 38069 38938 ex 2 2 CDSI	63.42 870 1583	
			CH.21_hs_gi 6682522		1.5	
	317460	EOS17391	AA926980	Hs.131347	ESTs	1.5
	332344	EOS32275	W45574	Hs.252497	ESTs	1.5
15	328801	EOS28732	c_7_hs_gi 58668321 ref	gn 1 - 44492 44609 ex 2 3 CDSI	1.71 118 5525	
			CH.07_hs_gi 58668321		1.5	
	321677	EOS21608	N44545	Hs.251865	ESTs	1.5
	331858	EOS31769	AA421163	Hs.163848	ESTs	1.5
	309243	EOS09174	AI972052	EST singleton (not in UniGene) with exon hit		
20	326213	EOS26144	c17_hs_gi 5867224 ref	gn 3 - 60751 60927 ex 1 4 CDSI	2.06 177 2687	
			CH.17_hs_gi 5867224		1.5	
	321632	EOS21563	AA419617	EST cluster (not in UniGene)	1.5	
	321424	EOS21355	AA057301	EST cluster (not in UniGene)	1.5	
	322465	EOS22396	AA137152	Hs.3784	ESTs; Highly similar to phosphoserine aminotransferase [H.sapiens]	1.5
25	333391	EOS33322	CH22_637FG_144_6_LINK_EM:AC005500.GENSCAN.25-6	CH22_FGENES.144_6	1.5	
	333384	EOS33315	CH22_630FG_143_23_LINK_EM:AC005500.GENSCAN.24-17	CH22_FGENES.143_23	1.5	
	334784	EOS34715	CH22_2096FG_432_9_LINK_EM:AC005500.GENSCAN.293-12	CH22_FGENES.432_9	1.5	
30	334078	EOS34009	CH22_1356FG_327_33_LINK_EM:AC005500.GENSCAN.181-35	CH22_FGENES.327_33	1.5	
	335158	EOS35089	CH22_2494FG_502_2_LINK_EM:AC005500.GENSCAN.396-17	CH22_FGENES.502_2	1.5	
35	335062	EOS34993	CH22_2388FG_482_17_LINK_EM:AC005500.GENSCAN.376-16	CH22_FGENES.482_17	1.5	
	333243	EOS33174	CH22_482FG_111_7_LINK_EM:AC000097.GENSCAN.120-6	CH22_FGENES.111_7	1.5	
	306380	EOS06311	AA968861	EST singleton (not in UniGene) with exon hit	1.5	
40	320809	EOS20740	AI540299	EST cluster (not in UniGene)	1.5	
	332813	EOS32744	CH22_29FG_8_1_LINK_C65E1.GENSCAN.2-2	CH22_FGENES.8_1	1.5	
	335817	EOS35748	CH22_3189FG_618_5_LINK_EM:AC005500.GENSCAN.510-5	CH22_FGENES.618_5	1.5	
45	319551	EOS19482	AA761668	EST cluster (not in UniGene)	1.5	
	334472	EOS34403	CH22_1771FG_394_3_LINK_EM:AC005500.GENSCAN.257-3	CH22_FGENES.394_3	1.5	
	333029	EOS32960	CH22_255FG_68_3_LINK_EM:AC000097.GENSCAN.40-3	CH22_FGENES.68_3	1.5	
50	308055	EOS07986	AI468091	Hs.119252	tumor protein; translationally-controlled 1	1.5
	302882	EOS02813	AW403330	EST cluster (not in UniGene) with exon hit	1.5	
	314033	EOS13964	AA167125	EST cluster (not in UniGene)	1.5	
	324928	EOS24859	AI932285	Hs.160569	ESTs	1.5
	329524	EOS29455	c10_p2_gi 3983507 gb A	gn 6 - 38025 38143 ex 3 3 CDSI	2.40 119 170	
			CH.10_p2_gi 3983507		1.5	
55	333131	EOS33062	CH22_360FG_83_6_LINK_EM:AC000097.GENSCAN.67-10	CH22_FGENES.83_6	1.5	
	332085	EOS32016	AA600353	Hs.173933	ESTs; Weakly similar to NUCLEAR FACTOR 1/X [H.sapiens]	1.5
60	305369	EOS05300	AA714040	EST singleton (not in UniGene) with exon hit	1.5	
	300344	EOS00275	AW291487	Hs.213659	ESTs	1.5
	325071	EOS25002	H08693	EST cluster (not in UniGene)	1.5	
	323693	EOS23624	AW297758	Hs.249721	ESTs	1.5
	321899	EOS21830	N55158	Hs.135252	ESTs	1.5
	331857	EOS31788	AA421160	Hs.9456	SWI/SNF related; matrix associated; actin dependent regulator of chromatin; subfamily a; member 5	1.5
65	334850	EOS34781	CH22_2164FG_439_36_LINK_EM:AC005500.GENSCAN.311-13	CH22_FGENES.439_36	1.5	
	322610	EOS22541	AF180919	EST cluster (not in UniGene)	1.5	
	335332	EOS35263	CH22_2677FG_535_6_LINK_EM:AC005500.GENSCAN.426-6	CH22_FGENES.535_6	1.5	
70	307565	EOS07496	AI282468	EST singleton (not in UniGene) with exon hit	1.5	
	314140	EOS14071	AI216473	Hs.154297	ESTs	1.5
	323011	EOS22942	AA580288	EST cluster (not in UniGene)	1.5	
	325366	EOS25297	c12_hs_gi 5866920 ref	gn 9 - 920962 921713 ex 1 8 CDSI	15.95 752 167	
			CH.12_hs_gi 5866920		1.5	
75	322306	EOS22237	W75935	Hs.146083	ESTs	1.5
	311034	EOS10965	AI564023	Hs.171467	ESTs; Highly similar to NKG2-D TYPE II INTEGRAL MEMBRANE PROTEIN [H.sapiens]	1.5
	305081	EOS05012	AA641638	EST singleton (not in UniGene) with exon hit	1.5	
	322933	EOS22864	AA099759	EST cluster (not in UniGene)	1.5	
	335221	EOS35152	CH22_2560FG_513_4_LINK_EM:AC005500.GENSCAN.406-4	CH22_FGENES.513_4	1.5	
80	304948	EOS04879	AA613107	EST singleton (not in UniGene) with exon hit	1.5	
	334900	EOS34831	CH22_2217FG_452_14_LINK_EM:AC005500.GENSCAN.341-17	CH22_FGENES.452_14	1.5	
	318404	EOS18335	AI654108	Hs.135125	ESTs	1.5
85	339358	EOS39289	CH22_8328FG_LINK_BA354112.GENSCAN.31-3	CH22_BA354112.GENSCAN.31-3	1.5	
	327074	EOS27005	c21_hs_gi 6531965 ref	gn 58 + 4039993 4040096 ex 3 4 CDSI	0.68 104 1284	

			CH.21_hs gi 6531965	1.5	
326054	EOS25985	c17_hs gi 5867184 ref  gn 2 - 146342 146469 ex 3 4 CDSi 10.00 128 426	CH.17_hs gi 5867184	1.5	
5	326892	EOS26823	c20_hs gi 6682511 ref  gn 5 + 119424 119500 ex 29 30 CDSi 18.89 77 2313	CH.20_hs gi 6682511	1.5
328767	EOS28698	c_7_hs gi 6017031 ref  gn 1 - 35625 35723 ex 4 4 CDSf 5.63 99 5262	CH.07_hs gi 6017031	1.5	
10	337772	EOS37703	CH22_6125FG__LINK_EM:AC000097.GENSCAN.119-11	CH22_EM:AC000097.GENSCAN.119-11	1.5
312199	EOS12130	AW438602 Hs.191179	ESTs	1.5	
303506	EOS03437	AA340605 Hs.105887	ESTs	1.5	
325176	EOS25107	T52843	EST cluster (not in UniGene)	1.5	
302023	EOS01954	AF060567 Hs.126782	sushi-repeat protein	1.5	
305833	EOS05764	AA857836 Hs.181165	eukaryotic translation elongation factor 1 alpha 1	1.5	
15	309131	EOS09062	AI929175 Hs.119122	ribosomal protein L13a	1.5
334184	EOS34115	CH22_1465FG_350_15_LINK_EM:AC005500.GENSCAN.209-17	CH22_FGENES_350_15	1.5	
335188	EOS35119	CH22_2524FG_507_3_LINK_EM:AC005500.GENSCAN.400-3	CH22_FGENES_507_3	1.5	
20	304813	EOS04744	AA584540	EST singleton (not in UniGene) with exon hit	1.5
315359	EOS15290	AA608808 Hs.225118	ESTs	1.5	
324434	EOS24365	AA707249 Hs.98789	ESTs	1.5	
327910	EOS27841	c_6_hs gi 5868162 ref  gn 1 + 21622 21748 ex 6 7 CDSi 3.69 127 449	CH.06_hs gi 5868162	1.4	
25	335671	EOS35602	CH22_3031FG_592_3_LINK_EM:AC005500.GENSCAN.485-4	CH22_FGENES_592_3	1.4
334943	EOS34874	CH22_2264FG_465_8_LINK_EM:AC005500.GENSCAN.359-8	CH22_FGENES_465_8	1.4	
30	326393	EOS26324	c19_hs gi 5867341 ref  gn 2 + 41702 41841 ex 5 5 CDSi 20.15 140 504	CH.19_hs gi 5867341	1.4
305296	EOS05227	AA687181	EST singleton (not in UniGene) with exon hit	1.4	
307243	EOS07174	AI199957	EST singleton (not in UniGene) with exon hit	1.4	
320066	EOS19997	AW364885 Hs.112442	ESTs	1.4	
311465	EOS11396	AI758660 Hs.206132	ESTs	1.4	
35	302822	EOS02753	AW404176 Hs.111611	ribosomal protein L27	1.4
304987	EOS04918	AA618044	EST singleton (not in UniGene) with exon hit	1.4	
330892	EOS30823	AA149579 Hs.118258	ESTs	1.4	
333385	EOS33316	CH22_631FG_143_24_LINK_EM:AC005500.GENSCAN.24-18	CH22_FGENES_143_24	1.4	
40	302626	EOS02557	AB021870	EST cluster (not in UniGene) with exon hit	1.4
318042	EOS17973	AW294522 Hs.149991	ESTs	1.4	
339361	EOS39292	CH22_8331FG__LINK_EM:BA354112.GENSCAN.32-3	CH22_BA354112.GENSCAN.32-3	1.4	
45	309000	EOS08931	AI880489	EST singleton (not in UniGene) with exon hit	1.4
306004	EOS05935	AA889992	EST singleton (not in UniGene) with exon hit	1.4	
329539	EOS29470	c10_p2_gi 3983503 gb U gn 1 - 1 326 ex 1 3 CDSi 41.66 326 212	CH.10_p2_gi 3983503	1.4	
50	313663	EOS13594	AI953261 Hs.169813	ESTs	1.4
323538	EOS23469	AW247696	EST cluster (not in UniGene)	1.4	
337595	EOS37526	CH22_5884FG__LINK_EM:C20H12.GENSCAN.8-1	CH22_C20H12.GENSCAN.8-1	1.4	
55	303149	EOS03080	AA312995	EST cluster (not in UniGene) with exon hit	1.4
308484	EOS08415	AI679292	EST singleton (not in UniGene) with exon hit	1.4	
300912	EOS00843	AW138724 Hs.168974	ESTs	1.4	
315158	EOS15089	AA744438 Hs.142476	ESTs; Weakly similar to !!! ALU CLASS D WARNING ENTRY !!! [H.sapiens]	1.4	
300462	EOS00393	AA746501 Hs.14217	ESTs	1.4	
312730	EOS12661	AI804372 Hs.208661	ESTs	1.4	
316868	EOS16799	AI660898 Hs.195602	ESTs	1.4	
60	337629	EOS37560	CH22_5933FG__LINK_EM:C20H12.GENSCAN.28-35	CH22_C20H12.GENSCAN.28-35	1.4
332518	EOS32449	D16562 Hs.155433	ATP synthase; H+ transporting; mitochondrial F1 complex; gamma polypeptide 1	1.4	
337422	EOS37353	CH22_5624FG_760_2	CH22_FGENES_760-2	1.4	
328835	EOS28766	c_7_hs gi 5868339 ref  gn 5 + 88053 88461 ex 3 3 CDSi 13.78 409 5775	CH.07_hs gi 5868339	1.4	
65	338282	EOS38213	CH22_6897FG__LINK_EM:AC005500.GENSCAN.291-4	CH22_EM:AC005500.GENSCAN.291-4	1.4
337895	EOS37826	CH22_6303FG__LINK_EM:AC005500.GENSCAN.56-2	CH22_EM:AC005500.GENSCAN.56-2	1.4	
70	320330	EOS20261	AF026004 Hs.141660	chloride channel 2	1.4
314302	EOS14233	AA813118 Hs.163230	ESTs	1.4	
313280	EOS13211	AI285537 Hs.222830	ESTs	1.4	
333222	EOS33153	CH22_459FG_105_2_LINK_EM:AC000097.GENSCAN.109-6	CH22_FGENES_105_2	1.4	
75	305726	EOS05657	AA828156	EST singleton (not in UniGene) with exon hit	1.4
312674	EOS12605	AI762475 Hs.151327	ESTs; Moderately similar to !!! ALU SUBFAMILY J WARNING ENTRY !!! [H.sapiens]	1.4	
315869	EOS15800	AI033547 Hs.132826	ESTs	1.4	
327010	EOS26941	c21_hs gi 5867664 ref  gn 12 + 941057 941139 ex 9 9 CDSi 7.44 83 790	CH.21_hs gi 5867664	1.4	
80	325892	EOS25823	c16_hs gi 5867088 ref  gn 1 - 10498 10652 ex 2 3 CDSi 3.94 155 870	CH.16_hs gi 5867088	1.4
302575	EOS02506	AF071164 Hs.249171	homeo box A11	1.4	
301970	EOS01901	AB028962 Hs.120245	KIAA1039 protein	1.4	
332207	EOS32138	H61475 Hs.237353	EST	1.4	
316024	EOS15955	AA707141 Hs.193388	ESTs	1.4	
314599	EOS14530	AW206512 Hs.186996	ESTs	1.4	
85	333585	EOS33516	CH22_846FG_203_4_LINK_EM:AC005500.GENSCAN.74-6		1.4

			CH22_FGENES.203_4	1.4		
324670	EOS24601	AI525557	EST cluster (not in UniGene)	1.4		
321307	EOS21238	R85409	EST cluster (not in UniGene)	1.4		
335170	EOS35101	CH22_2506FG_503_1_LINK_EM:AC005500.GENSCAN.397-1				
5			CH22_FGENES.503_1	1.4		
328274	EOS28205	c_7_hs_gi 5868219 ref  gn 2 - 31244 31439 ex 1 11 CDSi 13.06 196 9				
			CH.07_hs_gi 5868219	1.4		
336880	EOS36811	CH22_4619FG_318_8_-	CH22_FGENES.318-8	1.4		
10	313825	EOS13756	AA215470	EST cluster (not in UniGene)	1.4	
318410	EOS18341	AI138418	Hs.144935	ESTs	1.4	
335361	EOS35292	CH22_2710FG_541_11_LINK_EM:AC005500.GENSCAN.431-16				
			CH22_FGENES.541_11	1.4		
319802	EOS19733	AI701489	Hs.202501	ESTs	1.4	
15	334769	EOS34700	CH22_2081FG_429_4_LINK_EM:AC005500.GENSCAN.290-9			
			CH22_FGENES.429_4	1.4		
312709	EOS12640	AW069181	Hs.141146	ESTs; Weakly similar to transformation-related protein [H.sapiens]	1.4	
330004	EOS29935	c16_p2_gi 6623963 gb A gn 5 - 78872 78999 ex 2 6 CDSi 19.93 128 728				
			CH.16_p2_gi 6623963	1.4		
20	313103	EOS13034	AI184303	Hs.143806	ESTs	1.4
326359	EOS26290	c18_hs_gi 5867293 ref  gn 1 + 9436 9494 ex 2 3 CDSi 2.16 59 88				
			CH.18_hs_gi 5867293	1.4		
305211	EOS05142	AA668563	EST singleton (not in UniGene) with exon hit	1.4		
334628	EOS34559	CH22_1936FG_416_4_LINK_EM:AC005500.GENSCAN.277-4				
			CH22_FGENES.416_4	1.4		
25	326919	EOS26850	c21_hs_gi 6456782 ref  gn 2 - 40496 41046 ex 1 5 CDSi 17.70 561 157			
			CH.21_hs_gi 6456782	1.4		
315527	EOS15458	AI791138	Hs.116768	ESTs	1.4	
306090	EOS06021	AA0806209	EST singleton (not in UniGene) with exon hit	1.4		
303316	EOS03247	AF033122	Hs.14125	p53 regulated PA26 nuclear protein	1.4	
303642	EOS03573	AW299459	EST cluster (not in UniGene) with exon hit	1.4		
314357	EOS14288	AA781795	Hs.122587	ESTs	1.4	
337102	EOS37033	CH22_5033FG_472_-7	CH22_FGENES.472-7	1.4		
304384	EOS04315	AA235482	Hs.62954	ferritin; heavy polypeptide 1	1.4	
315117	EOS15048	AA828609	Hs.192044	ESTs	1.4	
35	305750	EOS05681	AA835250	EST singleton (not in UniGene) with exon hit	1.4	
311726	EOS11657	AW081766	Hs.253920	ESTs	1.4	
326996	EOS26927	c21_hs_gi 5867660 ref  gn 4 - 63212 63404 ex 2 6 CDSi 15.70 193 622				
			CH.21_hs_gi 5867660	1.4		
40	330257	EOS30188	c_5_p2_gi 6671881 gb A gn 2 - 143228 143393 ex 1 9 CDSi 11.31 166 586			
			CH.05_p2_gi 6671881	1.4		
323864	EOS23795	AA340724	Hs.214028	ESTs	1.4	
338204	EOS38135	CH22_6773FG__LINK_EM:AC005500.GENSCAN.241-3				
			CH22_EM:AC005500.GENSCAN.241-3	1.4		
45	314025	EOS13956	AI983981	Hs.189114	ESTs	1.4
315974	EOS15905	AW029203	Hs.191952	ESTs	1.4	
335599	EOS35530	CH22_2957FG_581_39_LINK_EM:AC005500.GENSCAN.476-37				
			CH22_FGENES.581_39	1.4		
50	335364	EOS35295	CH22_2713FG_543_2_LINK_EM:AC005500.GENSCAN.432-4			
			CH22_FGENES.543_2	1.4		
303634	EOS03565	AI953377	Hs.169425	ESTs; Weakly similar to predicted using Genefinder [C.elegans]	1.4	
315626	EOS15557	AA808598	Hs.35353	ESTs; Weakly similar to H21P03.2 [C.elegans]	1.4	
329936	EOS29867	c16_p2_gi 6165200 gb A gn 4 - 82761 82920 ex 3 4 CDSi 1.15 160 199				
			CH.16_p2_gi 6165200	1.4		
55	328632	EOS28563	c_7_hs_gi 5868247 ref  gn 1 + 76734 76853 ex 1 4 CDSf 13.95 120 3764			
			CH.07_hs_gi 5868247	1.4		
330207	EOS30138	c_5_p2_gi 6013606 gb A gn 3 - 109912 110004 ex 2 4 CDSi 6.54 93 174				
			CH.05_p2_gi 6013606	1.4		
60	329919	EOS29850	c16_p2_gi 6223624 gb A gn 6 - 103492 103681 ex 1 8 CDSi 6.18 190 93			
			CH.16_p2_gi 6223624	1.4		
331916	EOS31847	AA446131	Hs.124918	ESTs	1.4	
317617	EOS17548	T58194	EST cluster (not in UniGene)	1.4		
331943	EOS31874	AA453418	Hs.178272	ESTs	1.4	
306413	EOS06344	AA973288	EST singleton (not in UniGene) with exon hit	1.4		
313607	EOS13538	N94169	Hs.194258	ESTs; Moderately similar to !!! ALU SUBFAMILY SC WARNING ENTRY !!! [H.sapiens]	1.4	
65	336292	EOS36223	CH22_3691FG_783_3_LINK_EM:BA35412.GENSCAN.4-7			
			CH22_FGENES.783_3	1.4		
70	330453	EOS30384	HG3976-HT4246	Pou-Domain Dna Binding Factor Pit1, Pituitary-Specific	1.4	
324602	EOS24533	AA503620	Hs.212329	ESTs	1.4	
332183	EOS32114	H08225	Hs.177181	ESTs	1.4	
320032	EOS19963	AI699772	Hs.202361	ESTs; Weakly similar to X-linked retinopathy protein [H.sapiens]	1.4	
333156	EOS33087	CH22_387FG_89_6_LINK_EM:AC000097.GENSCAN.84-8				
			CH22_FGENES.89_6	1.4		
75	334156	EOS34087	CH22_1435FG_340_6_LINK_EM:AC005500.GENSCAN.190-7			
			CH22_FGENES.340_6	1.4		
334303	EOS34234	CH22_1594FG_373_6_LINK_EM:AC005500.GENSCAN.233-5				
			CH22_FGENES.373_6	1.4		
325513	EOS25444	c12_hs_gi 6017035 ref  gn 1 - 34295 34490 ex 2 7 CDSi 6.49 196 2471				
			CH.12_hs_gi 6017035	1.4		
80	302758	EOS02689	AA984563	EST cluster (not in UniGene) with exon hit	1.4	
329557	EOS29488	c10_p2_gi 3962492 gb A gn 6 - 53197 53647 ex 2 2 CDSf 37.68 451 247				
			CH.10_p2_gi 3962492	1.4		
85	331717	EOS31648	AA190888	Hs.153881 ESTs; Highly similar to NY-REN-62 antigen [H.sapiens]	1.4	
325885	EOS25816	c16_hs_gi 5867087 ref  gn 11 + 193212 193377 ex 1 3 CDSf 43.19 166 792				
			CH.16_hs_gi 5867087	1.4		
85	312160	EOS12091	AA805903	Hs.184371 ESTs	1.4	
328882	EOS28813	c_7_hs_gi 6552423 ref  gn 2 - 157669 157826 ex 4 6 CDSi 4.91 158 6200				

				CH.07_hs gi 6552423	1.4
339028	EOS38959	CH22_7925FG__LINK_DA59H18.GENSCAN.22-8		CH22_DA59H18.GENSCAN.22-8	1.4
5	323497	EOS23428	AI523613 Hs.221544	ESTs	1.4
	316897	EOS16828	AA838114	EST cluster (not in UniGene)	1.4
	312479	EOS12410	AI950844 Hs.128738	ESTs; Weakly similar to non-lens beta gamma-crystallin like protein [H.sapiens]	1.4
	338535	EOS38466	CH22_7251FG__LINK_EM:AC005500.GENSCAN.404-3	CH22_EM:AC005500.GENSCAN.404-3	1.4
10	312754	EOS12685	R99834 Hs.250383	ESTs	1.4
	327527	EOS27458	c_2_hs gi 6381882 ref  gn 2 - 98950 99040 ex 4 8 CDSi 5.78 91 1768	CH.02_hs gi 6381882	1.4
	324714	EOS24645	AA574312 Hs.245737	ESTs	1.4
	302347	EOS02278	AF039400 Hs.194659	chloride channel; calcium activated; family member 1	1.4
15	338008	EOS37939	CH22_6490FG__LINK_EM:AC005500.GENSCAN.127-9	CH22_EM:AC005500.GENSCAN.127-9	1.4
	315590	EOS15521	AA640637 Hs.225817	ESTs	1.4
	320825	EOS20756	NM_004751	EST cluster (not in UniGene)	1.4
	300930	EOS00861	AI289481 Hs.136371	ESTs	1.4
20	335225	EOS35156	CH22_2564FG_513_10_LINK_EM:AC005500.GENSCAN.406-9	CH22_FGENES.513_10	1.4
	337303	EOS37234	CH22_5442FG_681_5	CH22_FGENES.681-5	1.4
	317198	EOS17129	AI810384 Hs.128025	ESTs	1.4
	308991	EOS08922	AI879831	EST singleton (not in UniGene) with exon hit	1.4
25	325472	EOS25403	c12_hs gi 6017034 ref  gn 7 - 289581 289657 ex 2 6 CDSi 4.74 77 1786	CH.12_hs gi 6017034	1.4
	301266	EOS01197	AA829774	EST cluster (not in UniGene) with exon hit	1.4
	330901	EOS30832	AA157818 Hs.238380	Human endogenous retroviral protease mRNA; complete cds	1.4
	313406	EOS13337	AI248314 Hs.132932	ESTs	1.4
30	301454	EOS01385	AI751738	EST cluster (not in UniGene) with exon hit	1.4
	317269	EOS17200	AA906411 Hs.127378	ESTs	1.4
	338876	EOS38807	CH22_7733FG__LINK_DJ32I10.GENSCAN.4-2	CH22_DJ32I10.GENSCAN.4-2	1.4
	328481	EOS28412	c_7_hs gi 5868449 ref  gn 1 - 8987 9180 ex 4 31 CDSi 10.00 194 2103	CH.07_hs gi 5868449	1.4
35	314022	EOS13953	AW452420 Hs.248678	ESTs	1.4
	307640	EOS07571	AI301992	EST singleton (not in UniGene) with exon hit	1.4
	315541	EOS15472	AI168233 Hs.123159	ESTs; Weakly similar to KIAA0668 protein [H.sapiens]	1.4
	315489	EOS15420	AA628245 Hs.191847	ESTs	1.4
40	327815	EOS27746	c_5_hs gi 5867968 ref  gn 6 + 70804 71401 ex 2 2 CDSi 27.99 598 1000	CH.05_hs gi 5867968	1.4
	339319	EOS39250	CH22_8280FG__LINK_BA354I12.GENSCAN.22-19	CH22_BA354I12.GENSCAN.22-19	1.4
	322564	EOS22495	W86440 Hs.118344	ESTs	1.4
45	323812	EOS23743	AW081373 Hs.199199	ESTs	1.4
	303540	EOS03471	AA355607 Hs.173590	ESTs; Weakly similar to MMSET type I [H.sapiens]	1.4
	337902	EOS37833	CH22_6314FG__LINK_EM:AC005500.GENSCAN.56-13	CH22_EM:AC005500.GENSCAN.56-13	1.4
	335289	EOS35220	CH22_2631FG_527_2_LINK_EM:AC005500.GENSCAN.421-2	CH22_FGENES.527_2	1.4
50	327919	EOS27850	c_6_hs gi 5868165 ref  gn 6 + 547701 547800 ex 14 14 CDSi -0.20 100 505	CH.06_hs gi 5868165	1.4
	337674	EOS37605	CH22_6005FG__LINK_EM:AC000097.GENSCAN.67-4	CH22_EM:AC000097.GENSCAN.67-4	1.4
55	320087	EOS20018	AF032387 Hs.113265	small nuclear RNA activating complex; polypeptide 4; 190KD	1.4
	334939	EOS34870	CH22_2259FG_465_3_LINK_EM:AC005500.GENSCAN.359-3	CH22_FGENES.465_3	1.4
	303443	EOS03374	AA320525	EST cluster (not in UniGene) with exon hit	1.3
	325929	EOS25860	c16_hs gi 5867125 ref  gn 2 - 51715 51996 ex 1 1 CDSi 29.05 282 1594	CH.16_hs gi 5867125	1.3
60	327745	EOS27676	c_5_hs gi 6531959 ref  gn 1 - 229066 229124 ex 3 6 CDSi 3.01 59 177	CH.05_hs gi 6531959	1.3
	335166	EOS35097	CH22_2502FG_502_10_LINK_EM:AC005500.GENSCAN.396-25	CH22_FGENES.502_10	1.3
65	324497	EOS24428	AW152624 Hs.136340	ESTs	1.3
	338374	EOS38305	CH22_7017FG__LINK_EM:AC005500.GENSCAN.327-1	CH22_EM:AC005500.GENSCAN.327-1	1.3
	313601	EOS13532	R32458 Hs.257711	ESTs	1.3
	321415	EOS21346	AI377596 Hs.3337	transmembrane 4 superfamily member 1	1.3
70	305309	EOS05240	AA699717	EST singleton (not in UniGene) with exon hit	1.3
	330447	EOS30378	HG3546-HT3744	Pre-Mrna Splicing Factor Sf2p33, Alt. Splice Form 1	1.3
	308578	EOS08509	AI708573	EST singleton (not in UniGene) with exon hit	1.3
	315344	EOS15275	AW292176 Hs.245834	ESTs	1.3
	330503	EOS30434	M55024	Human cell surface glycoprotein P3.58 mRNA, partial cds	1.3
	308227	EOS01858	AI559126 Hs.195188	glyceraldehyde-3-phosphate dehydrogenase	1.3
75	332222	EOS32153	N28271 Hs.176618	ESTs	1.3
	323961	EOS23892	AL044428 Hs.207345	ESTs	1.3
	314530	EOS14461	AI052358 Hs.131741	ESTs	1.3
	320503	EOS20434	NM_005897	EST cluster (not in UniGene)	1.3
	306820	EOS06751	AI074408	EST singleton (not in UniGene) with exon hit	1.3
80	304165	EOS04096	H73265	EST singleton (not in UniGene) with exon hit	1.3
	324302	EOS24233	AA543008 Hs.136806	ESTs; Weakly similar to !!! ALU SUBFAMILY J WARNING ENTRY !!! [H.sapiens]	1.3
	319128	EOS19059	AA393820	EST cluster (not in UniGene)	1.3
	317092	EOS17023	AI286162 Hs.125657	ESTs	1.3
	304998	EOS04929	AA621203	EST singleton (not in UniGene) with exon hit	1.3
85	331433	EOS31364	H68097 Hs.161023	EST	1.3
	333348	EOS33279	CH22_594FG_140_2_LINK_EM:AC005500.GENSCAN.20-2	CH22_EM:AC005500.GENSCAN.20-2	1.3

			CH22_FGENES.140_2	1.3	
333619	EOS33550	CH22_880FG_219_3_LINK_EM:AC005500.GENSCAN.87-2	CH22_FGENES.219_3	1.3	
5	335903	EOS35834	CH22_3280FG_635_11_LINK_EM:AC005500.GENSCAN.525-14	CH22_FGENES.635_11	1.3
	326219	EOS26150	c17_hs gi 5867226 ref  gn 11 - 264008 264274 ex 3 5 CDSi 5.74 267 2847	CH.17_hs gi 5867226	1.3
	324456	EOS24387	AW500954	EST cluster (not in UniGene)	1.3
10	316405	EOS16336	AA757900 Hs.202624	ESTs	1.3
	314361	EOS14292	AL038765 Hs.161304	ESTs	1.3
	328546	EOS28477	c_7_hs gi 5868487 ref  gn 1 - 17547 17722 ex 2 3 CDSi 9.96 176 3284	CH.07_hs gi 5868487	1.3
	335871	EOS35802	CH22_3246FG_629_19_LINK_EM:AC005500.GENSCAN.519-18	CH22_FGENES.629_19	1.3
15	303735	EOS03666	AA707750 Hs.202616	ESTs; Weakly similar to cis-Golgi matrix protein GM130 [R.norvegicus]	1.3
	324048	EOS23979	AA378739	EST cluster (not in UniGene)	1.3
	326720	EOS26651	c20_hs gi 6552456 ref  gn 1 + 84526 84677 ex 5 7 CDSi 11.78 153 1031	CH.20_hs gi 6552456	1.3
20	322309	EOS22240	AF086372	EST cluster (not in UniGene)	1.3
	322136	EOS22067	AF075083	EST cluster (not in UniGene)	1.3
	313460	EOS13391	AW028655 Hs.136033	ESTs	1.3
	306275	EOS05206	AA936312	EST singleton (not in UniGene) with exon hit	1.3
	321974	EOS21905	N76794	EST cluster (not in UniGene)	1.3
25	327600	EOS27531	c_3_hs gi 6004462 ref  gn 1 - 2621 2862 ex 1 4 CDSi -4.01 242 1407	CH.03_hs gi 6004462	1.3
	329086	EOS29017	c_x_hs gi 5868604 ref  gn 1 - 35489 35588 ex 2 9 CDSi 2.55 100 719	CH.X_hs gi 5868604	1.3
	336919	EOS36850	CH22_4690FG_346_6	CH22_FGENES.346-6	1.3
30	302767	EOS02698	H94900 Hs.17882	ESTs	1.3
	334786	EOS34717	CH22_2098FG_432_11_LINK_EM:AC005500.GENSCAN.293-14	CH22_FGENES.432_11	1.3
	302472	EOS02403	AA317451 Hs.241451	SWI/SNF related; matrix associated; actin dependent regulator of chromatin; subfamily e; member 1	1.3
	333033	EOS32964	CH22_259FG_68_8_LINK_EM:AC000097.GENSCAN.40-8	CH22_FGENES.68_8	1.3
35	330493	EOS30424	M27826 Hs.238380	Human endogenous retroviral protease mRNA; complete cds	1.3
	330506	EOS30437	M61906 Hs.6241	phosphoinositide-3-kinase; regulatory subunit; polypeptide 1 (p85 alpha)	1.3
	313932	EOS13863	AI147601 Hs.154087	ESTs	1.3
	314394	EOS14325	AI380563 Hs.130816	ESTs	1.3
	323033	EOS22964	AI744284 Hs.221727	ESTs	1.3
40	326431	EOS26362	c19_hs gi 5867371 ref  gn 1 + 15855 15971 ex 4 6 CDSi 7.79 117 1108	CH.19_hs gi 5867371	1.3
	335547	EOS35478	CH22_2902FG_576_8_LINK_EM:AC005500.GENSCAN.467-8	CH22_FGENES.576_8	1.3
45	300548	EOS00479	AI026836 Hs.114689	ESTs	1.3
	316504	EOS16435	AW135854 Hs.132458	ESTs	1.3
	335756	EOS35687	CH22_3123FG_604_5_LINK_EM:AC005500.GENSCAN.493-10	CH22_FGENES.604_5	1.3
	301209	EOS01140	AI809912 Hs.159354	ESTs	1.3
50	306610	EOS06541	AI000635	EST singleton (not in UniGene) with exon hit	1.3
	314439	EOS14370	AI539443 Hs.137447	ESTs	1.3
	315396	EOS15327	AW296107 Hs.152686	ESTs	1.3
	335914	EOS35845	CH22_3291FG_636_10_LINK_EM:AC005500.GENSCAN.526-10	CH22_FGENES.636_10	1.3
55	333734	EOS33665	CH22_1000FG_260_2_LINK_EM:AC005500.GENSCAN.119-7	CH22_FGENES.260_2	1.3
	312370	EOS12301	AA744692 Hs.166539	ESTs	1.3
	304636	EOS04567	AA524031	EST singleton (not in UniGene) with exon hit	1.3
	323166	EOS23097	AA291001	EST cluster (not in UniGene)	1.3
60	338702	EOS38633	CH22_7482FG__LINK_EM:AC005500.GENSCAN.480-1	CH22_EM:AC005500.GENSCAN.480-1	1.3
	322331	EOS22262	AF086467	EST cluster (not in UniGene)	1.3
	318706	EOS18637	AI383593 Hs.159148	ESTs	1.3
	331186	EOS31117	T41159 Hs.8418	ESTs	1.3
65	334764	EOS34695	CH22_2076FG_428_13_LINK_EM:AC005500.GENSCAN.289-13	CH22_FGENES.428_13	1.3
	327565	EOS27496	c_3_hs gi 5867811 ref  gn 1 + 32516 32778 ex 2 3 CDSi 0.20 263 368	CH.03_hs gi 5867811	1.3
	335524	EOS35455	CH22_2879FG_572_4_LINK_EM:AC005500.GENSCAN.461-4	CH22_FGENES.572_4	1.3
70	308050	EOS07981	AI460004	EST singleton (not in UniGene) with exon hit	1.3
	334172	EOS34103	CH22_1452FG_349_5_LINK_EM:AC005500.GENSCAN.208-6	CH22_FGENES.349_5	1.3
	315674	EOS15605	AA651923 Hs.191850	ESTs	1.3
75	334876	EOS34807	CH22_2190FG_450_6_LINK_EM:AC005500.GENSCAN.339-6	CH22_FGENES.450_6	1.3
	315606	EOS15537	AW298724 Hs.202639	ESTs	1.3
	338779	EOS38710	CH22_7610FG__LINK_EM:AC005500.GENSCAN.526-15	CH22_EM:AC005500.GENSCAN.526-15	1.3
80	333511	EOS33442	CH22_766FG_171_5_LINK_EM:AC005500.GENSCAN.51-5	CH22_FGENES.171_5	1.3
	329254	EOS29185	c_x_hs gi 5868733 ref  gn 1 + 4133 4214 ex 1 2 CDSi -0.36 82 2833	CH.X_hs gi 5868733	1.3
	319510	EOS19441	W88633 Hs.254562	ESTs	1.3
85	339418	EOS39349	CH22_8411FG__LINK_DJ579N16.GENSCAN.11-4	CH22_DJ579N16.GENSCAN.11-4	1.3
	321012	EOS20943	AA737314	EST cluster (not in UniGene)	1.3

333217	EOS33148	CH22_454FG_104_9_LINK_EM:AC000097.GENSCAN.108-8 CH22_FGENES.104_9		1.3
338561	EOS38492	CH22_7294FG__LINK_EM:AC005500.GENSCAN.421-5 CH22_EM:AC005500.GENSCAN.421-5		1.3
5	335742	EOS35673	CH22_3105FG_601_13_LINK_EM:AC005500.GENSCAN.491-14 CH22_FGENES.601_13	1.3
	334993	EOS34924	CH22_2314FG_469_14_LINK_EM:AC005500.GENSCAN.365-16 CH22_FGENES.469_14	1.3
10	323430	EOS23361	AW062479 EST cluster (not in UniGene)	1.3
	306069	EOS06000	AA906983 EST singleton (not in UniGene) with exon hit	1.3
	331681	EOS31612	W85712 Hs.119571 collagen; type III; alpha 1 (Ehlers-Danlos syndrome type IV; autosomal dominant)	1.3
	337986	EOS37917	CH22_6441FG__LINK_EM:AC005500.GENSCAN.110-7 CH22_EM:AC005500.GENSCAN.110-7	1.3
15	313204	EOS13135	AI800518 Hs.118158 ESTs	1.3
	323189	EOS23120	AL121194 Hs.120589 ESTs	1.3
	318171	EOS18102	AA381202 EST cluster (not in UniGene)	1.3
	307156	EOS07087	AI186762 EST singleton (not in UniGene) with exon hit	1.3
20	332713	EOS32644	AA349792 Hs.78489 mutY (E. coli) homolog	1.3
	312828	EOS12759	AI865455 Hs.211818 ESTs; Moderately similar to !!! ALU SUBFAMILY J WARNING ENTRY !!! [H.sapiens]	1.3
	301127	EOS01058	AA758109 Hs.121072 ESTs	1.3
	311260	EOS11191	AI672509 Hs.196582 ESTs	1.3
	338364	EOS38295	CH22_7007FG__LINK_EM:AC005500.GENSCAN.323-7 CH22_EM:AC005500.GENSCAN.323-7	1.3
25	337904	EOS37835	CH22_6318FG__LINK_EM:AC005500.GENSCAN.56-17 CH22_EM:AC005500.GENSCAN.56-17	1.3
	329347	EOS29278	c_x_hs gi 6456785[refl] gn 1 + 18433 18997 ex 4 4 CDSi 43.39 465 3718 CH.X_hs gi 6456785	1.3
30	313329	EOS13260	AW293704 Hs.122658 ESTs	1.3
	314367	EOS14298	AA535749 EST cluster (not in UniGene)	1.3
	317098	EOS17029	AI123513 Hs.125456 ESTs	1.3
	306462	EOS05393	AA983397 EST singleton (not in UniGene) with exon hit	1.3
	301254	EOS01185	AI049624 EST cluster (not in UniGene) with exon hit	1.3
	335504	EOS35435	CH22_2856FG_571_15_LINK_EM:AC005500.GENSCAN.460-34 CH22_FGENES.571_15	1.3
35	334270	EOS34201	CH22_1559FG_368_2_LINK_EM:AC005500.GENSCAN.228-3 CH22_FGENES.368_2	1.3
	334324	EOS34255	CH22_1616FG_375_1_LINK_EM:AC005500.GENSCAN.235-1 CH22_FGENES.375_1	1.3
40	304254	EOS04185	AA046273 Hs.111334 ferritin; light polypeptide	1.3
	305731	EOS05662	AA829363 EST singleton (not in UniGene) with exon hit	1.3
	323284	EOS23215	AA279381 Hs.190010 ESTs	1.3
	322007	EOS21938	AW410646 Hs.165739 ESTs	1.3
	334537	EOS34468	CH22_1839FG_403_2_LINK_EM:AC005500.GENSCAN.268-2 CH22_FGENES.403_2	1.3
45	302360	EOS02291	AJ010901 Hs.198267 mucin 4; tracheobronchial	1.3
	311641	EOS11572	AI948829 Hs.213786 ESTs	1.3
	324643	EOS24574	AI436356 Hs.130729 ESTs	1.3
	327554	EOS27485	c_3_hs gi 5867801[refl] gn 2 - 23092 23191 ex 2 6 CDSi 10.44 100 107 CH.03_hs gi 5867801	1.3
50	312165	EOS12096	AW292139 Hs.115789 ESTs	1.3
	304679	EOS04610	AA548741 EST singleton (not in UniGene) with exon hit	1.3
	319564	EOS19495	AA026777 Hs.169732 ESTs	1.3
	310860	EOS10791	AW015920 Hs.161359 ESTs	1.3
55	337161	EOS37092	CH22_5180FG_561_3 CH22_FGENES.561-3	1.3
	311155	EOS11086	AI634410 Hs.197608 EST	1.3
	336846	EOS36777	CH22_4540FG_263_5 CH22_FGENES.263-5	1.3
	310985	EOS10916	T51842 EST cluster (not in UniGene)	1.3
	329499	EOS29430	c10_p2 gi 3983518 gb A gn 5 + 33463 33789 ex 1 1 CDSo 34.50 327 97 CH.10_p2 gi 3983518	1.3
60	334924	EOS34855	CH22_2244FG_459_2_LINK_EM:AC005500.GENSCAN.351-2 CH22_FGENES.459_2	1.3
	330861	EOS30792	AA084064 Hs.185747 ESTs	1.3
	324658	EOS24589	AI694767 Hs.129179 ESTs	1.3
65	323362	EOS23293	AL135067 Hs.117182 ESTs	1.3
	330468	EOS30399	L10343 Hs.112341 protease inhibitor 3; skin-derived (SKALP)	1.3
	314198	EOS14129	AA897581 Hs.128773 ESTs	1.3
	339436	EOS39367	CH22_8431FG__LINK_EM:DJ579N16.GENSCAN.19-1 CH22_DJ579N16.GENSCAN.19-1	1.3
70	312483	EOS12414	AI417526 Hs.184636 ESTs	1.3
	321505	EOS21436	H73183 Hs.129885 ESTs	1.3
	332254	EOS32185	N64702 Hs.194140 ESTs	1.3
	328253	EOS28184	c_6_hs gi 6381894[refl] gn 1 - 4411 4509 ex 1 5 CDSi 4.20 99 4561 CH.06_hs gi 6381894	1.3
75	332357	EOS32288	W73417 Hs.103183 EST	1.3
	329017	EOS28948	c_x_hs gi 6682532[refl] gn 7 - 255591 255672 ex 3 3 CDSf 12.94 82 22 CH.X_hs gi 6682532	1.3
	337504	EOS37435	CH22_5739FG_803_2 CH22_FGENES.803-2	1.3
	316625	EOS16556	AA780307 Hs.122156 ESTs	1.3
80	335389	EOS35320	CH22_2739FG_545_1_LINK_EM:AC005500.GENSCAN.436-1 CH22_FGENES.545_1	1.3
	310017	EOS09948	AI188739 Hs.148488 ESTs	1.3
	314354	EOS14285	AI037984 Hs.208982 ESTs; Weakly similar to !!! ALU SUBFAMILY J WARNING ENTRY !!! [H.sapiens]	1.3
	324641	EOS24572	AJ732615 Hs.189218 ESTs	1.3
85	335207	EOS35138	CH22_2546FG_510_4_LINK_EM:AC005500.GENSCAN.402-3 CH22_FGENES.510_4	1.3
	333673	EOS33604	CH22_934FG_246_5_LINK_EM:AC005500.GENSCAN.101-3	1.3

			CH22_FGENES.246_5	1.3
334370	EOS34301	CH22_1664FG_378_18_LINK_EM:AC005500.GENSCAN.240-1		1.3
		CH22_FGENES.378_18		1.3
5	328890	EOS28621 c_7_hs gi 6588001[ref] gn 7 - 571207 571274 ex 1 3 CDSI 3.34 68 4325	CH.07_hs gi 6588001	1.3
	323208	EOS23139 AA203415 Hs.136200 ESTs		1.3
	307010	EOS06941 AI140014 EST singleton (not in UniGene) with exon hit		1.3
	316563	EOS16494 AI587083 Hs.200558 ESTs; Weakly similar to !!!! ALU SUBFAMILY SP WARNING ENTRY !!!! [H.sapiens]		1.3
10	312219	EOS12150 H73505 Hs.117874 ESTs		1.3
	319884	EOS19815 T73234 EST cluster (not in UniGene)		1.3
	334720	EOS34651 CH22_2030FG_421_31_LINK_EM:AC005500.GENSCAN.282-31	CH22_FGENES.421_31	1.3
	335836	EOS35767 CH22_3210FG_621_3_LINK_EM:AC005500.GENSCAN.513-3	CH22_FGENES.621_3	1.3
15	305448	EOS05379 AA737894 Hs.29797 ribosomal protein L10		1.3
	314885	EOS14816 AI049878 Hs.133032 ESTs		1.3
	320130	EOS20061 AI820675 Hs.203804 ESTs		1.3
	310567	EOS10498 AI691065 Hs.155780 ESTs		1.3
20	323898	EOS23829 AA347566 EST cluster (not in UniGene)		1.3
	336132	EOS36063 CH22_3522FG_703_2_LINK_EM:AC005500.GENSCAN.9-2	CH22_FGENES.703_2	1.3
	337958	EOS37889 CH22_6403FG__LINK_EM:AC005500.GENSCAN.98-6	CH22_EM:AC005500.GENSCAN.98-6	1.3
25	305630	EOS05561 AA804508 EST singleton (not in UniGene) with exon hit		1.3
	334916	EOS34847 CH22_2235FG_457_7_LINK_EM:AC005500.GENSCAN.347-1	CH22_FGENES.457_7	1.3
	333542	EOS33473 CH22_799FG_178_4_LINK_EM:AC005500.GENSCAN.59-4	CH22_FGENES.178_4	1.3
30	331151	EOS31082 R82331 Hs.164599 ESTs		1.3
	315095	EOS15026 AA831815 Hs.243788 ESTs		1.3
	331593	EOS31524 N72150 Hs.50193 EST		1.3
	323767	EOS23698 AI807408 Hs.166368 ESTs		1.3
	334561	EOS34492 CH22_1865FG_405_1_LINK_EM:AC005500.GENSCAN.270-5	CH22_FGENES.405_1	1.3
35	308191	EOS08122 AI538878 EST singleton (not in UniGene) with exon hit		1.3
	319571	EOS19502 N91399 Hs.220826 ESTs		1.3
	316200	EOS16131 AI914535 Hs.221377 ESTs		1.3
	305996	EOS05927 AA889338 Hs.163356 EST		1.2
40	318055	EOS17986 AI249193 Hs.145945 ESTs		1.2
	315570	EOS15501 AI860360 Hs.160316 ESTs		1.2
	320792	EOS20723 AW236504 Hs.247020 ESTs		1.2
	331649	EOS31580 W20364 Hs.55412 ESTs; Weakly similar to c29 [M.musculus]		1.2
	303839	EOS03770 Z45939 EST cluster (not in UniGene) with exon hit		1.2
45	324399	EOS24330 AA814768 Hs.21396 ESTs		1.2
	317172	EOS17103 AI741232 Hs.206744 ESTs		1.2
	312452	EOS12383 AI692643 Hs.172749 ESTs		1.2
	325482	EOS25413 c12_hs gi 5866957[ref] gn 3 + 47957 48078 ex 5 7 CDSI 10.25 122 1896	CH.12_hs gi 5866957	1.2
50	3111395	EOS11326 R23313 EST cluster (not in UniGene)		1.2
	336124	EOS36055 CH22_3513FG_701_9_LINK_EM:AC005500.GENSCAN.8-9	CH22_FGENES.701_9	1.2
	320082	EOS20013 AA487678 Hs.189738 ESTs		1.2
	312168	EOS12099 T92251 Hs.198882 ESTs		1.2
55	338000	EOS37931 CH22_6472FG__LINK_EM:AC005500.GENSCAN.119-5	CH22_EM:AC005500.GENSCAN.119-5	1.2
	338852	EOS38783 CH22_7705FG__LINK_DJ246D7.GENSCAN.12-1	CH22_DJ246D7.GENSCAN.12-1	1.2
	312090	EOS12021 N57692 Hs.118064 ESTs		1.2
60	316480	EOS16411 AI749921 Hs.205377 ESTs		1.2
	333259	EOS33190 CH22_500FG_118_7__LINK_EM:AC005500.GENSCAN.2-7	CH22_FGENES.118_7	1.2
	335211	EOS35142 CH22_2550FG_511_2__LINK_EM:AC005500.GENSCAN.403-2	CH22_FGENES.511_2	1.2
65	321950	EOS21881 AA594780 Hs.172318 ESTs		1.2
	337937	EOS37868 CH22_6370FG__LINK_EM:AC005500.GENSCAN.86-1	CH22_EM:AC005500.GENSCAN.86-1	1.2
	316576	EOS16507 AI732114 Hs.193046 ESTs; Weakly similar to !!!! ALU SUBFAMILY J WARNING ENTRY !!!! [H.sapiens]		1.2
	322770	EOS22701 AA045796 Hs.159971 SW/SNF related; matrix associated; actin dependent regulator of chromatin; subfamily b; member 1		1.2
70	329369	EOS29300 c_x_hs gi 5868842[ref] gn 1 - 121148 121516 ex 3 4 CDSI 8.50 369 3910	CH.X_hs gi 5868842	1.2
	304183	EOS04114 H91161 EST singleton (not in UniGene) with exon hit		1.2
	339370	EOS39301 CH22_8343FG__LINK_BA232E17.GENSCAN.1-12	CH22_BA232E17.GENSCAN.1-12	1.2
75	303941	EOS03872 AW473878 Hs.156110 Immunoglobulin kappa variable 1D-8		1.2
	302245	EOS02176 H18835 EST cluster (not in UniGene) with exon hit		1.2
	335255	EOS35186 CH22_2597FG_517_2__LINK_EM:AC005500.GENSCAN.411-2	CH22_FGENES.517_2	1.2
80	316610	EOS16541 AW087973 Hs.126731 ESTs		1.2
	314915	EOS14846 AA573072 Hs.187748 ESTs; Weakly similar to !!!! ALU SUBFAMILY J WARNING ENTRY !!!! [H.sapiens]		1.2
	315426	EOS15357 AI391486 Hs.128171 ESTs		1.2
	334003	EOS33934 CH22_1281FG_310_28__LINK_EM:AC005500.GENSCAN.167-28	CH22_FGENES.310_28	1.2
85	304350	EOS04281 AA186871 EST singleton (not in UniGene) with exon hit		1.2
	325173	EOS25104 AI133215 Hs.144662 ESTs; Moderately similar to !!!! ALU SUBFAMILY J WARNING ENTRY !!!! [H.sapiens]		1.2
	312313	EOS12244 AW293341 Hs.122505 ESTs		1.2
	333366	EOS33297 CH22_612FG_142_3__LINK_EM:AC005500.GENSCAN.22-26		1.2

			CH22_FGENES.142_3	1.2	
334970	EOS34901	CH22_2291FG_466_3_LINK_EM:AC005500.GENSCAN.361-2	CH22_FGENES.466_3	1.2	
5	338668	EOS38599	CH22_7441FG__LINK_EM:AC005500.GENSCAN.465-1	CH22_EM:AC005500.GENSCAN.465-1	1.2
336502	EOS36433	CH22_3926FG_833_8_LINK_DJ579N16.GENSCAN.5-9	CH22_FGENES.833_8	1.2	
10	309438	EOS09369	AW102802 Hs.225787 ESTs; Moderately similar to hypothetical protein [H.sapiens]		1.2
336194	EOS36125	CH22_3591FG_717_20_LINK_DA59H18.GENSCAN.20-19	CH22_FGENES.717_20	1.2	
336678	EOS36609	CH22_4156FG_43_6_	CH22_FGENES.43-6	1.2	
321401	EOS21332	W90406 Hs.35962 ESTs		1.2	
306026	EOS05957	AA902309 EST singleton (not in UniGene) with exon hit		1.2	
15	336434	EOS36365	CH22_3854FG_826_1_LINK_BA232E17.GENSCAN.8-1	CH22_FGENES.826_1	1.2
315257	EOS15188	AW157431 Hs.248941 ESTs		1.2	
328349	EOS28280	c_7_hs gil5868383[ref] gn 7 - 260704 260804 ex 2 9 CDSI 4.37 101 621	CH.07_hs gil5868383	1.2	
20	326112	EOS26043	c17_hs gil5867192[ref] gn 1 + 2151 2725 ex 1 1 CDSI 54.87 575 1272	CH.17_hs gil5867192	1.2
333995	EOS33926	CH22_1272FG_310_19_LINK_EM:AC005500.GENSCAN.167-18	CH22_FGENES.310_19	1.2	
323683	EOS23614	AI380045 Hs.225033 ESTs		1.2	
330143	EOS30074	c21_p2 gil4210430[emb] gn 3 + 184737 184848 ex 4 4 CDSI 1.71 112 111	CH.21_p2 gil4210430	1.2	
25	329789	EOS29720	c14_p2 gil6469354[emb] gn 2 - 118977 119036 ex 1 3 CDSI 1.19 60 1517	CH.14_p2 gil6469354	1.2
324397	EOS24328	AA307836 Hs.118758 ESTs; Weakly similar to RLF [H.sapiens]		1.2	
308729	EOS08660	AI799766 Hs.208627 ESTs		1.2	
323939	EOS23870	AW499632 Hs.115696 ESTs		1.2	
30	333444	EOS33375	CH22_694FG_153_1_LINK_EM:AC005500.GENSCAN.34-1	CH22_FGENES.153_1	1.2
306302	EOS06233	AA937901 EST singleton (not in UniGene) with exon hit		1.2	
313693	EOS13624	AW469180 Hs.170651 ESTs		1.2	
316652	EOS16583	AA789249 EST cluster (not in UniGene)		1.2	
332325	EOS32256	T79428 Hs.191264 ESTs		1.2	
336235	EOS36166	CH22_3633FG_740_2_LINK_DA59H18.GENSCAN.44-2	CH22_FGENES.740_2	1.2	
40	319436	EOS19367	R02750 EST cluster (not in UniGene)		1.2
312335	EOS12266	AW043620 Hs.236993 ESTs		1.2	
322109	EOS22040	AI884327 Hs.244737 ESTs		1.2	
328466	EOS28397	c_7_hs gil5868434[ref] gn 1 - 15543 15900 ex 1 2 CDSI 2.36 258 1608	CH.07_hs gil5868434	1.2	
45	323244	EOS23175	T70731 EST cluster (not in UniGene)		1.2
312510	EOS12441	AA779907 Hs.117558 ESTs		1.2	
314853	EOS14784	AA729232 Hs.153279 ESTs		1.2	
336946	EOS36877	CH22_4731FG_355_2_ ESTs	CH22_FGENES.355-2	1.2	
50	303874	EOS03805	AA258921 EST cluster (not in UniGene) with exon hit		1.2
312658	EOS12589	AA730280 Hs.120936 ESTs		1.2	
308354	EOS08285	AI611044 EST singleton (not in UniGene) with exon hit		1.2	
310073	EOS10004	AI335004 Hs.148558 ESTs		1.2	
324777	EOS24708	AA744046 Hs.133350 ESTs		1.2	
300897	EOS00828	AI890356 Hs.127804 ESTs		1.2	
55	308371	EOS08302	AI620666 Hs.242510 EST		1.2
306358	EOS06289	AA961821 EST singleton (not in UniGene) with exon hit		1.2	
312295	EOS12226	AA578233 Hs.173863 ESTs		1.2	
319792	EOS19723	R20317 Hs.22968 ESTs		1.2	
338546	EOS38477	CH22_7267FG__LINK_EM:AC005500.GENSCAN.410-1	CH22_EM:AC005500.GENSCAN.410-1	1.2	
60	314546	EOS14477	AW007211 Hs.186672 ESTs		1.2
338494	EOS38425	CH22_7184FG__LINK_EM:AC005500.GENSCAN.385-5	CH22_EM:AC005500.GENSCAN.385-5	1.2	
65	331131	EOS31062	R54797 Hs.26238 EST; Weakly similar to reverse transcriptase homolog [H.sapiens]		1.2
309939	EOS09870	AW419122 EST singleton (not in UniGene) with exon hit		1.2	
332932	EOS32863	CH22_153FG_38_6_LINK_C20H12.GENSCAN.29-6	CH22_FGENES.38_6	1.2	
70	309653	EOS09584	AW196800 Hs.180842 ribosomal protein L13		1.2
318647	EOS18578	AI526152 EST cluster (not in UniGene)		1.2	
304044	EOS03975	T52479 Hs.252259 ribosomal protein S3		1.2	
75	330307	EOS30238	c_7_p2 gil4877982[gb]A gn 2 + 107384 107559 ex 2 4 CDSI 9.96 176 4	CH.07_p2 gil4877982	1.2
314499	EOS14430	AL044570 Hs.147975 ESTs		1.2	
338053	EOS37984	CH22_6552FG__LINK_EM:AC005500.GENSCAN.158-1	CH22_EM:AC005500.GENSCAN.158-1	1.2	
75	332991	EOS32922	CH22_215FG_56_4_LINK_EM:AC000097.GENSCAN.17-4	CH22_FGENES.56_4	1.2
306308	EOS06239	AA946870 EST singleton (not in UniGene) with exon hit		1.2	
338120	EOS38051	CH22_6655FG__LINK_EM:AC005500.GENSCAN.195-1	CH22_EM:AC005500.GENSCAN.195-1	1.2	
80	313703	EOS13634	AI161293 Hs.146862 ESTs; Weakly similar to KIAA0525 protein [H.sapiens]		1.2
330563	EOS30494	U50553 Hs.147916 DEAD/H (Asp-Glu-Ala-Asp/His) box polypeptide 3		1.2	
332886	EOS32817	CH22_106FG_33_7_LINK_C20H12.GENSCAN.22-9	CH22_FGENES.33_7	1.2	
85	303844	EOS03775	U94362 Hs.58589 glycogenin 2		1.2
321755	EOS21686	AI215881 Hs.144042 ESTs		1.2	
333532	EOS33463	CH22_789FG_175_19_LINK_EM:AC005500.GENSCAN.53-25	CH22_EM:AC005500.GENSCAN.53-25	1.2	

			CH22_FGENES.175_19	1.2
332863	EOS32794	CH22_81FG_28_3_LINK_C20H12.GENSCAN.18-3		1.2
		CH22_FGENES.28_3		1.2
5	33254	EOS33185	CH22_495FG_118_2_LINK_EM:AC005500.GENSCAN.2-2	1.2
		CH22_FGENES.118_2		1.2
317459	EOS17390	AI367254 Hs.131248 ESTs		1.2
315353	EOS15284	AW452608 Hs.129817 ESTs		1.2
300732	EOS00663	AI369956 Hs.257891 ESTs		1.2
10	303502	EOS03433	AA488528 EST cluster (not in UniGene) with exon hit	1.2
	333126	EOS33057	CH22_355FG_82_3_LINK_EM:AC000097.GENSCAN.66-10	1.2
		CH22_FGENES.82_3		1.2
15	332929	EOS32860	CH22_150FG_38_3_LINK_C20H12.GENSCAN.29-3	1.2
		CH22_FGENES.38_3		1.2
329502	EOS29433	c10_p2 gi 3983517 gb U gn 1 + 75 338 ex 1 1 CDSs 46.82 264 100		1.2
		CH.10_p2 gi 3983517		1.2
15	333408	EOS33339	CH22_657FG_145_6_LINK_EM:AC005500.GENSCAN.26-6	1.2
		CH22_FGENES.145_6		1.2
20	315472	EOS15403	AA828850 Hs.165469 ESTs	1.2
	328290	EOS28221	c_7_hs gi 5868363 ref  gn 2 - 127366 127496 ex 1 5 CDSI 5.24 131 289	1.2
		CH.07_hs gi 5868363		1.2
20	328662	EOS28593	c_7_hs gi 6004473 ref  gn 22 + 1184773 1184855 ex 7 8 CDSI 12.72 83 3916	1.2
		CH.07_hs gi 6004473		1.2
25	319808	EOS19739	T58960 EST cluster (not in UniGene)	1.2
	303929	EOS03860	AW470753 EST singleton (not in UniGene) with exon hit	1.2
	315712	EOS15643	AI950133 Hs.120882 ESTs; Moderately similar to !!! ALU SUBFAMILY J WARNING ENTRY !!! [H.sapiens]	1.2
	307391	EOS07322	AI225058 EST singleton (not in UniGene) with exon hit	1.2
	335499	EOS35430	CH22_2851FG_571_8_LINK_EM:AC005500.GENSCAN.460-28	1.2
		CH22_FGENES.571_8		1.2
30	303792	EOS03723	C75094 Hs.199839 ESTs; Highly similar to NG22 [H.sapiens]	1.2
	327287	EOS27218	c_1_hs gi 5867479 ref  gn 1 - 62838 63024 ex 4 5 CDSI 11.66 187 1628	1.2
		CH.01_hs gi 5867479		1.2
30	317713	EOS17644	AI733306 Hs.128071 ESTs	1.2
	330137	EOS30068	c21_p2 gi 4210430 emb  gn 1 - 21220 21377 ex 2 3 CDSI 1.89 158 104	1.2
		CH.21_p2 gi 4210430		1.2
35	308157	EOS08088	AI510824 Hs.75968 thymosin; beta 4; X chromosome	1.2
	314452	EOS14383	AI042699 Hs.209222 ESTs	1.2
	308268	EOS08199	AI567509 Hs.172928 collagen; type I; alpha 1	1.2
	321467	EOS21398	X13075 EST cluster (not in UniGene)	1.2
40	320993	EOS20924	AL050145 Hs.225986 Homo sapiens mRNA; cDNA DKFZp586C2020 (from clone DKFZp586C2020)	1.2
	336778	EOS36709	CH22_4367FG_159_4_- CH22_FGENES.159-4	1.2
	319827	EOS19758	T62778 EST cluster (not in UniGene)	1.2
	308249	EOS08180	AI560998 EST singleton (not in UniGene) with exon hit	1.2
	310094	EOS10025	AW450967 Hs.235240 ESTs	1.2
45	336902	EOS36833	CH22_4655FG_331_2_- CH22_FGENES.331-2	1.2
	339044	EOS38975	CH22_7944FG__LINK_DA59H18.GENSCAN.27-5	1.2
		CH22_DA59H18.GENSCAN.27-5		1.2
	336675	EOS36606	CH22_4153FG_43_3_- CH22_FGENES.43-3	1.2
	303563	EOS03494	AA367699 Hs.118787 transforming growth factor; beta-induced; 68kD	1.2
50	330673	EOS30604	D57823 Hs.92962 Sec23 (S. cerevisiae) homolog A	1.2
	311814	EOS11745	AW377113 Hs.119640 ESTs; Moderately similar to zinc finger protein [H.sapiens]	1.2
	335481	EOS35412	CH22_2833FG_570_10_LINK_EM:AC005500.GENSCAN.460-4	1.2
		CH22_FGENES.570_10		1.2
55	314775	EOS14706	AI149880 Hs.188809 ESTs	1.2
	324961	EOS24892	AA613792 EST cluster (not in UniGene)	1.2
	313458	EOS13389	AA007259 Hs.255853 ESTs	1.2
	307074	EOS07005	AI150989 EST singleton (not in UniGene) with exon hit	1.2
	337964	EOS37895	CH22_6410FG__LINK_EM:AC005500.GENSCAN.100-9	1.2
		CH22_EM:AC005500.GENSCAN.100-9		1.2
60	326519	EOS26450	c19_hs gi 5867439 ref  gn 4 + 166004 166243 ex 4 5 CDSI 4.49 240 2534	1.2
		CH.19_hs gi 5867439		1.2
	337366	EOS37297	CH22_5551FG_736_1_- CH22_FGENES.736-1	1.2
	322340	EOS22271	AF088076 EST cluster (not in UniGene)	1.2
	307954	EOS07885	AI419692 EST singleton (not in UniGene) with exon hit	1.2
65	326615	EOS28546	c_7_hs gi 5868239 ref  gn 2 + 35214 35347 ex 3 4 CDSI 11.49 134 3651	1.2
		CH.07_hs gi 5868239		1.2
	317787	EOS17718	AW339612 Hs.249364 ESTs	1.2
	335288	EOS35219	CH22_2630FG_527_1_LINK_EM:AC005500.GENSCAN.421-1	1.2
		CH22_FGENES.527_1		1.2
70	323175	EOS23106	AI827137 Hs.184023 ESTs	1.2
	330893	EOS30824	AA149620 Hs.71999 ESTs	1.2
	306810	EOS06741	AI057294 EST singleton (not in UniGene) with exon hit	1.2
	338239	EOS38170	CH22_6833FG__LINK_EM:AC005500.GENSCAN.264-5	1.2
		CH22_EM:AC005500.GENSCAN.264-5		1.2
75	332347	EOS32278	W60326 Hs.221716 ESTs	1.2
	309782	EOS09713	AW275156 Hs.156110 Immunoglobulin kappa variable 1D-8	1.2
	322518	EOS22449	AI133446 EST cluster (not in UniGene)	1.2
	301187	EOS01118	AA806542 EST cluster (not in UniGene) with exon hit	1.2
	312129	EOS12060	AW300867 EST cluster (not in UniGene)	1.2
80	334714	EOS34645	CH22_2024FG_421_25_LINK_EM:AC005500.GENSCAN.282-25	1.2
		CH22_FGENES.421_25		1.2
	316586	EOS16517	AI205077 Hs.144689 ESTs	1.2
	320488	EOS20419	R31386 EST cluster (not in UniGene)	1.2
	327458	EOS27389	c_2_hs gi 6004455 ref  gn 3 + 173267 173378 ex 5 7 CDSI 4.03 122 1184	1.2
		CH.02_hs gi 6004455		1.2
85	336707	EOS36638	CH22_4212FG_64_3_- CH22_FGENES.64-3	1.2
	313561	EOS13492	AA040155 EST cluster (not in UniGene)	1.2

	330906	EOS30837	AA169498	Hs.72804	ESTs	1.2
	330987	EOS30918	H40988	Hs.131965	ESTs; Weakly similar to !!! ALU SUBFAMILY J WARNING ENTRY !!! [H.sapiens]	1.2
	325041	EOS24972	AI809182	Hs.130907	ESTs	1.2
	313225	EOS13156	AA502384	Hs.151529	ESTs	1.2
5	305295	EOS05226	AA687131		EST singleton (not in UniGene) with exon hit	1.2
	306896	EOS05827	AI093383		EST singleton (not in UniGene) with exon hit	1.2
	326981	EOS26912	c21_hs gil6588016[ref] gn 3 + 105091 106038 ex 1 1 CDSs 122.69 948 567			
				CH.21_hs gil6588016		1.2
10	332225	EOS32156	N33213	Hs.100425	ESTs	1.2
	318802	EOS18733	R19443	Hs.92414	ESTs	1.2
	318413	EOS18344	AI138592	Hs.144936	ESTs	1.2
	312292	EOS12223	AW451893	Hs.151124	ESTs	1.2
	323753	EOS23684	AA327102		EST cluster (not in UniGene)	1.2
15	313582	EOS13513	AW207684	Hs.13583	ESTs	1.2
	317836	EOS17767	AA983913	Hs.128929	ESTs	1.2
	332868	EOS32799	CH22_86FG_28_8_LINK_C20H12.GENSCAN.18-8			
				CH22_FGENES.28_8		1.2
	336924	EOS36855	CH22_4699FG_347_9		CH22_FGENES.347-9	1.2
	327791	EOS27722	c_5_hs gil5867977[ref] gn 1 + 22491 22610 ex 6 7 CDSs 11.29 120 658			1.2
20				CH.05_hs gil5867977		1.2
	330717	EOS30648	AA233926	Hs.23635	ESTs	1.2
	322944	EOS22875	AA112573		EST cluster (not in UniGene)	1.2
	312108	EOS12039	T82331	Hs.127453	ESTs	1.2
	332570	EOS32501	AA401376	Hs.26176	ESTs	1.2
25	330880	EOS30811	AA132420	Hs.53542	KIAA0986 protein	1.2
	310341	EOS10272	AW302773		EST cluster (not in UniGene)	1.2
	334012	EOS33943	CH22_1290FG_313_3_LINK_EM:AC005500.GENSCAN.169-3			
				CH22_FGENES.313_3		1.2
30	318230	EOS18161	AA558125		EST cluster (not in UniGene)	1.2
	336071	EOS36002	CH22_3457FG_685_3_LINK_DJ3210.GENSCAN.21-6			1.2
				CH22_FGENES.685_3		1.2
	338510	EOS38441	CH22_7208FG__LINK_EM:AC005500.GENSCAN.391-22			1.2
				CH22_EM:AC005500.GENSCAN.391-22		1.2
35	334487	EOS34418	CH22_1786FG_395_9_LINK_EM:AC005500.GENSCAN.258-10			1.2
				CH22_FGENES.395_9		1.2
	320661	EOS20592	AA864846		EST cluster (not in UniGene)	1.2
	335200	EOS35131	CH22_2538FG_508_9_LINK_EM:AC005500.GENSCAN.401-9			1.2
				CH22_FGENES.508_9		1.2
40	333582	EOS33513	CH22_842FG_201_2_LINK_EM:AC005500.GENSCAN.72-3			1.2
				CH22_FGENES.201_2		1.2
	320789	EOS20720	R78712		EST cluster (not in UniGene)	1.2
	321185	EOS21116	H51659	Hs.189854	ESTs	1.2
	337740	EOS37671	CH22_6085FG__LINK_EM:AC000097.GENSCAN.100-6			1.2
45	315064	EOS14995	AA775208	Hs.136423	ESTs	1.2
	334883	EOS34814	CH22_2197FG_451_6_LINK_EM:AC005500.GENSCAN.340-6			1.2
				CH22_FGENES.451_6		1.2
	331825	EOS31756	AA411144	Hs.104768	ESTs	1.2
	319141	EOS19072	F12377		EST cluster (not in UniGene)	1.1
50	333682	EOS33613	CH22_944FG_247_10_LINK_EM:AC005500.GENSCAN.102-10			1.1
				CH22_FGENES.247_10		1.1
	336140	EOS36071	CH22_3530FG_705_2_LINK_DA59H18.GENSCAN.10-2			1.1
				CH22_FGENES.705_2		1.1
55	320727	EOS20658	U96044		EST cluster (not in UniGene)	1.1
	323947	EOS23878	AA649842	Hs.186667	ESTs	1.1
	324746	EOS24677	AA603367	Hs.222294	ESTs	1.1
	306744	EOS06675	AI031882		EST singleton (not in UniGene) with exon hit	1.1
	326511	EOS26448	c19_hs gil5867439[ref] gn 1 + 44732 46356 ex 6 6 CDSs 148.22 1625 2512			1.1
				CH.19_hs gil5867439		1.1
60	333597	EOS33528	CH22_858FG_211_5_LINK_EM:AC005500.GENSCAN.79-5			1.1
				CH22_FGENES.211_5		1.1
	330135	EOS30066	c21_p2 gil4456470[emb] gn 2 - 121583 121885 ex 2 2 CDSs 18.67 303 102			1.1
				CH.21_p2 gil4456470		1.1
65	315118	EOS15049	AA564921	Hs.143899	ESTs	1.1
	302893	EOS02824	AL117539	Hs.173515	Homo sapiens mRNA; cDNA DKFZp586H021 (from clone DKFZp586H021)	1.1
	337169	EOS37100	CH22_5189FG_563_1		CH22_FGENES.563-1	1.1
	336121	EOS36052	CH22_3510FG_701_6_LINK_DA59H18.GENSCAN.8-6			1.1
				CH22_FGENES.701_6		1.1
70	323332	EOS23263	AI829520	Hs.227513	ESTs	1.1
	320911	EOS20842	AI056872	Hs.133386	ESTs	1.1
	327990	EOS27921	c_6_hs gil5868218[ref] gn 2 - 36225 36503 ex 1 2 CDSs 16.35 279 1419			1.1
				CH.06_hs gil5868218		1.1
	320425	EOS20356	C14069	Hs.201627	ESTs; Moderately similar to !!! ALU SUBFAMILY SQ WARNING ENTRY !!! [H.sapiens]	1.1
	327075	EOS27006	c21_hs gil6531965[ref] gn 58 + 4041318 4041431 ex 4 4 CDSs 1.79 114 1285			1.1
75				CH.21_hs gil6531965		1.1
	314384	EOS14315	AA535840	Hs.162203	ESTs; Weakly similar to alternatively spliced product using exon 13A [H.sapiens]	1.1
	338716	EOS38647	CH22_7502FG__LINK_EM:AC005500.GENSCAN.488-9			1.1
80	330886	EOS30817	AA135606	Hs.189384	ESTs; Weakly similar to !!! ALU SUBFAMILY J WARNING ENTRY !!! [H.sapiens]	1.1
	327331	EOS27262	c_1_hs gil5867516[ref] gn 4 - 55606 55737 ex 2 6 CDSs 7.01 132 2349			1.1
				CH.01_hs gil5867516		1.1
	326714	EOS26645	c20_hs gil5867595[ref] gn 2 + 124490 124568 ex 5 6 CDSs 0.11 79 1020			1.1
				CH.20_hs gil5867595		1.1
85	316734	EOS16665	AW080237	Hs.252884	ESTs	1.1
	311660	EOS11591	AI978583	Hs.232161	ESTs	1.1
	312757	EOS12688	AI285970	Hs.183817	ESTs	1.1

331686	EOS31617	W88502	Hs.182258	ESTs		1.1
337840	EOS37771	CH22_6223FG_LINK_EM:AC005500.GENSCAN.26-9		CH22_EM:AC005500.GENSCAN.26-9		1.1
5	332093	EOS32024	AA608794	Hs.112592	ESTs	1.1
	319595	EOS19526	H81361	Hs.194485	ESTs	1.1
	315990	EOS15921	AI800041	Hs.190555	ESTs	1.1
	322438	EOS22369	W44531	Hs.167851	ESTs	1.1
	332965	EOS32896	CH22_189FG_50_3_LINK_EM:AC000097.GENSCAN.3-5		CH22_FGENES.50_3	1.1
10	337182	EOS37113	CH22_5204FG_570_2		CH22_FGENES.570-2	1.1
	334948	EOS34879	CH22_2269FG_465_15_LINK_EM:AC005500.GENSCAN.359-13		CH22_FGENES.465_15	1.1
	325864	EOS25795	c16_hs_gi 5867069 refl	gn 2 - 110834 110904 ex 3 3 CDSf	9.76 71 457	1.1
				Ch.16_hs_gi 5867069		1.1
15	337760	EOS37691	CH22_6110FG_LINK_EM:AC000097.GENSCAN.116-8		CH22_EM:AC000097.GENSCAN.116-8	1.1
	315422	EOS15353	AW135357	Hs.192374	ESTs	1.1
	338899	EOS38820	CH22_7746FG_LINK_DJ32I10.GENSCAN.7-1		CH22_DJ32I10.GENSCAN.7-1	1.1
20	332961	EOS32892	CH22_185FG_48_18_LINK_EM:AC000097.GENSCAN.2-14		CH22_FGENES.48_18	1.1
	314703	EOS14634	AI791249		EST cluster (not in UniGene)	1.1
	317791	EOS17722	AI801500	Hs.128457	ESTs	1.1
25	333680	EOS33611	CH22_942FG_247_7_LINK_EM:AC005500.GENSCAN.102-7		CH22_FGENES.247_7	1.1
	322419	EOS22350	AA248987	Hs.14084	ESTs; Highly similar to zinc RING finger protein SAG [M.musculus]	1.1
	338124	EOS38055	CH22_6661FG_LINK_EM:AC005500.GENSCAN.196-2		CH22_EM:AC005500.GENSCAN.196-2	1.1
30	308884	EOS08815	AI833131	Hs.179100	ESTs	1.1
	333349	EOS33280	CH22_595FG_140_3_LINK_EM:AC005500.GENSCAN.20-3		CH22_FGENES.140_3	1.1
	313150	EOS13081	AA824410	Hs.165003	ESTs	1.1
	339208	EOS39139	CH22_8146FG_LINK_FF113D11.GENSCAN.6-3		CH22_FF113D11.GENSCAN.6-3	1.1
35	335653	EOS35584	CH22_3013FG_590_4_LINK_EM:AC005500.GENSCAN.484-4		CH22_FGENES.590_4	1.1
	319524	EOS19455	AA682865	Hs.194441	ESTs	1.1
	301576	EOS01507	AI682905	Hs.146875	ESTs; Weakly similar to !!! ALU SUBFAMILY J WARNING ENTRY !!! [H.sapiens]	1.1
40	317598	EOS17529	AW206035	Hs.192123	ESTs	1.1
	333473	EOS33404	CH22_724FG_162_3_LINK_EM:AC005500.GENSCAN.42-10		CH22_FGENES.162_3	1.1
	333949	EOS33880	CH22_1225FG_303_5_LINK_EM:AC005500.GENSCAN.162-9		CH22_FGENES.303_5	1.1
45	339256	EOS39187	CH22_8207FG_LINK_BA354I12.GENSCAN.7-11		CH22_BA354I12.GENSCAN.7-11	1.1
	332884	EOS32815	CH22_104FG_33_5_LINK_C20H12.GENSCAN.22-7		CH22_FGENES.33_5	1.1
	314660	EOS14591	AA436007	Hs.188780	ESTs	1.1
50	333220	EOS33151	CH22_457FG_104_12_LINK_EM:AC000097.GENSCAN.108-11		CH22_FGENES.104_12	1.1
	308106	EOS08037	AI476803		EST singleton (not in UniGene) with exon hit	1.1
	320709	EOS20640	AA456660	Hs.154165	ESTs	1.1
	307612	EOS07543	AI290787		EST singleton (not in UniGene) with exon hit	1.1
55	330286	EOS30217	c_5_p2_gi 6671913 gb A	gn 2 - 31050 31171 ex 2 7 CDS1	8.84 122 791	1.1
	304495	EOS04426	AA446448		CH.05_p2_gi 6671913	1.1
	310583	EOS10514	AW205632	Hs.211198	ESTs	1.1
	332896	EOS32827	CH22_117FG_35_10_LINK_C20H12.GENSCAN.24-9		CH22_FGENES.35_10	1.1
60	337602	EOS37533	CH22_5895FG_LINK_C20H12.GENSCAN.15-1		CH22_C20H12.GENSCAN.15-1	1.1
	307626	EOS07557	AI300035		EST singleton (not in UniGene) with exon hit	1.1
	334696	EOS34627	CH22_2006FG_421_5_LINK_EM:AC005500.GENSCAN.282-5		CH22_FGENES.421_5	1.1
65	318662	EOS18583	T53259		EST cluster (not in UniGene)	1.1
	337844	EOS37775	CH22_6229FG_LINK_EM:AC005500.GENSCAN.30-9		CH22_EM:AC005500.GENSCAN.30-9	1.1
	334823	EOS34754	CH22_2137FG_437_5_LINK_EM:AC005500.GENSCAN.301-7		CH22_FGENES.437_5	1.1
70	333928	EOS33859	CH22_1201FG_299_2_LINK_EM:AC005500.GENSCAN.158-5		CH22_FGENES.299_2	1.1
	337503	EOS37434	CH22_5738FG_803_1		CH22_FGENES.803-1	1.1
	323044	EOS22975	AA148725	Hs.154190	ESTs	1.1
75	329164	EOS29095	c_x_hs_gi 5868691 refl	gn 1 + 62305 62517 ex 2 2 CDS1	17.51 213 1868	1.1
	335468	EOS35399	CH22_2819FG_567_4_LINK_EM:AC005500.GENSCAN.454-12		CH22_FGENES.567_4	1.1
	338962	EOS38893	CH22_7838FG_LINK_DJ32I10.GENSCAN.23-39		CH22_DJ32I10.GENSCAN.23-39	1.1
80	323570	EOS23501	AL038623	Hs.208752	ESTs; Weakly similar to !!! ALU SUBFAMILY SX WARNING ENTRY !!! [H.sapiens]	1.1
	333568	EOS33499	CH22_826FG_185_1_LINK_EM:AC005500.GENSCAN.64-1		CH22_FGENES.185_1	1.1
	331865	EOS31796	AA425756	Hs.98445	ESTs	1.1
85	336246	EOS36177	CH22_3644FG_746_5_LINK_DA59H18.GENSCAN.48-4		CH22_FGENES.746_5	1.1
	337238	EOS37169	CH22_5343FG_641_3		CH22_FGENES.641-3	1.1

5	305089	EOS05020	AA642622	EST singleton (not in UniGene) with exon hit	1.1
	300097	EOS00028	AI916973	Hs.213603 ESTs	1.1
	313134	EOS13065	N63406	Hs.258697 ESTs	1.1
	337452	EOS37383	CH22_5665FG_775_1	CH22_FGENES.775-1	1.1
	325433	EOS25364	c12_hs gj 5866936 ref	gn 4 - 480706 480826 ex 3 4 CDSi 1.99 121 818 CH.12_hs gj 5866936	1.1
10	335999	EOS35930	CH22_3380FG_657_1_LINK_EM	DJ246D7.GENSCAN.11-1 CH22_FGENES.657_1	1.1
	333580	EOS33511	CH22_840FG_199_2_LINK_EM	AC005500.GENSCAN.71-2 CH22_FGENES.199_2	1.1
	336836	EOS36767	CH22_4512FG_247_11	CH22_FGENES.247-11	1.1
	334677	EOS34608	CH22_1986FG_418_30_LINK_EM	AC005500.GENSCAN.279-31 CH22_FGENES.418_30	1.1
15	329062	EOS28993	c_X_hs gj 5866590 ref	gn 3 - 58977 59094 ex 4 11 CDSi -6.19 118 627 CH.X_hs gj 5866590	1.1
	333671	EOS33602	CH22_932FG_245_5_LINK_EM	AC005500.GENSCAN.100-12 CH22_FGENES.245_5	1.1
	304941	EOS04872	AA612612	EST singleton (not in UniGene) with exon hit	1.1
20	315772	EOS15703	AW515373	Hs.158893 ESTs	1.1
	301281	EOS01212	AA843986	Hs.190586 ESTs	1.1
	333520	EOS33451	CH22_777FG_174_3_LINK_EM	AC005500.GENSCAN.53-6 CH22_FGENES.174_3	1.1
25	315203	EOS15134	AI559820	Hs.199438 ESTs	1.1
	315927	EOS15858	AW025517	Hs.133250 ESTs	1.1
	317161	EOS17092	AA972165	Hs.150308 ESTs	1.1
	337692	EOS37623	CH22_6028FG__LINK_EM	AC000097.GENSCAN.78-12 CH22_EM:AC000097.GENSCAN.78-12	1.1
30	331472	EOS31403	N24830	yx70a02.s1 Soares melanocyte 2NbHomo sapiens cDNA clone IMAGE:267050 3' similar to gb M87912 HUMALNE562 Human carcinoma cell-derived Alu RNA transcript, (rRNA);contains Alu repetitive element; mRNA sequence.	1.1
	336439	EOS36370	CH22_3859FG_827_4_LINK_EM	DJ579N16.GENSCAN.1-3 CH22_FGENES.827_4	1.1
	326882	EOS26813	c20_hs gj 6682509 ref	gn 2 - 167988 168179 ex 4 4 CDSf 18.69 192 2238 CH.20_hs gj 6682509	1.1
35	336977	EOS36908	CH22_4793FG_380_9	CH22_FGENES.380-9	1.1
	333983	EOS33914	CH22_1260FG_310_7_LINK_EM	AC005500.GENSCAN.167-5 CH22_FGENES.310_7	1.1
	328878	EOS28809	c_7_hs gj 6552423 ref	gn 1 + 105580 105774 ex 6 7 CDSi 2.91 195 6195 CH.07_hs gj 6552423	1.1
40	330415	EOS30346	D83777	Hs.75137 KIAA0193 gene product	1.1
	324824	EOS24755	AI826999	Hs.224624 ESTs	1.1
	325815	EOS25746	c14_hs gj 6682483 ref	gn 1 - 129273 130754 ex 1 1 CDSi 11.82 1482 2225 CH.14_hs gj 6682483	1.1
45	300463	EOS00394	N52510	Hs.186470 ESTs	1.1
	335708	EOS35639	CH22_3069FG_599_8_LINK_EM	AC005500.GENSCAN.490-11 CH22_FGENES.599_8	1.1
	324575	EOS24506	AW502257	EST cluster (not in UniGene)	1.1
	337951	EOS37882	CH22_6391FG__LINK_EM	AC005500.GENSCAN.94-1 CH22_EM:AC005500.GENSCAN.94-1	1.1
50	335935	EOS35866	CH22_3313FG_646_6_LINK_EM	DJ246D7.GENSCAN.1-5 CH22_FGENES.646_6	1.1
	334914	EOS34845	CH22_2233FG_457_3_LINK_EM	AC005500.GENSCAN.346-2 CH22_FGENES.457_3	1.1
55	309527	EOS09458	AW150648	Hs.75621 protease inhibitor 1 (anti-elastase); alpha-1-antitrypsin	1.1
	318901	EOS18832	AW368520	Hs.24639 ESTs	1.1
	320484	EOS20415	AA094436	Hs.155712 follistatin-like 1	1.1
	333665	EOS33596	CH22_926FG_244_1_LINK_EM	AC005500.GENSCAN.99-1 CH22_FGENES.244_1	1.1
60	335860	EOS35791	CH22_3235FG_629_5_LINK_EM	AC005500.GENSCAN.519-4 CH22_FGENES.629_5	1.1
	313339	EOS13270	AI682536	Hs.163495 ESTs	1.1
	300149	EOS00080	AW448916	Hs.149018 ESTs	1.1
	318112	EOS18043	AI028162	Hs.132307 ESTs	1.1
65	337807	EOS37738	CH22_6178FG__LINK_EM	AC005500.GENSCAN.9-4 CH22_EM:AC005500.GENSCAN.9-4	1.1
	336917	EOS36848	CH22_4688FG_346_4	CH22_FGENES.346-4	1.1
	337489	EOS37420	CH22_5722FG_799_2	CH22_FGENES.799-2	1.1
	320112	EOS20043	T92107	Hs.188489 ESTs	1.1
	332975	EOS32906	CH22_199FG_51_10_LINK_EM	AC000097.GENSCAN.4-12 CH22_FGENES.51_10	1.1
70	327805	EOS27736	c_5_hs gj 5867968 ref	gn 2 + 19952 20019 ex 1 2 CDSf 9.47 68 988 CH.05_hs gj 5867968	1.1
	339215	EOS39146	CH22_8153FG__LINK_EM	FF113D11.GENSCAN.6-10 CH22_FF113D11.GENSCAN.6-10	1.1
75	311965	EOS11896	T69279	EST cluster (not in UniGene)	1.1
	314043	EOS13974	AA827082	EST cluster (not in UniGene)	1.1
	333447	EOS33378	CH22_697FG_154_5_LINK_EM	AC005500.GENSCAN.35-6 CH22_FGENES.154_5	1.1
80	333242	EOS33173	CH22_481FG_111_6_LINK_EM	AC000097.GENSCAN.120-5 CH22_FGENES.111_6	1.1
	338596	EOS38527	CH22_7343FG__LINK_EM	AC005500.GENSCAN.437-2 CH22_EM:AC005500.GENSCAN.437-2	1.1
	329989	EOS29920	c16_p2 gj 4567166 gb A	gn 2 + 72861 73052 ex 1 3 CDSf 18.02 192 590 CH.16_p2 gj 4567166	1.1
85	315675	EOS15606	AA652272	Hs.197320 ESTs	1.1
	336722	EOS36653	CH22_4245FG_84_2	CH22_FGENES.84-2	1.1

	334220	EOS34151	CH22_1503FG_359_4_LINK_EM:AC005500.GENSCAN.217-7 CH22_FGENES.359_4	1.1
5	336703	EOS36634	CH22_4201FG_56_3 CH22_FGENES.56-3	1.1
	336397	EOS36328	CH22_3812FG_823_12_LINK_EM:AC005500.GENSCAN.6-11 CH22_FGENES.823_12	1.1
	316105	EOS16036	AW295687 Hs.254420 ESTs	1.1
	334661	EOS34592	CH22_1969FG_418_9_LINK_EM:AC005500.GENSCAN.279-13 CH22_FGENES.418_9	1.1
10	307783	EOS07714	AI347274 EST singleton (not in UniGene) with exon hit	1.1
	333997	EOS33928	CH22_1275FG_310_22_LINK_EM:AC005500.GENSCAN.167-21 CH22_FGENES.310_22	1.1
	331903	EOS31834	AA436673 Hs.29417 Homo sapiens mRNA; cDNA DKFZp586B0323 (from clone DKFZp586B0323)	1.1
	328249	EOS28180	c_6_hs gi 6381891 ref  gn 2 - 96352 96527 ex 2 3 CDSI 6.19 176 4550 CH.06_hs gi 6381891	1.1
15	338251	EOS38182	CH22_6849FG__LINK_EM:AC005500.GENSCAN.270-1 CH22_EM:AC005500.GENSCAN.270-1	1.1
	323561	EOS23492	AA825426 Hs.238832 ESTs; Weakly similar to !!! ALU SUBFAMILY J WARNING ENTRY !!! [H.sapiens]	1.1
	301464	EOS01395	AA991519 Hs.253324 ESTs	1.1
20	335916	EOS35847	CH22_3293FG_636_12_LINK_EM:AC005500.GENSCAN.526-12 CH22_FGENES.636_12	1.1
	321828	EOS21759	X56197 EST cluster (not in UniGene)	1.1
	327413	EOS27344	c_2_hs gi 5867750 ref  gn 3 + 101410 101508 ex 4 5 CDSI 4.34 99 587 CH.02_hs gi 5867750	1.1
25	334474	EOS34405	CH22_1773FG_394_5_LINK_EM:AC005500.GENSCAN.257-5 CH22_FGENES.394_5	1.1
	336739	EOS36670	CH22_4291FG_117_3 CH22_FGENES.117-3	1.1
	316517	EOS16448	AI784315 Hs.123163 ESTs	1.1
	325519	EOS25450	c12_hs gi 6017036 ref  gn 5 - 186804 186915 ex 1 3 CDSI 8.36 112 2508 CH.12_hs gi 6017036	1.1
30	333875	EOS33806	CH22_1145FG_291_11_LINK_EM:AC005500.GENSCAN.149-6 CH22_FGENES.291_11	1.1
	338221	EOS38152	CH22_6797FG__LINK_EM:AC005500.GENSCAN.246-10 CH22_EM:AC005500.GENSCAN.246-10	1.1
35	336878	EOS36809	CH22_4617FG_318_5 CH22_FGENES.318-5	1.1
	337919	EOS37850	CH22_6338FG__LINK_EM:AC005500.GENSCAN.66-5 CH22_EM:AC005500.GENSCAN.66-5	1.1
	309828	EOS09759	AW293999 EST singleton (not in UniGene) with exon hit	1.1
	305259	EOS05190	AA679225 EST singleton (not in UniGene) with exon hit	1.1
40	333922	EOS33853	CH22_1195FG_296_13_LINK_EM:AC005500.GENSCAN.155-16 CH22_FGENES.296_13	1.1
	322092	EOS22023	AF085833 EST cluster (not in UniGene)	1.1
	313356	EOS13287	AI266254 Hs.132929 ESTs	1.1
	318847	EOS18778	Z42908 Hs.12308 ESTs	1.1
45	337175	EOS37106	CH22_5195FG_567_1 CH22_FGENES.567-1	1.1
	336979	EOS36910	CH22_4802FG_385_4 CH22_FGENES.385-4	1.1
	312169	EOS12100	AI064824 Hs.193385 ESTs	1.1
	336198	EOS36129	CH22_3595FG_719_2_LINK_EM:DA59H18.GENSCAN.21-2 CH22_FGENES.719_2	1.1
50	321948	EOS21879	AA309612 Hs.118797 ubiquitin-conjugating enzyme E2 D 3 (homologous to yeast UBC4/5)	1.1
	324692	EOS24623	AA557952 EST cluster (not in UniGene)	1.1
	330395	EOS30326	D10923 Hs.137555 putative chemokine receptor; GTP-binding protein	1.1
	333119	EOS33050	CH22_347FG_80_4_LINK_EM:AC000097.GENSCAN.65-4 CH22_FGENES.80_4	1.1
55	316012	EOS15943	AA764950 Hs.119898 ESTs	1.1
	300142	EOS00073	AI743419 Hs.205707 ESTs	1.1
	317215	EOS17146	AW014242 Hs.159998 ESTs	1.1
	329526	EOS29457	c10_p2 gi 3983506 gb U gn 2 + 12251 12325 ex 3 3 CDSI 7.37 75 178 CH.10_p2 gi 3983506	1.1
60	317409	EOS17340	AA764968 Hs.4864 KIAA0892 protein	1.1
	339230	EOS39161	CH22_8171FG__LINK_EM:BA354112.GENSCAN.1-6 CH22_BA354112.GENSCAN.1-6	1.1
	311598	EOS11529	AW023595 Hs.232048 ESTs	1.1
	339164	EOS39095	CH22_8091FG__LINK_EM:DA59H18.GENSCAN.69-4 CH22_DA59H18.GENSCAN.69-4	1.1
65	326725	EOS26656	c20_hs gi 6552456 ref  gn 2 - 223005 223125 ex 5 6 CDSI 6.10 121 1038 CH.20_hs gi 6552456	1.1
	330952	EOS30883	H02855 Hs.29567 ESTs	1.1
	334621	EOS34552	CH22_1928FG_412_4_LINK_EM:AC005500.GENSCAN.275-4 CH22_FGENES.412_4	1.1
70	301685	EOS01616	W67730 EST cluster (not in UniGene) with exon hit	1.1
	308718	EOS08712	AI811707 EST singleton (not in UniGene) with exon hit	1.1
	323413	EOS23344	AA248828 Hs.225676 ESTs	1.1
	306723	EOS06654	AI026151 EST singleton (not in UniGene) with exon hit	1.1
	331258	EOS31189	Z41777 Hs.27413 ESTs	1.1
	313028	EOS12959	A1355433 Hs.190856 ESTs	1.1
	333002	EOS32933	CH22_226FG_59_3_LINK_EM:AC000097.GENSCAN.21-3 CH22_FGENES.59_3	1.1
75	303011	EOS02942	AF090405 EST cluster (not in UniGene) with exon hit	1.1
	317687	EOS17618	AA972990 Hs.127904 ESTs	1.1
	328779	EOS28710	c_7_hs gi 5868309 ref  gn 4 + 41570 41639 ex 1 5 CDSF 2.65 70 5365 CH.07_hs gi 5868309	1.1
	338707	EOS38638	CH22_7487FG__LINK_EM:AC005500.GENSCAN.482-2 CH22_EM:AC005500.GENSCAN.482-2	1.1
80	337974	EOS37905	CH22_6427FG__LINK_EM:AC005500.GENSCAN.106-3 CH22_EM:AC005500.GENSCAN.106-3	1.1
	332854	EOS32785	CH22_71FG_22_1_LINK_EM:C20H12.GENSCAN.15-2	1.1

				CH22_FGENES.22_1	1.1
5	311225	EOS11156	AW451982	Hs.248613 ESTs	1.1
	337094	EOS37025	CH22_5018FG_465_19_	CH22_FGENES.465-19	1.1
	319357	EOS19288	F13425	Hs.26229 ESTs	1.1
	332958	EOS32889	CH22_182FG_48_15_LINK_EM:AC000097.GENSCAN.2-11	CH22_FGENES.48_15	1.1
10	309634	EOS09565	AW193825	EST singleton (not in UniGene) with exon hit	1.1
	321171	EOS21102	AI769410	Hs.221461 ESTs	1.1
	316440	EOS16371	AI954795	Hs.156135 ESTs	1.1
	311665	EOS11596	AW294254	Hs.223742 ESTs	1.1
	327548	EOS27479	c_3_hs gi 5867797 ref  gn 2 - 81067 81130 ex 3 7 CDSI 6.42 64 12 CH.03_hs gi 5867797		1.1
15	314940	EOS14871	AW452768	Hs.162045 ESTs	1.1
	326401	EOS26332	c19_hs gi 5867355 ref  gn 1 + 35165 35332 ex 9 11 CDSI 0.41 168 788 CH.19_hs gi 5867355		1.1
	336347	EOS36278	CH22_3759FG_815_3_LINK_EM:AC005500.GENSCAN.1-24	CH22_FGENES.815_3	1.1
20	322297	EOS22228	W76548	Hs.136026 ESTs; Moderately similar to !!!! ALU SUBFAMILY SC WARNING ENTRY !!!! [H.sapiens]	1.1
	309977	EOS09908	AW451663	EST singleton (not in UniGene) with exon hit	1.1
	333466	EOS33397	CH22_717FG_161_2_LINK_EM:AC005500.GENSCAN.42-2	CH22_FGENES.161_2	1.1
	329170	EOS29101	c_x_hs gi 5868693 ref  gn 2 + 67924 68019 ex 6 8 CDSI 3.30 96 1882 CH.X_hs gi 5868693		1.1
25	329479	EOS29410	c10_p2 gi 3983526 gb A gn 3 - 7425 7561 ex 13 CDSI 4.33 137 22 CH.10_p2 gi 3983526		1.1
	326668	EOS26599	c20_hs gi 6552455 ref  gn 1 + 146726 146838 ex 11 11 CDSI 1.84 113 767 CH.20_hs gi 6552455		1.1
	319364	EOS19295	H06538	Hs.12270 ESTs	1.1
30	302988	EOS02919	W23986	Hs.34578 alpha2;3-sialyltransferase	1.1
	327687	EOS27618	c_4_hs gi 5867847 ref  gn 1 - 169293 169362 ex 2 3 CDSI -0.28 70 782 CH.04_hs gi 5867847		1.1
	339413	EOS39344	CH22_8405FG__LINK_DJ579N16.GENSCAN.5-8	CH22_DJ579N16.GENSCAN.5-8	1.1
35	306156	EOS06087	AA918274	Hs.76067 heat shock 27kD protein 1	1.1
	320858	EOS20789	D59968	EST cluster (not in UniGene)	1.1
	325447	EOS25378	c12_hs gi 5866941 ref  gn 3 - 372480 372621 ex 2 3 CDSI 9.16 142 1026 CH.12_hs gi 5866941		1.1
40	322696	EOS22627	AI064724	Hs.228468 ESTs	1.1
	329959	EOS29890	c16_p2 gi 5103803 gb A gn 3 + 188050 188193 ex 8 8 CDSI 2.01 144 361 CH.16_p2 gi 5103803		1.1
	312628	EOS12559	AA632817	Hs.190316 ESTs	1.1
	339305	EOS39236	CH22_8262FG__LINK_EM:AC005500.GENSCAN.21-3	CH22_BA35412.GENSCAN.21-3	1.1
45	311829	EOS11760	AI078483	Hs.134549 ESTs	1.1
	303270	EOS03201	AL120518	Hs.105352 ESTs	1.1
	321226	EOS21143	AA311443	Homo sapiens mRNA; cDNA DKFZp586E2317 (from clone DKFZp586E2317)	1.1
	335827	EOS35758	CH22_3200FG_620_1_LINK_EM:AC005500.GENSCAN.512-1	CH22_FGENES.620_1	1.1
50	336677	EOS36608	CH22_4155FG_43_5	CH22_FGENES.43-5	1.1
	330081	EOS30012	c19_p2 gi 6015314 gb A gn 1 - 5768 5835 ex 4 9 CDSI 2.88 68 162 CH.19_p2 gi 6015314		1.1
	339313	EOS39244	CH22_8272FG__LINK_EM:AC005500.GENSCAN.22-11	CH22_BA35412.GENSCAN.22-11	1.1
55	319936	EOS19867	W22152	EST cluster (not in UniGene)	1.1
	332858	EOS32789	CH22_76FG_24_1_LINK_EM:AC005500.GENSCAN.16-6	CH22_FGENES.24_1	1.1
	315630	EOS15561	AA648355	Hs.185155 ESTs; Weakly similar to echinoderm microtubule-associated protein-like EMAP2 [H.sapiens]	1.1
	332995	EOS32926	CH22_219FG_58_2_LINK_EM:AC000097.GENSCAN.19-2	CH22_FGENES.58_2	1.1
60	333441	EOS33372	CH22_691FG_151_5_LINK_EM:AC005500.GENSCAN.32-5	CH22_FGENES.151_5	1.1
	333496	EOS33427	CH22_748FG_168_6_LINK_EM:AC005500.GENSCAN.47-5	CH22_FGENES.168_6	1.1
65	339188	EOS39119	CH22_8123FG__LINK_EM:AC005500.GENSCAN.72-16	CH22_DA59H18.GENSCAN.72-16	1.1
	336981	EOS36912	CH22_4818FG_397_7	CH22_FGENES.397-7	1.1
	312142	EOS12073	AW298359	Hs.221069 ESTs	1.1
	315779	EOS15710	AW015736	Hs.211378 ESTs	1.1
	318596	EOS18527	AI470235	Hs.172698 EST	1.1
70	335701	EOS35632	CH22_3062FG_599_1_LINK_EM:AC005500.GENSCAN.490-2	CH22_FGENES.599_1	1.1
	319395	EOS19326	AW062570	Hs.13809 ESTs	1.1
	304236	EOS04167	W93278	EST singleton (not in UniGene) with exon hit	1.1
	307264	EOS07195	AI202211	EST singleton (not in UniGene) with exon hit	1.1
	334066	EOS33997	CH22_1344FG_327_21_LINK_EM:AC005500.GENSCAN.181-23	CH22_FGENES.327_21	1.1
75	327042	EOS26973	c21_hs gi 6531965 ref  gn 18 - 1380806 1381443 ex 1 5 CDSI 30.85 638 943 CH.21_hs gi 6531965		1.1
	326025	EOS25956	c17_hs gi 5867176 ref  gn 1 + 70854 70915 ex 6 8 CDSI -1.46 62 127 CH.17_hs gi 5867176		1.1
	325609	EOS25540	c14_hs gi 5866996 ref  gn 28 - 981751 981849 ex 1 10 CDSI 1.46 99 101 CH.14_hs gi 5866996		1.1
80	319983	EOS19914	T81429	EST cluster (not in UniGene)	1.1
	334298	EOS34229	CH22_1589FG_372_4_LINK_EM:AC005500.GENSCAN.232-5	CH22_FGENES.372_4	1.1
85	323203	EOS23134	AA203135	Hs.130186 ESTs	1.1

305700	EOS05631	AA815428	EST singleton (not in UniGene) with exon hit	1.1	
313304	EOS13235	AI334078	Hs.152438 ESTs	1.1	
310716	EOS10647	AI589618	Hs.192413 ESTs	1.1	
5	327049	EOS26980	c21_hs gi 6531965 ref  gn 24 - 1924026 1924110 ex 2 6 CDSi 9.43 85 1012 CH.21_hs gi 6531965 ESTs	1.1	
313749	EOS13680	AW450376	Hs.130803 ESTs	1.1	
307041	EOS06972	AI144243	EST singleton (not in UniGene) with exon hit	1.1	
322394	EOS22325	AF077208	EST cluster (not in UniGene)	1.1	
10	326416	EOS26347	c19_hs gi 5867362 ref  gn 3 - 45283 45375 ex 3 3 CDSi 5.65 93 923 CH.19_hs gi 5867362 ESTs	1.1	
333947	EOS33878	CH22_1221FG_303_1_LINK_EM:AC005500.GENSCAN.162-5 CH22_FGENES.303_1		1.1	
324609	EOS24540	AW299534	EST cluster (not in UniGene)	1.1	
15	330057	EOS29988	c17_p2 gi 6478962 gb A gn 3 + 75145 75287 ex 3 3 CDSi -2.56 143 150 CH.17_p2 gi 6478962 ESTs	1.1	
337603	EOS37534	CH22_5896FG__LINK_C20H12.GENSCAN.16-2 CH22_C20H12.GENSCAN.16-2		1.1	
332913	EOS32844	CH22_134FG_36_18_LINK_C20H12.GENSCAN.28-17 CH22_FGENES.36_18		1.1	
20	310026	EOS09957	T24895 Hs.100691 ESTs	1.1	
330153	EOS30084	c21_p2 gi 4325335 gb A gn 2 + 146951 147475 ex 2 2 CDSi 25.45 525 233 CH.21_p2 gi 4325335 ESTs	1.1		
334118	EOS34049	CH22_1396FG_330_19_LINK_EM:AC005500.GENSCAN.185-20 CH22_FGENES.330_19		1.1	
25	324795	EOS24726	AI494481 Hs.141579 ESTs	1.1	
332530	EOS32461	M31682 Hs.1735 inhibin; beta B (activin AB beta polypeptide)	1.1		
332048	EOS31979	AA496019 Hs.201591 ESTs	1.1		
334532	EOS34463	CH22_1834FG_402_13_LINK_EM:AC005500.GENSCAN.266-13 CH22_FGENES.402_13		1.1	
30	329762	EOS29693	c14_p2 gi 6048280 emb  gn 3 + 127744 127878 ex 2 4 CDSi 11.66 135 1054 CH.14_p2 gi 6048280 ESTs	1.1	
332909	EOS32840	CH22_130FG_36_13_LINK_C20H12.GENSCAN.28-10 CH22_FGENES.36_13		1.1	
35	321253	EOS21184	AI699484 EST cluster (not in UniGene)	1.1	
336572	EOS36503	CH22_4007FG_843_12_LINK_DJ579N16.GENSCAN.15-13 CH22_FGENES.843_12		1.1	
328768	EOS28699	c_7_hs gi 6017031 ref  gn 5 - 223741 224238 ex 1 1 CDSi 30.00 498 5285 CH.07_hs gi 6017031 ESTs		1.1	
40	334335	EOS34266	CH22_1627FG_375_12_LINK_EM:AC005500.GENSCAN.235-12 CH22_FGENES.375_12		1.1
334063	EOS33994	CH22_1341FG_327_17_LINK_EM:AC005500.GENSCAN.181-20 CH22_FGENES.327_17		1.1	
45	333011	EOS32942	CH22_235FG_61_3_LINK_EM:AC000097.GENSCAN.23-3 CH22_FGENES.61_3		1.1
304677	EOS04608	AA548071 EST singleton (not in UniGene) with exon hit	1.1		
313948	EOS13879	AW452823 Hs.135268 ESTs	1.1		
334358	EOS34289	CH22_1652FG_378_1_LINK_EM:AC005500.GENSCAN.239-1 CH22_FGENES.378_1		1.1	
50	328479	EOS28410	c_7_hs gi 5868449 ref  gn 1 - 331 560 ex 1 31 CDSi 18.51 230 2100 CH.07_hs gi 5868449 ESTs		1.1
335813	EOS35744	CH22_3185FG_618_1_LINK_EM:AC005500.GENSCAN.510-1 CH22_FGENES.618_1		1.1	
55	312430	EOS12361	AW139117 Hs.117494 ESTs	1.1	
324783	EOS24714	AA640770 EST cluster (not in UniGene)	1.1		
337776	EOS37707	CH22_6132FG__LINK_EM:AC000097.GENSCAN.119-18 CH22_EM:AC000097.GENSCAN.119-18		1.1	
327205	EOS27136	c_1_hs gi 5867447 ref  gn 5 + 167335 167576 ex 9 9 CDSi 15.50 242 259 CH.01_hs gi 5867447 ESTs		1.1	
60	315198	EOS15129	AI741506 Hs.186753 ESTs; Weakly similar to !!! ALU SUBFAMILY J WARNING ENTRY !!! [H.sapiens]	1.1	
336135	EOS36066	CH22_3525FG_704_3_LINK_EM:DA59H18.GENSCAN.9-5 CH22_FGENES.704_3		1.1	
318558	EOS18489	AW402677 Hs.90372 ESTs	1.1		
328152	EOS28083	c_6_hs gi 5868060 ref  gn 1 - 73981 74203 ex 1 8 CDSi 31.69 223 3411 CH.06_hs gi 5868060 ESTs		1.1	
65	330211	EOS30142	c_5_p2 gi 6013592 gb A gn 1 + 59158 59215 ex 2 4 CDSi 4.20 58 184 CH.05_p2 gi 6013592 ESTs		1.1
339280	EOS39211	CH22_8234FG__LINK_BA354I12.GENSCAN.14-12 CH22_BA354I12.GENSCAN.14-12		1.1	
70	332045	EOS31976	AA491253 Hs.155045 bromodomain adjacent to zinc finger domain; 2A	1.1	
313597	EOS13528	AW162263 Hs.249990 ESTs	1.1		
329503	EOS29434	c10_p2 gi 3983517 gb U gn 2 - 1801 1937 ex 1 4 CDSi 4.33 137 101 CH.10_p2 gi 3983517 ESTs		1.1	
75	333488	EOS33419	CH22_740FG_167_3_LINK_EM:AC005500.GENSCAN.46-10 CH22_FGENES.167_3		1.1
311960	EOS11891	AW440133 Hs.189690 ESTs	1.1		
320590	EOS20521	U67058 Hs.168102 Human proteinase activated receptor-2 mRNA; 3'UTR	1.1		
334047	EOS33978	CH22_1325FG_326_5_LINK_EM:AC005500.GENSCAN.175-5 CH22_FGENES.326_5		1.1	
80	304782	EOS04713	AA582081 EST singleton (not in UniGene) with exon hit	1.1	
324231	EOS24162	W60827 EST cluster (not in UniGene)	1.1		
327212	EOS27143	c_1_hs gi 5867463 ref  gn 1 - 42308 42424 ex 5 13 CDSi 6.58 117 325 CH.01_hs gi 5867463 ESTs		1.1	
85	336857	EOS35788	CH22_3232FG_629_1_LINK_EM:AC005500.GENSCAN.519-1 CH22_FGENES.629_1		1.1
317775	EOS17706	AA974603 Hs.181123 ESTs	1.1		
331053	EOS30984	N70242 Hs.183146 ESTs	1.1		

	335940	EOS35871	CH22_3318FG_646_13_LINK_DJ246D7.GENSCAN.1-12				
			CH22_FGENES.646_13				1.1
5	322568	EOS22499	W87342	Hs.209652	ESTs		1.1
	314091	EOS14022	AI253112	Hs.133540	ESTs		1.1
	313570	EOS13501	AA041455	Hs.209312	ESTs		1.1
	300967	EOS00898	AA565209	Hs.190216	ESTs		1.1
	314544	EOS14475	AA399018	Hs.250835	ESTs		1.1
	328321	EOS28252	c_7_hs_gi 5868373 ref  gn 7 - 1029614 1029673 ex 1 3 CDSI -2.40 60 448				
10			CH.07_hs_gi 5868373				1.1
	310979	EOS10910	AW445166	Hs.170802	ESTs		1.1
	310730	EOS10661	AI939421	Hs.160900	ESTs		1.1
	318471	EOS18402	AW137725	Hs.146874	ESTs		1.1
	315533	EOS15464	AW206191	Hs.152774	ESTs		1.1
	325751	EOS25682	c14_hs_gi 6682474 ref  gn 4 + 130437 130520 ex 6 7 CDSI 0.22 84 1666				
			CH.14_hs_gi 6682474				1.1
15	318780	EOS18711	R90906	Hs.113307	ESTs		1.1
	313271	EOS13202	AW444819	Hs.144851	ESTs; Weakly similar to C09F5.2 [C.elegans]		1.1
	304546	EOS04477	AA486074		EST singleton (not in UniGene) with exon hit		1.1
20	330618	EOS30549	X55990	Hs.73839	ribonuclease; RNase A family; 3 (eosinophil cationic protein)		1.1
	332931	EOS32862	CH22_152FG_38_5_LINK_C20H12.GENSCAN.29-5				
			CH22_FGENES.38_5				1.1
	336602	EOS36533	CH22_4047FG_372_4_LINK_EM:AC005500.GENSCAN.232-4				1.1
			CH22_FGENES.372_4				
25	311185	EOS11116	AI638294	Hs.224665	ESTs		1.1
	337585	EOS37516	CH22_5873FG__LINK_C20H12.GENSCAN.5-3				
			CH22_C20H12.GENSCAN.5-3				1.1
	310249	EOS10180	AW071751	Hs.13179	ESTs; Moderately similar to !!! ALU SUBFAMILY SQ WARNING ENTRY !!!! [H.sapiens]		1.1
	314578	EOS14509	AA410183	Hs.137475	ESTs		1.1
	310750	EOS10681	AI373163	Hs.170333	ESTs		1.1
30	333968	EOS33899	CH22_1245FG_307_4_LINK_EM:AC005500.GENSCAN.165-5				
			CH22_FGENES.307_4				1.1
	316133	EOS16064	AI187742	Hs.125562	ESTs		1.1
	308337	EOS08268	AI608947		EST singleton (not in UniGene) with exon hit		1.1
	326160	EOS26091	c17_hs_gi 5867254 ref  gn 6 - 112000 112137 ex 2 4 CDSI 8.01 138 1952				
			CH.17_hs_gi 5867254				1.1
35	336023	EOS35954	CH22_3406FG_669_12_LINK_DJ32110.GENSCAN.9-17				
			CH22_FGENES.669_12				1.1
	323479	EOS23410	AA278246		EST cluster (not in UniGene)		1.1
	336090	EOS36021	CH22_3477FG_689_2_LINK_DJ32110.GENSCAN.23-20				
			CH22_FGENES.689_2				1.1
40	311192	EOS11123	AW237220	Hs.211130	ESTs		1.1
	335081	EOS35012	CH22_2409FG_488_4_LINK_EM:AC005500.GENSCAN.384-6				
			CH22_FGENES.488_4				1.1
45	309519	EOS09450	AW148940	Hs.248647	EST		1.1
	321172	EOS21103	H49160	Hs.133472	ESTs		1.1
	301976	EOS01907	T97905		EST cluster (not in UniGene) with exon hit		1.1
	323012	EOS22943	AI832201	Hs.211469	ESTs		1.1
	319528	EOS19459	R08673	Hs.177514	ESTs		1.1
	329838	EOS29769	c14_p2_gi 6672062 emb  gn 2 + 33990 34098 ex 3 4 CDSI 9.11 109 2222				
			CH.14_p2_gi 6672062				1.1
50	302623	EOS02554	AB019571		EST cluster (not in UniGene) with exon hit		1.1
	334433	EOS34364	CH22_1731FG_385_8_LINK_EM:AC005500.GENSCAN.249-6				
			CH22_FGENES.385_8				1.1
55	304747	EOS04678	AA577816		EST singleton (not in UniGene) with exon hit		1.1
	333270	EOS33201	CH22_513FG_121_1_LINK_EM:AC005500.GENSCAN.4-11				
			CH22_FGENES.121_1				1.1
	307054	EOS06985	AI148181	Hs.176835	EST		1.1
	320764	EOS20695	R73070	Hs.246927	ESTs		1.1
60	321523	EOS21454	H78472	Hs.191325	ESTs; Weakly similar to cDNA EST yk414c9.3 comes from this gene [C.elegans]		1.1
	322114	EOS22045	AA643791	Hs.191740	ESTs		1.1
	303582	EOS03513	AA377444		EST cluster (not in UniGene) with exon hit		1.1
	322924	EOS22855	AA669253	Hs.193971	ESTs		1.1
	311179	EOS11110	AI880843	Hs.223333	ESTs		1.1
	318601	EOS18553	T39921		EST cluster (not in UniGene)		1.1
65	309791	EOS09722	AW276176	Hs.73742	ribosomal protein; large; P0		1.1
	333882	EOS33813	CH22_1153FG_292_4_LINK_EM:AC005500.GENSCAN.150-4				
			CH22_FGENES.292_4				1.1
	337645	EOS37576	CH22_5960FG__LINK_EM:AC000097.GENSCAN.10-8				
			CH22_EM:AC000097.GENSCAN.10-8				1.1
70	335623	EOS35554	CH22_2983FG_584_2_LINK_EM:AC005500.GENSCAN.478-2				
			CH22_FGENES.584_2				1.1
	314745	EOS14676	AA564489	Hs.137526	ESTs		1.1
	330790	EOS30721	T48536	Hs.105807	ESTs		1.1
	332071	EOS32002	AA598594	Hs.112475	ESTs		1.1
	312005	EOS11936	T78450	Hs.13941	ESTs		1.1
	330694	EOS30626	AA019806	Hs.108447	spinocerebellar ataxia 7 (olivopontocerebellar atrophy with retinal degeneration)		1.1
	330739	EOS30670	AA293477	Hs.227591	ESTs		1.1
	303042	EOS02973	AF129532		EST cluster (not in UniGene) with exon hit		1.1
	323091	EOS23022	AW014094	Hs.210761	ESTs		1.1
80	328820	EOS28751	c_7_hs_gi 5868330 ref  gn 1 + 90446 90602 ex 3 4 CDSI 10.20 157 5634				
			CH.07_hs_gi 5868330				1.1
	300472	EOS00403	T90622	Hs.82609	hydroxymethylbilane synthase		1.1
	310645	EOS10576	AI420742	Hs.163502	ESTs		1.1
	332238	EOS32169	N53480	Hs.108622	ESTs		1.1
	300966	EOS00897	AA564740	Hs.258401	ESTs		1.1
85	330437	EOS30368	HG2730-HT2827		Fibrinogen, A Alpha Polypeptide, Alt. Splice 2, E		1.1

	302292	EOS02223	AF067797	EST cluster (not in UniGene) with exon hit	1.1
	330138	EOS30069	c21_p2 gi 4210430 emb  gn 1 - 22334 22460 ex 3 3 CDSf 16.56 127 105 CH.21_p2 gi 4210430		1.1
5	332952	EOS32883	CH22_176FG_48_8_LINK_EM:AC000097.GENSCAN.2-4 CH22_FGENES.48_8		1.1
	319901	EOS19832	T77136 Hs.8765 RNA helicase-related protein		1.1
	321166	EOS21097	AA411263 Hs.128783 ESTs		1.1
	336227	EOS36158	CH22_3625FG_730_2_LINK_DA59H18.GENSCAN.36-2 CH22_FGENES.730_2		1.1
	302332	EOS02263	AI833168 Hs.184507 Homo sapiens Chromosome 16 BAC clone CIT987SK-A-328A3		1.1
10	313800	EOS13731	AW296132 Hs.166674 ESTs		1.1
	339356	EOS39287	CH22_8326FG_LINK_BA35412.GENSCAN.31-1 CH22_BA35412.GENSCAN.31-1		1.1
	324512	EOS24443	AW502125 EST cluster (not in UniGene)		1.1
	319235	EOS19166	F11330 Hs.177633 ESTs		1.1
15	320252	EOS20283	Y13323 Hs.145296 disintegrin protease		1.1
	338316	EOS38247	CH22_6944FG_LINK_EM:AC005500.GENSCAN.304-2 CH22_EM:AC005500.GENSCAN.304-2		1.1
	333964	EOS33895	CH22_1241FG_305_2_LINK_EM:AC005500.GENSCAN.164-2 CH22_FGENES.305_2		1.1
	312758	EOS12689	AA721107 Hs.202604 ESTs		1.1
20	338178	EOS38109	CH22_6726FG_LINK_EM:AC005500.GENSCAN.219-6 CH22_EM:AC005500.GENSCAN.219-6		1.1
	315199	EOS15130	AA877996 Hs.125376 ESTs		1.1
	312321	EOS12252	R66210 Hs.186937 ESTs		1.1
	338765	EOS38696	CH22_7588FG_LINK_EM:AC005500.GENSCAN.518-1 CH22_EM:AC005500.GENSCAN.518-1		1.1
25	330547	EOS30478	U32989 Hs.183671 tryptophan 2,3-dioxygenase		1.1
	315368	EOS15299	AW291563 Hs.152495 ESTs		1.1
	328691	EOS28622	c_7_hs gi 6588001 ref  gn 7 - 579598 579664 ex 2 3 CDSi 12.78 67 4326 CH.07_hs gi 6588001		1.1
	329179	EOS29110	c_x_hs gi 5868704 ref  gn 2 + 181639 181815 ex 3 4 CDSi 0.32 177 1939 CH.X_hs gi 5868704		1.1
30	327072	EOS27003	c21_hs gi 6531965 ref  gn 55 - 3796429 3797197 ex 4 4 CDSi 9.33 769 1270 CH.21_hs gi 6531965		1.1
	312056	EOS11987	T83748 Hs.189712 ESTs		1.1
	339128	EOS39059	CH22_8046FG_LINK_DA59H18.GENSCAN.55-2 CH22_DA59H18.GENSCAN.55-2		1.1
	307646	EOS07577	AI302236 EST singleton (not in UniGene) with exon hit		1.1
35	319198	EOS19129	F07354 EST cluster (not in UniGene)		1.1
	338556	EOS38487	CH22_7283FG_LINK_EM:AC005500.GENSCAN.417-8 CH22_EM:AC005500.GENSCAN.417-8		1.1
	306143	EOS06074	AA916314 EST singleton (not in UniGene) with exon hit		1.1
	332384	EOS32315	M11433 Hs.101850 retinol-binding protein 1; cellular		1.1
40	325100	EOS25031	T10265 Hs.116122 ESTs; Weakly similar to coded for by C. elegans cDNA yk30b3.5 [C.elegans]		1.1
	309839	EOS09770	AW296076 EST singleton (not in UniGene) with exon hit		1.1
	312180	EOS12111	A1248285 Hs.118348 ESTs		1.1
	330385	EOS30316	AA449749 Hs.31386 ESTs; Highly similar to secreted apoptosis related protein 1 [H.sapiens]		1.1
45	315882	EOS15813	A1831297 Hs.123310 ESTs		1.1
	325843	EOS25774	c16_hs gi 6552453 ref  gn 1 - 7126 7232 ex 1 3 CDSi 1.87 107 182 CH.16_hs gi 6552453		1.1
	330783	EOS30714	D60050 Hs.34812 ESTs		1.1
	317224	EOS17155	D56760 Hs.8122 ESTs		1.1
50	316042	EOS15973	AW297979 Hs.170698 ESTs		1.1
	333524	EOS33455	CH22_781FG_175_10_LINK_EM:AC005500.GENSCAN.53-15 CH22_FGENES.175_10		1.1
	302357	EOS02288	X03178 Hs.198246 group-specific component (vitamin D binding protein)		1.1
	309830	EOS09761	AW294725 EST singleton (not in UniGene) with exon hit		1.1
55	321489	EOS21420	AW392474 Hs.172759 ESTs; Moderately similar to !!! ALU SUBFAMILY SQ WARNING ENTRY !!!! [H.sapiens]		1.1
	312304	EOS12235	A4491949 Hs.183359 ESTs		1.1
	322026	EOS21957	AA233527 Hs.213289 low density lipoprotein receptor (familial hypercholesterolemia)		1.1
	302357	EOS02288	X03178 Hs.198246 group-specific component (vitamin D binding protein)		1.1

**TABLE 1A**

Table 1A shows the accession numbers for those primekeys in Table 1 which lack a unigeneID. Listed for each probeset is the gene cluster (CAT) number from which the oligonucleotides were designed. Gene clusters were compiled using sequences derived from Genbank ESTs and mRNAs. These sequences were clustered based on sequence similarity using Clustering and Alignment Tools (DoubleTwist, Oakland California). The Genbank accession numbers for sequences comprising each cluster are listed in the "Accession" column.

Pkey:	Unique Eos probeset identifier number
CAT number:	Gene cluster number
Accession:	Genbank accession numbers

	<b>Pkey</b>	<b>CAT number</b>	<b>Accession</b>
20	300611	337193_1	N75450 AA877636 AW137945 W05248 AA514763 AW972399 AI758397 AW195051
	301187	434061_1	AV976692 AA806542 AA745856
	301254	463589_1	AI049624 AA814705 AW404856 BE078289 BE078292
	301266	468223_1	AA829774 AI082020
	301454	534162_2	AI751738 AA977930
25	301615	5613_2	W39477 AK002047 NM_015515 T58707 AA386214 C19007 AA295466 T49621 T47323
	301661	7974_1	AK001735 AF227906 AI815558 AW238991 AL133051 AW272417 AI083492 AI816503 AW888717 AA333166 AI925832 BE048352 BE048415 AI141922 AW805674 AW805578 AA633581 AA424632 R71439 AW020988 AW976735 AA883247 W37208 AI091039 AW317020 BE221788 AA502917 AW009024 AI141417 BE349081 AI421443 AI080490 AI003921 AI373690 AI379240 AA424587 AA740607 AA972391 AA620797 AW271656 AA400517 AI370902 AI680616 AA757270 AA909500
30	301685	326972_1	N32107 R43738 AI270464 AI870568 AI085139 AA225666 Z41046 AI767739 AI270546 N56779
	301804	61_1	W67730 Z44630 AA490699 W67596 W76661 R21207 AK001468 AA190315 AA374980 AW961179 AA307782 AA315295 AA347194 AW953073 AW368190 AW368192 AA280772 AA251247 N85676 AI215522 AI216389 N87835 R12261 R57094 AI660045 AA347193 R16712 AW119006 N55905 N87768 AW900167 AI341261 AI818674 D20285 AI475165 AA300756 R40626 AI122827 AA133250 AI952488 AA970372 AA889845 AV069517 AI524385 AA190314 AI673359 AA971105 AI351088 AI872789 AI919056 AI611216 AK001472 BE568761 AA581004
35	301976	128835_1	T97905 AA101672
	302245	9482_1	H18835 R47363 AI460004 N31660 AA454774 AA551759 AI417040 AA694490 AA633315 AI344661 AA708532 AA878567 AI802702 AI913465 AW001160 AA932133 AI092908 AA026974 AW628573 AA592910 H18836 AI274428 C00675 AK00048 BE313619
40	302292	27735_1	AF067797 AB013456 NM_001169 AI791955 AW843925 AI732659 AA577625 AW083143 AW138645
	302476	31932_3	AF182294 NM_016200 AL046942 AI354410 AI697029 AI859557 AW188855 AW105437 AI358735 AW000959 AI491813 AW023693
45	302623	9705_1	AW836724 BE243668 AB019571 H43803
	302626	10441_1	AK001553 AK001951 AB021870 NM_016282 F01168 AA211870 AA078889 AA312979 AL138385 R70844 AA165658 AA007279 AA194688 H65871 AA476639 F01095 AA300170 R39487 AA649126 AA193643 AA418300 BE173477 N84408 AV024465 AA406255 BE173412 BE173583 BE173470 AW069288 AA372937 BE504414 AA209472 AI262833 AI628359 AI458075 AI476266 AA397706 AI768605 AW243125 AI056436 AA838326 AA810651 AI472025 N35912 AA165622 AI985532 AI139528 AA626087 W16998 AI632833 AW130827 AW662551 AA731459 AW780188 AI653447 AI694970 AA810662 AI199987 AI587402 AI492972 H65872 AI805624 AW194835 AW994874 R70790 AA836506 N53285 F00181 R83595 AI290941 AW936750 AW936703 AW936623 AW936785 AW936691 AW936668 AW936713 AW936788 AW936744 AW936613 AW936614 AW936665 AW936702 AW936647 AW936643 AW936712 AW936791 AW936624 AW936672 AW936754 AW936696 AW936802 AW936792 AW936689 AW936692 AW936645 AW936746 AW936801 AW936748 AW936661 AW936612 AW936697 AW936704 AW936695 AW936626 AW936794 AW936629 AW936577 AW936798 T35617
50	55	AA375943 R29459 AW936717 AA342108 AW963351 Z24876 AW936708 AW374110 AW936586 W20080 AW936752 W31803 AA093709 AA431256 AW803610 AA424959 W76607 AA432267 W72009 R70817 AW778851 AA890563 AA194632 AI089644 AI373864 AA890333 AI745574 AI095714 AI567507 AI280712 AW864083 AW468991 N48087 AA860500 AA279471 AA993680 AA676504 AI360949 AI052134 AI038657 AI439836 AA629147 AA651840 AA435925 AA854457 AW796472 AA838725 AA193407 AA302403 AW958003 AA342107 AA639258 AI435811 AA410342 N25790 AA156454 AI539628 AI275854 N58849 AI858171 AW338576 W15321 AA418342 AA780577 W04701 AA630452 AW769154 AI274286 N23736 BE465020 AI554346 AI920804 AA969728 AW193440 AI368697 AA115096 AA564981 AA630461 N91475 BE464381 AA913741 AA757161 Z24907 C00067 AA649290 AI245223 AA363098 AI520754 AA887983 AI273015 AW878871 AW878981 AA480455 AA709267 AW959521 AW959523 N90014 N32441 F00193 AA115095 AA147583 W19813 AI333349 AI197937 R39488 AW750110	
60	65	302655 41899_1 AJ227892 AA338715 BE074475 BE074469 BE074474 AW006182 AW572953 AI831725 AI762923 AI341466 AW449335 BE551686 AI692895 AI040410 AI276881 AI891008	
	302758 24028_3 AK001841 H40087 H11121 AW408676 N99603 AA984563 H92041 H11226		
	302882 458_60 AW403330 AF062097		

	302977	47403_3	AW263124 AI925166 AW105732 AA804479 BE621436 AF086399 W79085 W74440 AW992181 AA389686 AA314311 AA173955 AA677564 D59895 D60771 AI887733 C14814 AW162193 D81894 AA732538 AW150919 AA748064 AA769465 AA708143 BE327613 AA092726 AI692476 T35673 Z33600 AA134036 AI671394 AI267461 AW362795 AI769759 AA909042 AA130042 AW156938 AI753129 AI246205 AI823883 AI752836 D60770 AI336386 AI584003 AW627976 AI348676 D59894 AI969795 AW073259 AI400534 AI081318 AI082427 BE550515
5	302980	47495_1	AI925740 AF086489 W93435 W93345 AA337166 AW966214 AA336257 T11355 AW842435
	303011	41689_1	AF090405 AF090407 AF090406
	303037	35681_1	AF118395 NM_014317 AW376657 AW848189 AI261617 AI963829 AW848591 AW848598 AW376696 AW848523 AW848450 AW848655 AW848183 AW848550 AW376675 AI632752 AI590245 AI431824 AI857990 AI953341 AA888092 AW364968
10			AI188545 AI217741 AW275906 AI311481 AI991404 AI364963 AA628392 AA927982 AW150563 AA503063 AW079470 AW512180 AA889371 AW390132 AW609052 AW390112 AW390112 R12337 R14020 H98970 AI474918 N56139 AL135669 AW067702 AW372065 AW631389 AA083416 AA287511 AA602923 AA488914 AI167215 AW946829 R82855 AI948792 AA371333 AW953883 AW956152 C02539 AA298280 AI932587 AA022742
15			A1983021 AA195252 N58991 R78733 AW083996 H39614 AI365249 AW615389 AI927744 AI089971 N52205 AA083417 BE326666 BE349514 AI743785 AI640148 AI378211 AW181881 AI949484 W31374 AW628233 AA418406 AW068010 AI708085 AI092696 AI089823 AI277828 AA022660 AI440527 AW054937 AW474104 AI017436 AI159819 AI356716 AW473140 AW316518 N34522 AI675092 AI866697 AA864593 AW511185 AA488844 AA04975 N49111 Z39951 R37265
20	303042	5058_1	AIW505345 AF129532 AF126028 AA852108 BE169359 R83701 Z43904 BE613543 AA283163 AA905463 AW067849 R13544 R12337 R14020 H98970 AI474918 N56139 AL135669 AW067702 AW372065 AW631389 AA083416 AA287511 AA602923 AA488914 AI167215 AW946829 R82855 AI948792 AA371333 AW953883 AW956152 C02539 AA298280 AI932587 AA022742 A1983021 AA195252 N58991 R78733 AW083996 H39614 AI365249 AW615389 AI927744 AI089971 N52205 AA083417 BE326666 BE349514 AI743785 AI640148 AI378211 AW181881 AI949484 W31374 AW628233 AA418406 AW068010 AI708085 AI092696 AI089823 AI277828 AA022660 AI440527 AW054937 AW474104 AI017436 AI159819 AI356716 AW473140 AW316518 N34522 AI675092 AI866697 AA864593 AW511185 AA488844 AA04975 N49111 Z39951 R37265 AI141362 T25856 R20664 F03163 AI767927 AA805942 D79905 AI914645 AW190553 AI934213 AI458796 AA195385 R82854 W31965
25	303072	4654_1	AI566718 AF157833 NM_012133 AI202415 AK002086 AF207598 AI214562 AI202184 AI865579 AA603481 AA483808 AA909166 AA774034 AW748102 AW176026 AW351472 BE164787 AA970983 BE622521 BE389817 AW366336 AW366328 AW366327 AW366329 AW366335 AW366337 BE269711 T11249 T11264 BE253295 BE256412 BE250882 BE255440 BE257663
30	303149	97393_1	AW963315 AA312995 AA037152 AA088607 AA064770 BE088067
	303306	11887_1	AB037732 AW503898 AA215297 BE547488 AW177355 AA046224 AA361664 AA773328 AW512704 AI283330 AI307357 AI138263 AA046116 AI219874 AA315431 AW169999 AA492006 AW298002 AA043140 AA131781 AA292383 AA031721 AA027867 R31381 AW023352 AI686186 AW467416 AA493914 AA483019 AA483081 AA040871 AA558288 AW070397 AW572828 AA693439 AW206584 AA761354 AA907254 AI671019 BE221791 AI915828 AA744724 AA027815 AA131769 AA031641 AA837288 AA737401 AI765196 AW086076 AW873024 AI567164 AA744556 AA888910 AI572276
35	303443	224022_1	AA320525 AW025411 AI684617 AI653685
	303502	325188_1	BE174240 AA488528 AL042253
	303582	647662_1	AA377444 AI458965
	303610	226089_1	BE247299 AA323288 AW966142 AA334916 AL046572 BE145095 AW751265
	303642	284260_1	AW299459 AA417112
	303777	244977_1	AA348491 BE246984 AW505247
	303839	1770217_1	Z45939 T54414 T06550
	303874	5013_1	AL050333 F08138 Z43325 H13393 AA258921 AA224232 BE439918 AL050018 AW363692 AA236615 AA746291 Z19312 AA428674 Z82579 T32527 AW952956 R59046 AA403173 AA403171 AW023058 AA461143 BE149531 AA428185 AI382812 H42659 AA406086 L48858 AW630177 Z24777 AW675297 AI393859 AI743022 AI669354 AW803015 AA401255 A952901 AW043848 AI808787 AI140662 AA194627 AI140997 AA007454 AA007318 AI469859 AI540581 C06482 AI277356 AI458423 AA460839 AA861452 AI080197 AA630781 AA845367 AI125582 AA411705 AA970524 AA699910 AW804640 AW805007 AA724226 AI128207 AI696852 AW673064 AA748404 AW771788 AW088185 AI026976 AI537560 AA224233 T24024 T50208 AI827319 R17235 T11904 AI816830 R41845 AA639828 Z41214 AA258158 H06057 F02752
40	303886	81595_1	AIW365963 BE141537 BE141535 BE141538 T19123 R57434 Z43870 AA298099 AA298004 AW963314 AI627790 AA298160 BE501485 AW271198 AA195563 AA195584 H28868 AA004370 Z42582 R21338
	303929		AW470753
	304084		T91986
50	304143	30606_-14	R88737
	304165		H73265
	304183		H91161
	304236		W93278
	304350		AA186871
	304439		AA398882
	304495		AA446448
	304518		AA461438
	304521		AA464716
	304546	14011_1	AF113676 X01683 K01396 M11465 NM_000295 K02212 J02619 AB004044 H60588 T72131 T74637 T70970 T73183 T62154 R93629 H50855 H80585 H78044 T69186 R95698 H59327 T54018 H83071 R99626 H89864 H91798 T72841 T71108 T72812 T54005 T50896 H56102 W1486 W01669 AA137076 H90340 T61854 T61840 N93360 T61844 N53576 T55852 X02920 H56350 T58720 H56351 R92748 H56914 H59279 H50665 H56928 T69144 H80448 R91066 H77829 R92479 T55014 T52174 T67613 AA845231 H95664 V00496 M26123 AA484470 AI114833 T58716 D64752 T50876 T67858 H48675 T51161 T70409 T61715 T72289 H51854 H72171 T50834 H81483 T72132 T58792 T51179 T72833 R29662 T60563 W23562 H94193 T55017 T74830 H78469 H90811 T61303 H75867 T71527 T68360 R91065 R91079 T71172 H52923 T50871 T86567 H94691 T69226 AV649730 T46850 T56587 T46849 T60552 BE043578 BE042051 T72296 T61001 T58918 T52107 H82324 H47453 R06725 D16856 T48282 T52250 R92117 AI287339 H73203 AA318670 D17206 H66626 T69268 H73485 R93078 H73533 R87097 T71529 AA885254 AA486074 R94242 H74033 T3643 D12131 AV655901 AA345387 W07278 AW371443 AW371484 AW371427 AW371435 AW371467 AW371474 AW371471 AW371453 AW371448 AW371468 AW371486 AW371521 AW371463 AW371485 AW371460 T55283 H58030 H68955 AA360298 H73996 T58919 N94213 T50911 H51045 T56077 T72988 T71819 H60101 T72439 T68079 T73548 D11609 T61008 H67597 N49781 T73190 H50843 T73140 H61124 AV657376 N39304 AW075086 AI247165 H73123 H60258 AA343450 X17122 AW470706 AA336649 AW392737 H75576 N76963 T64227 AI803294 W73727 AV649563 AI307406 AW075080 D11525 AI032826
60			AA486189
	304636		AA524031
70	304677	2822_15	AF119850 AI133021 BE561919 BE617370 BE61149 BE561526 BE616541 BE561473 BE618997 BE549320 BE515105 AA471325 BE61284 BE559668 BE547034 BE548383 BE299097 BE545965 BE614687 BE615783 BE561441 BE562777

BE515345 BE614972 BE560948 BE561499 BE615787 BE267297 BE545452 BE300347 BE514432 BE615122 BE561421  
 BE560145 BE618939 BE544951 BE395272 BE619057 BE396645 BE563309 BE544527 BE408111 BE615925 BE254022  
 BE549220 BE562856 BE279501 BE616173 AA486917 BE560738 BE547802 BE548127 BE561779 BE314412 BE542474  
 BE249907 R35879 BE561536 BE544036 BE549153 BE560885 BE545111 BE256731 BE548221 BE619339 AA130747  
 BE259251 BE294371 BE546522 BE548227 BE612845 BE543601 N87541 H45619 BE547387 BE259822 W28818 BE542131  
 BE547778 BE292868 AW580223 BE513458 T60201 AW364101 BE293099 BE296447 BE270504 AW794068 BE266461  
 BE617287 BE277701 BE265741 BE267576 BE293068 BE265316 BE253499 BE261049 BE513754 BE252038 AA652484  
 BE255820 BE171777 BE393701 BE253247 BE564673 BE261164 AA308384 BE314208 BE293686 BE393327 BE263931  
 BE019750 AW937098 BE315499 AA079590 AA247598 BE387154 AW601680 AA152060 BE253235 BE258730 BE076040  
 BE549252 W05061 NM\_001404 X63526 BE314801 BE397574 AA353735 BE622279 BE561832 BE256022 BE297838  
 BE542783 AA346189 Z11531 AA159202 N27603 H68019 BE616749 BE249954 AA206881 BE619172 AL079889 T50884  
 T73176 H84175 N42762 AA190934 AA199914 BE541310 BE267209 BE304598 BE256955 BE257013 AA213850 BE546575  
 BE410825 BE257518 BE304451 BE392426 BE249978 AW938636 BE280611 BE269434 AA018801 BE249981 AA356501  
 BE546691 BE543217 BE535759 BE548149 BE274370 BE298124 BE621502 BE548075 BE562746 BE383394 BE262657  
 BE544530 BE619215 AI904614 BE546044 BE256949 BE560570 BE538936 BE548558 BE269263 BE296993 BE260919  
 AW862760 AA011045 BE563467 BE396985 AA934616 BE378952 AA100761 BE251832 BE254887 BE545870 BE396771  
 BE543862 BE538869 BE266240 BE542730 BE383556 BE502067 BE552014 AW602866 BE261720 AA016106 BE258671  
 BE281263 AW882888 W56843 AA090219 AW883193 BE618734 BE539981 BE616532 AA338063 BE618675 M55409  
 BE267147 BE545495 BE622854 BE618741 BE299535 BE512709 BE619549 BE621826 BE619650 BE271084 BE304968  
 BE620704 F00658 BE278020 BE543196 AA359668 AW959325 BE296029 BE256156 BE613346 BE617836 BE304900  
 AI142782 BE619732 BE621566 BE396956 AW887816 BE168703 AI219282 BE613404 BE088705 AI379660 AI620510  
 AA659784 W79449 AI767892 AA908586 AW863561 AI142797 BE278846 AA299822 AW954154 AA826854 W32788  
 AW801176 BE257998 BE569129 AA167694 H95000 AI431695 AA613987 T53824 BE514817 AA227510 AA877445  
 AW882770 AA507451 AW820977 BE174939 AI200307 AI445023 AI569411 AA548618 AA724243 AA653140 AA522801  
 AW578153 AA177136 AW192960 R76536 AA548071 AW969645 N86444 AI273891 R49511 AA860967 AA916058 AA826867  
 H25586 BE548884 AI911255 AA199842 AA115835 AA976259 H69325 AA055300 AI474601 W60531 AI719943 H26633  
 H67377 AW410942 AA133585 AA069276 BE260996 AA668262 AA172175 AW973378 AA084253 AA989530 AI250640  
 BE168951 BE564839 AW440725 BE564468 AA070688 BE315049 AA224349 F31228 AW517818 AA523027 AA190811  
 AA161089 AA088923 AA468923 AA129110 AA205266 AI220801 AA181780 AA121359 AA483551 AA501345 AA081217  
 AA865545 AA151850 AA604041 AA583670 AA807148 AA056615 AA486229 AA206889 R43764 AA187051 N88700 T49757  
 AW247901 AA582333 AW194553 AW387155 AA086260 AW166834 AA535771 AA001541 AI275743 AW589611 BE539505  
 AI472623 AA724240 AI270077 R26806 AA018915 AW979353 AI702378 AW591470 AW015803 AW265338 F27619 F37352

304679 AA548741  
 304747 AA577816  
 304782 AA582081  
 304813 AA584540  
 304852 27797\_1 NM\_005534 U05875 BE244633 AW163768 AW163746 AI879080 U05877 BE274096 BE390182 BE409310 BE302866  
 AA03965 W31129 AA496185 F07599 Z43842 R24838 N29198 AW877047 AW877121 AW877125 AW363656 AW877123  
 AA377595 BE207637 W20297 AA315151 W35239 T89516 AA058408 AW408191 W23815 AA448929 AI793330 AA173035  
 AI567425 C02282 AW602173 R73773 AI870491 AA287175 AI140367 AI025534 AA370611 AI824138 AI300591 AI300537  
 H04055 AA039282 AI921705 AA738464 AA376968 AW474501 AW157734 AW190741 AW474549 AI590825 BE302836  
 AA128344 AW151252 AI813665 AW514266 W78209 AI368371 AI41996 AA834439 AA085743 AA779622 AW264599  
 BE313816 AA470375 AA572697 AI970559 AI445034 N91836 AI160431 W72709 AI248975 AA757418 AI719633 BE300540  
 AI811310 AI368446 AI619458 AI186860 AA037665 AI126764 AI985833 AA579211 AI126756 AI140467 AA470989 AI188863  
 AA913322 AA825360 AA848066 AA683483 BE206448 AW008445 AA741327 AI022347 AA740593 AW117190 AI906053  
 AA843509 AI139173 AI733305 AI492747 N50093 AA862085 D20407 AI094789 AA449440 AA719906 T89696 AA838546  
 AI140848 AA503840 AI580923 AA828526 AI970569 AA993118 AA082836 AA825533 AA699833 AI263685 AA922853  
 AI160301 AI078729 AA570145 BE046096 AI371141 AI436611 AI186016 AI088757 AI141470 AI187126 AI342299 AI373836  
 AI827816 AI201090 AA970120 AA551892 AA287086 AW841869 AI492741 AI073355 AI192343 AI344429 AI879541 AI423770  
 N94310 AI051206 AI344504 AA854553 AI126700 AA775124 AW572702 W19288 AA748746 AI344385 H38212 AI130719  
 AA626341 AW771310 AW019997 AW771346 AI922093 AA75114 AA445452 AA703616 AI362485 AW072333 AI186652  
 AW675704 AI638453 AA709431 R43128 AA588595 AA832391 AI865833 R73681 AA515501 AW235701 N94608 AI798771  
 BE045450 AI369476 AI886680 AW197390 Z39908 H03368 AI766761 T28478 AI424369 F03847 N90677 F32611 BE041313  
 AI420363 AW008689 AI933460 AA614856 AI401046 AA056737 AI620906 AI797411 AA039629 BE243267 W23784 AI311585  
 W70312 AA081965 AA643695 AA541686 AI249750 AW298157 AA939276 N63266 AW000886 AA678042 F19175 AI499969  
 AI924421 T25501 Z20802 AI972829 AI826621 AW891571 Z20801 W26292 AW190699 AI640652 AA970137 W77882  
 AI567514 AI570317 D45596 AA905904

304941 AA612612  
 304948 AA613107  
 304987 AA618044  
 304998 AA621203  
 305081 AA641638  
 305089 AA642622  
 305211 AA668563  
 305257 AA679005  
 305259 AA679225  
 305271 33276\_16 T81656 AI174932 T82847 T19334 BE563893 BE563101 BE407270 BE386893 AW075527 BE616551 AI879322 AI929728  
 AI557023 AA657885 AI207449 H74068 AA135820 BE560404 BE250148 BE619362 BE612714 BE276295 X55715  
 NM\_001005 T71701 BE513229 BE385149 BE387298 R27196 T55650 AI546980 BE276020 AA483994 AI525170 BE384198  
 AA158297 BE206964 AA259269 H51936 S42658 AA187098 AA186388 AA205135 AA533926 BE083862 BE563679 N88835  
 AA096117 AA225077 BE543870 U14991 U14992 U14990 W40297 W17363 AA377786 AA199796 H14178 H74025 W05492  
 AA187571 BE537410 BE277109 BE515193 AW674485 AI433617 AW797409 BE297553 T52590 BE264322  
 BE407417 AW583518 BE392397 AA564507 AA187843 BE620435 BE274727 BE544676 BE559616 BE407160 T59911  
 BE083719 AA375543 N58722 BE562491 BE616109 AA876152 AI439321 AA570592 AA381913 AA381480 AA381930  
 AA381960 AA381565 AA381447 AA381519 AA381183 AA381267 AI453809 AA652710 BE378907 AW276979 AA211342  
 T48442 T56312 N42678 AA134859 BE538136 AA166796 AI698339 AI679955 AI679515 BE551797 T56162 BE409734

BE397723 BE410190 N54738 AA176837 AI200414 AW005899 T90127 BE383057 AI193324 AW265248 AA563942  
 AW970872 AI110677 R66333 AF063551 AI433082 AW162201 AA533266 H46197 AI347056 H80563 R38464 BE378960  
 AA642986 H51899 AA578725 AW172802 BE043720 BE041229 BE043205 AI345540 AW301354 BE041779 BE043807  
 AV302305 AW301424 AW268315 AA588176 BE171107 AI312967 AI148696 AI307710 AW272066 AW302329 AI054237  
 AI054050 AA588383 T39250 AI546935 AI174280 AI589974 AI246958 AW301874 AI306015 AI284741 AI349605 AI307731  
 AI340494 AI340550 AI251655 AI370084 AI224265 AW271029 AI224705 AI343155 AI224236 AA076493 BE392932 BE387057  
 AI537940 AA738136 AA744193 AI933432 BE043117 BE392795 AW452224 AA860693 AI193219 AI273441 AW748113  
 AW883829 N84473 AW249041 AW970861 AA079174 T50227 AW576854 AW883209 AA420780 AA076320 AW883211  
 AW883853 AW883637 AW883470 AW883331 AA156171 H26617 N90099 AI371291 AA181064 AI286302 T52479 R94956  
 AA678887 AA158113 T54355 N71149 AA18675 H71939 AI308306 BE620398 AA523799 AI094727 AW411258 AW381687  
 AA494005 T48633 AA976344 AI472484 AI500002 AI497712 T70949 AI005589 AA555014 T47971 AA679895 AW572711  
 AI345573 AA541349 AA885550 N69050 AA664708 AI348824 AW072863 AW303049 AI784034 BE138654 BE139281  
 AW072723 AI312363 AW572720 AA177047 AW327859 AA713813 AI908759 AI474689 AI890275 AW606076 AA523658  
 AA523366 AA522640 AA340801 AW268340 AA730765 BE349872 AI014737 AW300665 AW392212 AI040590 AA046630  
 AA045165 AI972954 BE550191 AA044995 AI191439 AI191437  
 305295 AA687131  
 305296 AA687181  
 305309 15215\_2 N25117 BE567374 AA188062 R76474 BE394149 AA346106 BE268871 W32999 AW793468 AI075912 D53484 AW512628  
 AA829843 AA831923 AA187162 AI375868 AA576000 U41448 NM\_001029 X69654 AW068471 AI262598 AI719560 N24138  
 AA588388 F22286 AA385512 AA155919 AI347783 F22561 AI016457 AI708902 AA622797 AW084953 W35385 AA845254  
 AA683406 AA773365 AA443835 AI626039 AA175725 AW900657 AW900718 AW148625 AA187280 AA583906 AA916044  
 AI500256 N75427 N87626 AA921883 N89069 AW327807 AI377311 AI735263 AI039117 H51501 AA384041 AA908565  
 AI022990 AA530944 AW169092 AA640128 F31984 AI421320 AA902400 T47242 AA187283 AA676731 BE466262 AI358868  
 AI829098 AI720169 AA602834 AA075051 AI016333 AI832618 AI626112 AW028704 AI807675 AI554601 AI735043  
 AW275957 AA135844 AA102517 AI202054 AA836808 BE536365 H71341 AI1719486 AI497920 AI581707 AA601142 AI626130  
 AA991422 AA344639 AA740155 AI350162 AW780123 AI056774 AI057132 AA532368 AI524528 AA720807 AI553829  
 AI688278 R76475 AA136006 AA614598 AA224772 AA181050 AI278337 AA583702 AA932358 AI708568 AA657866  
 BE173137 BE174600 AA181944 T48413 AA718293 H87639 T86323 T86324 AA885811 N35961 AA186975 AI221285  
 AI707796 AW273582 AA775295 AI719122 AA091156 H23786 AA225182 AA779355 AI718506 AA148325 T47243 AA358519  
 AA001821 F28492 AA737274 F25722 AI038467 H51521 AA576106 F35015 AA548697 BE073926 AA748127 T48414 D54761  
 AA863282 AW384185 AA086265 AA508410 AW297113 AA342512 AI884745 T55298 AA758944 AI719674 AA501428  
 AI718704 BE165424 AA961799 AA962669 AA983161 BE173686 AA886117 AA872983 AW793184 AA665189 T55130  
 AA613898 H91152 AW797007 AA071280 AA071377 AI880677 AA659794 F35193 C15194 T51001 T70211 AI625921  
 AA908590 AA551606 T69882 AA077153 AA346105 AA225274 AA358552 F18375 AW002113 D55051 AW368652 AW368778  
 AI721092 AI335641 AI202753 T81175 AI720731 AA747418 AA304776 AW022506 AI299308 T25496 AI749286 AI460263  
 D53639 AI124989 AA593610 T68867 AI460138 AA358484 AW975054 AA321580 F28232 AA353257 AA303644 AW068878  
 AW068730 F25874 AA300373 D56049 F26553 BE568636 AA385266 AA316385 AA303470 X77770 AA090283 AW327251  
 AA316866 F37411 BE385312 BE253018 AA320071 AA308568 D51956 T27782 BE393178 D51473 AA364881 D52094  
 AA303753 AA314074 AA369588 H91202 D52058 BE393600 BE389958 F33433 BE389989 F28039 H51520 AW239409  
 F32141 BE304844 AA316043 F30160 AA352329 F19450 F19208 AA187556 AA961772 F31331 AA075050 AA187320  
 AI374585 F27581 F32548 AA308515 T51089 AW964109 AA128451 BE386435 H51500 AA570763 AA224794 AA157613  
 T68942 AA096244 AA186484 AA360942 W78176 AA147724 AA699717 AI052202 AA095483 D52120 T25967 AW880072  
 AA903320 AA082357 H71393  
 45 305350 AA706676  
 305369 AA714040  
 305371 AA714180  
 305611 AA782331  
 305630 AA804508  
 50 305700 AA815428  
 305715 AA826884  
 305726 AA828156  
 305731 AA829363  
 305750 AA835250  
 55 306004 AA889992  
 306026 AA902309  
 306069 AA906983  
 306090 AA908609  
 306143 AA916314  
 60 306235 AA932299  
 306275 AA936312  
 306302 AA937901  
 306308 AA946870  
 306336 AA954198  
 306358 AA961821  
 65 306380 24593\_27 BE271766 AF116719 BE513150 H49670 BE397881 BE614531 BE562280 BE387230 BE270556 AA333494 U43901  
 AW630924 U43901 BE620882 X15005 AI630308 AA224482 BE285468 BE566970 BE542957 BE567127 BE564341  
 BE257276 N85360 R47201 AA130008 BE393122 T57407 BE304440 BE538285 BE543610 W04320 BE261521 BE395193  
 BE227139 BE513224 BE542480 BE298611 BE542479 BE545346 BE546436 T70965 BE620480 J03799 BE397891 BE562251  
 AA373403 T55389 BE393268 BE562409 AA324299 AA317559 BE393578 AA064690 BE562188 AI525344 AW672959  
 BE561902 R19651 BE616792 AA308270 BE563600 N31380 T74744 T58212 N31805 BE542361 AA357300 BE543041  
 BE560993 N76928 BE389448 BE262375 BE538256 R56424 R20398 BE615985 S37431 T72413 NM\_002295 BE281623  
 AA076198 AI205269 AA334045 T55370 AW800811 T57971 AA134763 AA209447 R19067 AA053200 BE542788 BE299054  
 BE268607 N49627 T51676 BE561411 BE619391 AL031311 BE388868 BE279523 BE560315 BE271784 BE253962  
 BE619010 BE258112 BE382896 BE535693 BE620389 AA204831 BE267276 BE547775 BE249951 BE537902 BE298296  
 BE537758 BE612439 BE267227 BE295417 H81474 AA083251 BE545562 BE279950 AW389264 BE545054 BE294352  
 BE382915 BE259414 BE542092 BE298128 BE273820 BE271588 BE253890 BE563184 BE546606 BE263148 BE614968

5		BE542402 H67612 BE560882 BE546130 BE543147 BE540497 BE537524 BE613753 BE547997 N50015 BE614047 BE618974 H94126 AA113031 BE546168 AA102514 BE565376 BE543414 BE272689 BE545988 BE537457 BE617006 T86726 BE512888 X61156 BE560027 BE538834 AA076331 T59466 BE559529 BE257047 T59764 AA191701 AA1385844 BE562585 AA366436 AA335298 BE019631 BE378577 BE535521 BE396621 BE279958 BE549122 BE537798 AA1338812 AA961833 H21616 W04906 AA928220 BE548328 AA089878 T71120 T56899 BE542669 AA082136 T82381 T94025 H81332 AA772988 BE545437 R56691 H93222 H70111 AA090016 M55990 AA184974 BE620968 BE539322 AA160696 AA346768 BE389278 N86754 AA092482 AA086091 N83880 AA147808 AA147849 BE544762 AA342408 AA204832 AA147998 R28783 AA328498 H79896 R91237 AA301536 AA206450 AA358599 N84489 AA094379 R29091 AA206157 AA489490 AW889947 AA167032 W19474 AI241644 AI432647 AA665658 AI718549 AA737795 AI207622 AA599406 AW514994 W35312 T59884 AA643137 AI200347 AW591918 T57822 AI190984 T59412 AA658064 AW973080 AI370110 BE176259 AA730904 AA872869 AW869253 AA227371 AA968861 AI281565 M14199 AI266346 T55798 H99950 W23712 AI033135 T96423 AA128434 R97053 AI266269 AA114134 AI264933 AA593164 R99334 R44109 R71185 H81360 AA532483 R56847 H66334 AA083252 AA135211 R44437 AA320995 AA983284 T96327 AA622534 D51200 AI469759 N89876 AA129995 AI168036 R43516 AA835456 AA508702 AA989565 AA552538 AA491480 T56856 R58891 AA112376 H75933 AA099846 AI553658 AI538107 AA197339 W40255 AI358440 AA506867 AW176499 N89447 AA548318 AW089158 AW800311 AA205200 H47125 H61801 AI817550 AA885896 H61800 AW516691 AW089792 AA206057 AW411349 AW301332
10		306413 AA973288 306462 AA983397 306469 AA983792 306501 AA987294 306563 AA952926 306610 A1000635 306723 A1026151 306744 A1031882 306792 A1042426 306810 A1057294 306820 A1074408 306896 A1093383 306953 A1124971 307010 A1140014 307012 A1140212 307034 A1142526 307041 A1144243 307056 A1148675 307064 A1149335 307074 23309_6 BE171526 AI159895 AI362169 AI557175 AI816211 W84586 AA443432 AA478090 BE273860 AW067965 BE616889 A1905033 AW161521 W07464 AA402174 T72985 AA086142 AI816076 AI929055 AW404703 X05607 NM_000099 X52255 A1362017 N31169 AW964703 AI346361 AA430624 AI419306 AI089436 AA478163 AI422740 AA405246 AA581187 M78181 A1A11423 A1569697 AA456462 AI056312 AI394439 AI056388 AI394465 AI419320 AI096478 AA554829 AW003877 AI394482 A1149531 AI014323 AI096480 AI816174 AA454916 AI050879 AA723285 AW055076 M27891 X61681 AI564229 AA464371 A1809078 AA284648 AA994535 AW026440 AI421217 AA379815 AW026352 AI218999 AA713818 AI424074 AI003039 AA947130 AW051798 AA371755 AI633210 AI342529 AW007903 AI949112 AA776633 BE379201 AI361017 AI589045 AW026349 AI421109 AI361213 AI356896 AI356628 AW027415 AW005034 AW304939 AW175887 AA622325 AI206007 A1422765 AI953860 AA055990 AA363573 AW027888 AI498831 AI937335 AA385758 AW880771 AA523539 AA315198 AA371942 AI420662 AI418260 AI422729 AI424095 AA335397 W93398 AA335301 AA336083 AI088933 AA933630 AI207597 A1422814 AW880787 H46870 AI422104 AI784528 AI422028 AI423492 AI418908 AI476176 AI418491 AI262265 AW054997 A1401222 AW935107 R75680 AI356763 AA086053 AA320998 AW959124 AW951566 AA582437 AA618600 AI928791 A141394 AI380094 AI077661 AA570651 AI418220 AA586896 AW051500 AW628321 AW168616 AI423428 AA775816 AA464260 AA363434 AA808067 AI356585 AA365044 H39961 AI421421 AI084346 AI421590 AI678845 AI401337 T68178 AA533499 AI420682 AI422976 AA7040329 H24665 AA452987 AI041100 AI423091 AA704378 AA482076 AE688207 AA588778 A1582532 AA594350 AI799230 A1817567 AA365706 AA526950 AA325184 AI6247928 AA489696 AI148572 AA854613 A1499904 AA320404 AW005422 AA533158 A1469702 AA291655 AI149169 BE327985 AW328286 AA922081 AA729897 AW799758 AA364028 AA843331 F20435 AW001524 AI285253 AA916618 AI422358 AI094747 AI094754 AI262629 AI597587 A1281139 AI096859 AI537257 AA846506 AA973472 AA845740 AA977830 AW157016 AI174258 AI460190 AI089868 A1095037 AI089864 AA723298 AI074244 AI096812 AA687428 AI422334 AI359247 AA782354 AI460199 AA454568 AI096464 AA426217 AI097391 AA434378 AI097359 H93031 AI160095 AI373821 AA477632 AI564398 AI150904 AI346637 AW026849 A1A573822 AW192676 AI3639739 AI094919 AI290053 AA319991 AI627370 AA702191 AA363695 AI096607 AA992885 A183336 AI361555 AA890199 AW768597 AI421336 AW001458 AI095758 AW027909 AI150989 AA620294 AA974335 A1306477 AI146532 AW103696 AW025355 AA588754 AA772569 T51534 AI077588 AI051136 H45665 AA641555 AA335249 A1359662 AA335173 AI160230 AI359451 AI096516 AI832913 AA856626 AW129193 AI057277 AI299349 AI801789 AI360811 AW026246 AW009786 AI146563 H54066 AI141357 AI290397 AI359760 AI073390 AA369438 AA782546 AI360379 AW1561525 AW026180 R73226 AI057139 AA918124 AA652886 AI359602 AW008584 AI361599 AA858102 AI075082 AI361575 AI360860 A1363088 AI095068 W74631 AI369419 AI359287 AA617644 AI027185 AI721209 AA579636 AI535703 AI535742 AA411125 A1535680 BE206595 AI076856 AI084271 AI356202 AI076863 AW104687 F22288 F37131 AA135467 AA722050 AI283930 AW193140 AA584226 T68253 AI362193 W46392 AI419269 AA533764 AA565340 N29369 AI282034 AI074482 AA464433 AA335520 AI816542 AI311529 AA402731 AI361536 AA936821 AI090604 AI278884 AI272311 AA454842 AW511753 AW316946 AA577547 AI625298 AW008545 BE206455 F35252 AA478251 AA894702 AW055036 AA318997 AI051266 AA916075 AA468716 AW575608 AI097241 AA478250 AI813393 AA961716 AW168580 AI362156 H56144 AI351645 A1W051887 AA464434 AA782238 AW069713 AW328287 AI361335 H22417 H28467 AA774593 H68981 AA948632 AA987632 AW026296 AW069809 AI270651 AA290771 AW027914 AA402315 W02152 H23679 F34337 AI367943 AI422932 AA464435 AA630257 AA630372 AA216227 AI354596 AA725375 AI832369 AI864731 AA642707 AI143086 N90262 AI471066 AI475343 AA478008 N81104 AA838768 AI147250 AA195967 AI097264 AI418176 AI041946 H64634 AA304727 AI245016 AA947243 AA320023 AA411983 AI285077 AI055044 AA602808 AA954547 AA844184 AW900840 AI356806 AA934005 AA658077 R54958 T29906 AI983056 AA364947 F27923 N69886 AI418676 AI149599 AI418632 AI636479 AA114824 AW168203 A1A584263 AA290780 AW008659 AW149229 H18245 AA559148 AW008735 AI088875 W84605 AI149397 AW515564
15		
20		
25		
30		
35		
40		
45		
50		
55		
60		
65		
70		
75		

AA722497 AI422059 AI394499 AI475192 AI589945 H24677 AA548688 AI130975 AI130862 F30448 AA313754 AI148377  
 AA996252 AI915211 AA886378 AI371520 AI879469 F24100 F28104 AA503929 F19709 H28541 AA993350 AA320635  
 AI783926 D11806 D11707 T60062 H62085 F36915 H19496 AA558311 AA485384 AA037016 AA435523 T60162 AA602795  
 AA477963 AA335919 AA335626 AW008995 AA991946 AI554752 AI370939 AI400912 AA250788 AA320310 AA99770  
 AI13034  
 5 307084 AI160527  
 307095 AI167910  
 307156 AI186762  
 307243 AI199957  
 10 307264 AI202211  
 307288 AI205169  
 307391 AI225058  
 307542 AI280859  
 307565 AI282468  
 15 307586 9205\_7 AW367324 AA513829 BE300206 AA134200 AA126689 AW975636 AA375190 AA229666 BE562156 AW957307 AI133468  
 BE562075 BE293171 H43213 AA095435 AW405074 N28598 AA579221 AA096022 AA315868 AI557910 AA096172  
 BE616496 R83253 NM\_006013 M64241 BE386761 W03711 BE261517 BE298374 BE278491 BE255200 BE304448  
 AA206687 BE386934 BE618525 T49742 AA532479 H92515 BE299890 N84711 W80584 W04247 AV654899 BE313651  
 AA301932 BE616507 BE568551 H18202 W93773 S35960 M81806 AA352773 BE281085 H69766 N87825 AA090275 M73791  
 20 AA090218 H79923 H85682 N89270 AI001784 N83949 BE567780 R73099 H63621 AA714817 BE536488 BE407585  
 BE394061 BE619086 T71244 BE563254 C04286 BE293916 T67271 BE544286 BE538828 S64169 BE254604 BE295847  
 BE407999 BE616137 BE618994 AL121934 AA084264 AA084260 BE615364 BE541737 R30878 BE537874 BE293082  
 BE622544 BE280736 BE560886 BE512983 BE512970 AI469402 BE279821 BE615519 BE563424 AI027614 S35959 T67270  
 AA377416 H20002 W04857 AB019572 H91139 BE279255 W02004 W05580 AA524537 AW970052 H91379 AI223104  
 25 AI219245 AI190973 R73100 AA618062 AA583590 F37254 AA054426 AW951771 AI582983 AI535778 AW183619 AI285499  
 AI205618 R18051 T50234 BE252578 AI025348 AA738325 AI190989 AW732196 AA531213 AI439364 AA292351 N70489  
 AA126581 AI5677584 AI264944 C14996 AI351267 W65440 H68843 AA548110 H60635 T50114 AA187603 AA404627  
 AA614381 AA835418 AA192999 T49333 AI246998 AA070742 AW388661 T49876 AI865171 BE261134 T50166 AW752163  
 30 A1866246 AI497721 AA022533 AW388584 C05870 AW388551 AW375685 AA079408 AW388685 AW388565 AW388289  
 AA346577 AW752158 AW752161 AW752157 AW752160 AW388563 AL031276 AW375687 AA528266 AW578215 BE250977  
 H66751 AA054338 AA084261 AW972379 AA084265 AA446308 AL046840 AW580988 T50626 AA634027 AA492347 D51630  
 BE261889 AW581036 AA302508 AB007170 AW752162 AW375688 AW375709 AW752152 AI499767 AA936623 AW455210  
 AW388395 AW388383 AW581024 AW388313 AW797015 T25776 AA503342 BE378698 AA501932 D17268 AA493357  
 AA932910 AW075203 BE171178 AW388379 AW385863 AW388335 AW581005 AA369640 AW388419 AW388324 AW579474  
 35 AW388588 AW388719 AA582262 AW388360  
 307612 AI290787  
 307626 AI300035  
 307640 AI301992  
 307646 AI302236  
 40 307783 697809\_1 AI347274 AW844024  
 307856 AI366158  
 307954 AI419692  
 308047 AI459633  
 45 308050 9482\_1 H18835 R47363 AI460004 N31660 AA454774 AA551759 AI417040 AA694490 AA633315 AI344661 AA708532 AA878567  
 AI802702 AI913465 AW001160 AA932133 AI092908 AA026974 AW628573 AA592910 H18836 AI274428 C00675 AK000048  
 BE313619  
 50 308106 AI476803  
 308139 AI494477  
 308186 33276\_16 T81656 AI174932 T82847 T19334 BE563893 BE563101 BE407270 BE386893 AW075527 BE616551 AI879322 AI929728  
 AI557023 AA657885 AI207449 H74068 AA135820 BE560404 BE250148 BE619362 BE612714 BE276295 X55715  
 NM\_001005 T71701 BE513229 BE385149 BE387298 R27196 T55650 AI546980 BE276020 AA483994 AI525170 BE384198  
 AA158297 BE206964 AA289269 H51936 S42658 AA187098 AA186388 AA205135 AA533926 BE083862 BE563679 N88835  
 AA096117 AA225077 BE543870 U14991 U14992 U14990 W40297 W17363 AA377786 AA199796 H14178 H74025 W05492  
 AA187571 BE537410 BE270902 BE277109 BE515193 AW674485 AI433617 AW797409 BE297553 T52590 BE264322  
 55 BE407417 AW583518 BE392397 AA564507 AA187843 BE620435 BE274727 BE544676 BE559616 BE407160 T59911  
 BE083719 AA375543 N58722 BE562491 BE616109 AA876152 AI439321 AA507052 AA381913 AA381480 AA381930  
 AA381960 AA381565 AA381447 AA381519 AA381183 AA381267 AI453809 AA652710 BE378907 AW276979 AA211342  
 T48442 T56312 N42678 AA134859 BE538136 AA166796 AA698339 AI679955 AI679515 BE551797 T56162 BE409734  
 60 BE397723 BE410190 N54738 AA176837 AI200414 AW005899 T90127 BE383057 AI193324 AW265248 AA563942  
 AW970872 AI110677 R66333 AF063551 AI433082 AW162201 AA533266 H46197 AI347056 H80563 R38464 BE378960  
 AA642986 H51899 AA578725 AW172802 BE043720 BE041229 BE043205 AI345540 AW301354 BE041779 BE043807  
 AW302305 AW301424 AW268315 AA588176 BE171107 AI312967 AI148696 AI307710 AW272066 AW302329 AI054237  
 AI054050 AA588383 T39250 AI546935 AI174280 AI589974 AI246958 AW301874 AI306015 AI284741 AI349605 AI307731  
 AI340494 AI340550 AI251655 AI370084 AI224265 AW271029 AI224705 AI343155 AI224236 AA076493 BE392932 BE387057  
 AI537940 AA738136 AA744193 AI933432 BE043117 BE392795 AW452224 AA860693 AI193219 AI273441 AW748113  
 AW883829 N84473 AW249041 AW970861 AA079174 T50227 AW576854 AW883209 AA420780 AA076320 AW883211  
 AW883853 AW836337 AW883470 AW883331 AA156171 H26617 N90099 AI371291 AA181064 AI286302 T52479 R94956  
 AA678887 AA158113 T54355 N71149 AA181675 H71939 AI308306 BE620398 AA523799 AA094727 AW411258 AW381687  
 70 AA494005 T48633 AA976344 AI472484 AI50002 AI497712 T70949 AI005589 AA555014 T49791 AA679895 AW572711  
 AI345573 AA541349 AA885550 N69050 AA664708 AI348824 AW072863 AW303049 AI784034 BE138654 BE139281  
 AW072723 AI312363 AW572720 AA177047 AW327859 AA713813 AI908759 AI474689 AI890275 AW606076 AA523658  
 AA523366 AA522640 AA340801 AW268340 AA730765 BE349872 AI014737 AW300665 AW392212 AI040590 AA046630  
 AA045165 AI972954 BE550191 AA044995 AI191439 AI191437  
 75 308191 AI538878  
 308243 AI560037  
 308249 AI560998

308256		AI565498
308337	3328_20	J00124 AA158848 AW452899 NM_000526 AW384139 BE140708 BE140690 H28274 BE140404 AW844922 BE157342 AW580043 AW384049 BE157333 BE157179 H44127 BE140671 BE614947 AW380727 BE184532 AA808075 R73201 AW949948 AW950242 BE183306 BE183297 AA587174 BE183243 AI591112 AI609229 AI590690 AW606369 AI609238 AI608848 AI590885 AW084944 AI609220 AI609210 AA583890 AI609274 AI609226 AI608878 AA550914 AW170430 AI608947 AI608606 R88146 M28646 AA587248 AA583985 AW872988 AA583993 AA586980 AI284481 AA584051 AA587024 AW800015 AA587313 AI283394 AI143314 AA595434 AA583917 AA922332 AA584021 AA583576 AA862004 AW873168 AW796060 AA862049 AW873161 AW079705 AI366769 AA583724 AA583889 D29065 AA113288 AW083705 AA584334 AA586711 AA583558 AW117878 R72295 AA583842 AW969598 R73138 H28224 R88223 AA586985 AW369022 AA602252 AA641063 R72845 AA861952 AA587149 AA587308 AW369033 AA584344 AW799955 AA583732 AW368990 AA862083 D29595 R53012 AA587254 D29462 AA158849 R72649 BE181541 D29604 AI127631 D29297 H44051 AA583575 AA586762 D29122 AA583878 AA595946 AW368880 AA587318 AW368881
5		AI611044
10		308354 792518_1 AW998989 AI613519
15		308380 AI623988 308457 AI669859
20	308462 816335_1	AI671311 BE501055 AW813781 AW813651 AW836924
25	308484	AI679292
30	308578	AI708573
35	308615 33893_1	AIK000142 AW243187 AI738593 AW505395 BE009209
40	308667	AI758754
45	308676	AI761036
50	308781	AI811707
55	308928	AI863908
60	308957	AI869642
65	308991	AI879831
70	309000	AI880489
75	309243 309398	AI972052 AW081820
	309582	AW169657
	309634	AW193825
	309628	AW293999
	309830	AW294725
	309839	AW296076
	309939	AW419122
	309977	AW451663
	310341 651442_1	AW302773 AW303087 AI254651
	310985 114598_1	T51842 T51888 T51099 T51008
	311395 252948_1	R23313 R23323 Z25059 AA359123 AW965886 BE167187 AI808503
	311736 193321_1	AA330047 AW962512 AI983896 AI590126 AA837373 AI638125 AW139338 AA765161 AW194751 AA765897 AA261818 F13656
	311965 404771_1	BE385785 T69279 AA703325 AA678398
	312129 338029_1	T87431 AW300867 T87330 AA515973 AI242970
	312339 151127_1	AA524394 AW015969 AA158731 AI831401 AI955800 AW272596 AW272595 AI281799 R34231 R35558 AW378527
	312405 765247_1	AI523875 R45782 R45781
	312996 187327_1	AW368634 AI702169 AI245179 AW368646 BE545574 AA249018 AW368633 N27553
	313078 1653847_1	N49730 N52659
	313139 255206_1	N59555 AA362113 AA362136
	313561 98898_1	BE396243 AA040818 AA134134 AA040155
	313603 199797_1	AA284333 AW468119 AA284334 AA810992
	313781 116301_1	AA079229 AA079201 AA078874
	313825 173444_1	AA215470 AA215547 AW470551
	314033 153186_1	AW752247 AA167125 AA161495
	314043 155125_1	AA827082 AA732246 AA167611 AA830741
	314138 179960_1	AA740616 AA654854 AA229923
	314183 31294_3	AW971564 AA251226 AA835824 AA748600
	314367 210554_1	AA535749 AA574021 AA302428
	314703 303778_1	AI791249 AA452038 AW817263 AW817051
	315021 344728_1	AA533447 AF158241 AI240598
	315858 406384_1	AA737345 AA682286 AI799378
	316055 409389_1	AW105663 AA693880 AW517398 AI768507 BE220851 AW978538 AA831489
	316652 454745_1	AA789249 AA904217 AA904142
	316897 474090_1	AA838114 AW629478 AA883713 AI620552
	317617 1660859_1	N58024 T58194 T11693 N64222 T05848
	318171 269002_1	AW868221 T11286 H53526 AW630814 AA381202
	318175 139564_1	AW992356 D82445 D82303 T11347 AW995841 AW995909 AA133068 AW859596 AW859595 AA644624 AW631466 AI903549 AI903513 AI903561 AI380423 AW364047 AW373615 AW373755 AA634799 AW860107 AA336073 AW960638 AI284099 AI284098 AI201463 AI972977 AA224831 BE183312 AI567336 AW389340 H39906 AW770710 AA411736 AI763378 AW340972 AW516306 AI400359 AI745530 AW511407 AA406542 AA524982 AI628029 AW008882 AA858439 AW074041 AI379597 AA132792 AW189012 AI538874 AA857364 AA224830 AW590976 AI678604 AW512052 AA593133 AI142902 AI992380 AA888921 AW517569 AI679729 AA577041 AI60743 AI282492 AA494400 AW074288 AA551421 AA904079 AA505483 AW513312 AI469669 AI872908 AI290109 AI610272 AA829570 AI611723
	318230 193526_1	AA457028 AI674737 AI688648 N92749 AI439620 AI218005 AA731321 AA828303 T12590 AI685371 AA504636 AA825573 AW172614 AA291292 AA649919 AW576505 AA262152 H29888 AA830757 AA910433 AA808481 AI971807 AI767874 AI216422 AA831483 AI566351 H41305 AA879438 AA807123

		318581	162718_1A1633044 AW016212 AW241143 AA769058 R43272 AW068958 AA210918 AA293774 AI748815 AI763294 AI333114 AI277384 AI088297 AI468477 AI824624 AW189606 AI631751 Z40749 AI984673 AI671316 AA189024 AW235412 BE178426 R24677 R40635 H05100 R40597
5	318601	366837_1	AA975853 AI915867 AW341040 AI985422 AI733001 AW300448 AA865842 AI913757 AI283772 AI928188 AA593410 AW771149 AI732663 T39921
	318647	767064_1	AI557774 AI526159 AI526153 AI541503 AI541531 AI526149 AI526018 AI526177 AI557786 AI526140 AI557743 AI541492
	318652	73477_1	T53260 AI659829 AI620887 BE619759
	318721	250974_1	Z28504 H85748 AA418060 AA356371 AW965781 W45671
10	318740	15117_1	NM_002543 AF079167 AB010710 AF035776 AA114093 AJ131757 R07930 AI243883 R62556 AA682386 AA620341 R78357 R31345 H67054 AI300045 BE174919 AW082681 N75085 AI928113 R23598 AA044216
	318975	458663_1	Z44110 F13225 R13941 BE542680 T75468 N36522 BE207328 AW136340 AA808805 AI633843 AW875667 AI637851 AI990628 AW608757 AI636889 AW136314 R58192
	319128	273945_1	H09041 H10254 AA968912 Z45974 AA393820
15	319141	1533572_1	F12377 Z46143 F05137 T74119
	319198	1535512_1	F07354 R11946 H16080
	319436	1596398_1	T70298 H58072 R02750
	319478	765461_1	AI524124 R06841 R06842
	319488	10887_1	AK001330 AA356435 BE313393 BE293644 BE251929 AA808340 BE409475 AA331948 N91096 AW402232 AW402994 AW386322 AA004739 AA459479 AW579400 W68758 AW673556 BE313041 AA455698 AL045680 AA134589 AW606254 BE301261 AW976697 AW968467 AW976703 W68453 AA004688 AW976701 AA223724 AA565953 AA215565 AA744555 AW193840 AI086227 AW970769 BE300513 AI458782 AI183406 AI309531 AA455644 AA128908 AA588705 AI138389 AI476292 AA515291 AA524425 AA459254 AA600279 AA614836 AA769786 AA492544 AL045681 AA765178 AI864425 AW780369 BE246640 AA926793 AW054669 N63744 AA206610 AA729135 AA766112 AA553635
20			
25	319551	357371_1	AA761668 AA573621 R92814 R09670
	319599	798430_1	H56112 H58047 AI630710 N58742
	319808	7069_3	T58960 AA609180 AA621130 AI927236 AA431075
	319827	776890_1	T62926 AI565392 T62778
	319884	171016_1	AA325579 AW961004 BE004785 T73234 AA209403 N54886
30	319936	57653_1	W22152 AV647377 AV647331 AA320693 T79025 F23202
	319983	1814568_1	T81429 T95572 T95563
	320001	331474_1	AA873350 T82429 T82428
	320445	132254_1	R26830 R33029 AA115761 AW118148 AI743741 AA954284 AI934165 AI088310 AI123759 AW340232 AI089180 AA700861 AI129973 AI088552 AA963119 AA359156 R33916
35	320488	368456_1	AI817336 R32883 AA595590 AI743065 R31386
	320503	25164_1	NM_005897 AF156857 AA346876 BE545147 AI003306 N45644 AW889728 BE007236
	320661	159_1	AB030034 BE304778 NM_016653 AF251441 AA307684 AW957882 AF238255 NM_016595 R57782 AA244505 AA864846 AW601475 AA232750 AI475399 AA232253 AW294490 AA626441 AW814670 AW814669 N95341 AI123874 AA100075 AW337275 AW804295 AI922069 BE161875 AI470677 AI242841 AA402558 AI435815 AA402496 AI359093 AA505991 AW197200 AA234622 AA258509 H17033 AI799498 AI263346 AA236466 AA258354 N24807 R14272 AA100160
40	320727	36759_63	AIW003360 AI971548 AA017585 X80306 X91133 AJ276100 X91132 AF064499 AF064495 AF063768 X83713 AW951310 AW975565 AA721610 AA715972
	320746	29157_2	AW975814 AA282765 AA811755 AA731129 BE219297 AA128302 R71285 AA095218
	320789	252516_1	R78712 AA603646 R78713
	320809	254953_1	W16480 AA376361 N83837 R81853 AA361779
45	320825	29807_1	NM_004751 AF102542 AW360893 AF038650 BE304708 AW360892 AW360931 AW842622 AA307800 BE292814 AW582119 AW582122 AW374998 AW374874 AI587061 AA583339 AW662377 AW192901 AW887756 AW887761 AI955582 AI150400 AA568218 AA583146 AI832775 AA294858 AI445680
50	320858	1509963_1	D59968 D81035 C15620 D80887 D81432 C15618 D60320 D80661
	320909	996651_1	D62269 AW022615
	321012	84379_2	AA737314 AA682280 AA010792 AA143573 AA953433 AA745273 AA188649 AA011221
	321253	375160_1	AA610649 AI699484 H59558
	321286	236538_2	BE245833 BE539992 AI380940 AW952644 AA535470 R84610
	321307	69583_1	AI696519 BE464779 AW296343 BE550149 AW470402 AA129660 T78937 AA342648 N71662 H82431 AI302712 AV660681 R85409 AA962323 AI680732 AA889147 AA932629 AW103527
55	321354	116028_-2	AA078493
	321370	41905_1	AJ227900 AI094933 AW051119 F00947
	321412	624592_1	AI674383 AI865710 AI201451 AI659387 U25919 BE093109 AW366305 BE141926 BE141913 AW854334 AW854342 BE141916
60	321424	107996_1	D59886 AA779752 AI655936 AW976526 AA235034 AA744353 T26888 AA235103 R96569 AA057301 AA057286 D61635 H87227
	321467	43034_1	X13075 X13076
	321524	45508_1	R39382 BE467537 AI657156 AI375103 AW021134 AI479241 BE326541 AW150836 AI684065 R35463 AA678409 AI694321 AW470057 AW608873 N62359 AI702778 AI701838 AI655208 BE465196 R51845 R38307 AW39336 AW043913 AA782285 AI205974 L13824 L23311 AI635429 L13826
65	321632	286374_1	AW812795 AA419617 H87827 AW299775 AW382168 AW382133 BE171659 AW392392 BE171641 AA541393
	321828	44964_2	R59890 R60548 N64863 AI224545 N69114 AI811204 AW518902 AI184866 AI440169 AA809472 H63089 AW952971 AW337382 AI872923 N73882 AA334161 AI537113 H63731 AI383952 N41701
70	321875	430904_2	N78520 AW606984 AI287235 AA973956 N49122
	321920	1023421_1	AW089866 N63915
	321974	1280819_1	N76794 N94221 W04156 AW897535
	322035	33334_1	AI137517 BE072492 AI127076 AW196207 AW294979
	322092	46678_1	AF085833 R69689 AW341677 AA923375 BE327566 AW630415 R69601 AW615339
	322136	46802_1	AF075083 H52291 H52528
	322175	46877_1	AF085975 H53458 H53459
75	322303	704603_1	AI357412 AI870708 AI590539 W07459
	322309	47372_1	W76622 AF086372 W72660

322331	47467_1	AF086467 W81444 W81445	
322340	47509_1	AF088076 W95222 W92523	
322394	27492_1	<p>AW068287 AA310079 BE336702 AA356318 AA306059 AA346785 AW402633 AA311210 AW402909 N76879 AW402913            AW401920 AA321636 AA354474 C17297 C16938 AA311774 M29871 NM_002872 Z82188 AW405674 H94176 R89281            AA214723 AI014482 AW949347 T27749 AW804226 AW796964 AW404581 AF077208 NM_014029 W68830 W79652            AA353375 AW575218 AA552192 AA521232 AA702695 AA033975 AW407827 AA829948 N94402 AW628604 AI523308            N57605 AA641662 H42477 N52784 AI753478 AA768493 AA845729 W47391 N55270 AI090117 R89282 BE206172            AA076650 AA595650 AI218931 BE049397 AI433110 W74114 H94277 AI358627 AI085221 AI862818 AA835967 AW103905            AI640644 AA835507 AA856887 AA694392 AW337542 AI524410 BE045500 AI440060 AI358801 AW028238 AW205248            AI718264 R48618 AA357358 AI695002 AA897549 AW081065 AI433360 AI810783 AI620963 Z82188 AA360224</p>	
5		<p>AI133446 T50819 AF147343 T50665            AF155108 AW877241 AW393512 BE160738 AW384889 AW610272 BE160915 BE160774 BE160744 AW836696 AW384919            AW836739 BE160743 BE160814 AW610275 BE160965 AW580785 AA662739 BE160941 AW821136 AW592083 AA449860            AI798661 AA310698 AW302768 AW268932 AW268741 AI250559 AW302879 AW821181 AW580243 AW384922 AW606812            AI345641 AW821143 AW384873</p>	
10	322518	38914_1	<p>AI133446 T50819 AF147343 T50665            AF155108 AW877241 AW393512 BE160738 AW384889 AW610272 BE160915 BE160774 BE160744 AW836696 AW384919            AW836739 BE160743 BE160814 AW610275 BE160965 AW580785 AA662739 BE160941 AW821136 AW592083 AA449860            AI798661 AA310698 AW302768 AW268932 AW268741 AI250559 AW302879 AW821181 AW580243 AW384922 AW606812            AI345641 AW821143 AW384873</p>
15	322610	21773_1	<p>BE242847 AA159840 NM_016216 AF180919 BE262663 BE312610 W53026 BE093965 BE004620 AW992549 AA069408            R66803 BE002445 T80130 N67797 AA765401 AA765829 AW837997 AW837993 AW838011 AW838012 AW837996            AA069435 W52118 AI457469 AA954977 R39354</p>
20	322694	35627_1	<p>NM_014125 AF090919 AF075371 AI110872 BE070571            AA086123 AA026296 AA086041 AA026297            AA777274 AI761381 AI738617 C02420            AA649792 AA640427</p>
25	322890	117967_2	<p>AA099759 AA100511 AA687172</p>
	322933	127352_1	<p>AA112573 AA112574 AA984323</p>
	322944	130324_1	<p>AI267606 AA121045 AA126521</p>
	322959	132817_1	<p>AI272141 AI879676 AF070669 W25179 AA534016 AA533386 AA010740 AF124147 W16493 W56636 AA258911 AA321677            H44503 AA642777 AA081800 W69885 H82507 AA536128 AA326782 AA326783 AA353693 AA354642 R73311 AA354400            W79820 W16502 AA301647 AI202303 AA453926 AA705795 AA011128 AA929033 AI393389 AA845133 AI445640 AI677727</p>
30	322968	17218_2	<p>A1818296 AI369820 AI539292 AI870541 W69797 AI871096 BE550803 N35853 AA640199 N27809 R49769 AA738197            AI565700 AW207656 AA587216 AA669237 AI906947 AI809956 AI740905 AW043811 AW182476 AA659844 AI742797            AI832103 AI660967 R86125 AI674667 AI808074 AI869284 AI336214 AI218002 AI338629 AI857930 AW183986 BE044333            AW135467 AI826077 AI357643 AI475486 AA478855 AW172550 AI553942 AA868731 AW268850 AI123793 AI887022            AA046935 AI361954 AI091737 AI682235 AI367076 AI088882 AI808682 AI312679 AA0469556 AW027546 AI660019 AI696174            AW008626 AI266337 AI568959 AW027409 AI040014 AW134559 AA479953 AA910082 AI301458 AW028352 AI017863</p>
35		<p>AI268915 AI185866 AI265907 AI274195 AW051540 AW027515 AI380435 AA883117 AI279396 AA846628 AW628235            AW206201 AW628510 AA954276 AI301405 AI827185 AI53978 AI200301 AI470343 AA933953 AI914937 AI362849            AW085066 AI204021 AA631192 AI351701 AA748663 AA993806 AA580146 AW027744 AA580016 AA897344 AI042638            AI473196 AA995065 AW027720 AI217421 AI935604 AW449411 AW237094 AI653348 NM_003107 X70683 AI470473            AI765137 AI193479 AI253050 AI470510 AI399828 AI371461 AI185518 N20940 R49816 N79977 W56599 N24649 W78113</p>	
40		<p>N78761 AI817673 AI911482 AW205984 AI240186 AI828016 AI942449 X65661 AW751587 AI392808 AI624192 AI950969            AA573260 AI203361 AI479942 BE041834 AW305351 AI918327 BE048713 AW071712 BE041565 AI139260 BE466360            BE502737 AW007819 AW071887 AI742130 AI344020 AW772112 AI932275 AI992189 AI197801 BE219990 AI990863            AI536934 AI336275 AI971955 AI798204 AI870429 AI652390 AI080187 BE219486 AI185434 AW628564 AW072399 AI656370</p>	
45		<p>AI498606 BE041559 AI743591 AI515805 AI087833 AI917506 AI123191 AI858043 AI334046 AI242585 AI636670 AI919478            AW771487 AI417185 AI468527 AI137861 AI554782 AI130733 AW005164 AI910551 AI189135 AI963934 AI985482 AI660396            AI497963 AW204662 AW137602 AI382505 AI493485 AI185987 AI078841 AI830054 AI378223 AI351299 AA937301            AA242817 AA258359 AW027603 AI935204 AI500360 AI569741 BE551058 AW275536 AI457854 AI142093 AW028288</p>	
50		<p>AI286002 AI279114 AI364121 AI341323 AI190436 AW002607 AI242488 AI338122 AI368600 AI340276 AI417994 AI190234            AI275527 AI934886 AI498274 AI813630 AI075339 AI087976 AI459251 AI989477 AW004046 AI992190 AI885279 AI479475            AI698030 AI473294 AI951648 AI699587 AI660602 AI873018 AW613987 AI808297 AW270159 AW572955 AW195908            AW469034 AW197100 AA885164 AW611668 AI143038 AI910560 AA418374 AW341092 AI871169 AI937136 AI204003            AA775707 AW590759 AW593350 AW572981 AI197905 AI660941 AI743469 AW237017 AI808587 AI984962 AA418254</p>	
55		<p>AI828104 AA625231 AI832151 H84232 AI240215 AI911775 AI219668 AI336801 AA232630 AI343471 W69129 N93602            AA768883 W04386 AI086277 AA983433 W07646 AA458584 N86625 AI384055 AI928089 W25479 AA242952 AI763303            AI225039 AI740896 AA953758 W69240 AA558331 AI760593 AA558712 AW992121 AW992157 W69115 BE328596 AI953190            W95311 AI950195 AI739605 AI857262 W69185 AA884586 AI198104 AI127451 AI905932 AA723310 AI936623 AI732940            AI332918 AI221396 AI336095 AI200067 AI824853 D558933 D526975 D562025 AA232764 T53299 H84555 AA076539 AA158347            BE298430 AL134493 AW732398 AW750740 AW578208 N36572 AA453861 AA252914 AA234197 AW576988 AW577034</p>	
60		<p>AA025199 AW577052 AW385538 AW576996 AW577021 T83230 AA421529 AI918492 AA909038 AA507060 AA654561            AA064597 AW001594 AI469192 AI368002 AI42435 AW379382 W93438 AA076387 AI802344 AI097013 AA987215            AA635282 W93349 AI017818 AI421564 AI158348 AI140004 AA506259 AW473184 AA236350 AI138669 H96873 AA974889            AA643735 AA995463 AA995471 AA809555 AA253225 AI298682 AI572515 T53300 AA064596 AA193589 AA025118            AI669682 AA610638 T90774 AI972322 AI280776 T27980 AW136058 BE000428 AI378691 AA961520 BE049142 AI311424            AA283211 AI344071 AI344007 AI582410 AL036314 AW798038 AI905228 C15325 AA380386 AW958417</p>	
65		<p>AW630531 BE538239 T70488 AA088296 T34175 T31626 D54331 D53142 AA029415 AW948823 AI914128 AA355446            T34322 BE006559 M85677 AA034335 T31463 AW804007 AA256591 D55128 AI535884 D55192 N23605 T31802 AA326899            AW999156 AA355201 AW999306 AI091590 BE172021 AA029490 BE000255 AW339939 AW150093 AI872098 AI274876            T06303 AA857909 N23606 AA922714 AI914104 AI285281 AW999919 AI39803 AI081354 AA972184 AI049566 AW151583            AI682455 AA088257 AI217050 BE551774 AI277033 AI252627 AA910406 AI369422 H46634 AI873113 AA033710 AW078579            AI636452 N23010 AA357263 AA256592 T05786 AA884195 AA406145 AA907807 AA482840 AI637691 AA654523 AA911495            T06601 AW594370 AW016524 C15324 AA622519 AA340191 AA174168 C13992 T69433 T96576 AW166622 T96575</p>	
70	323011	139750_1	<p>AA580288 AA315655 AA133031 AA377748</p>
	323166	162676_1	<p>AA291001 AA188974 AA290616</p>
75	323216	6526_1	<p>AA332145 AA331790 AW962563 AA868189 L13837 T34468 AA055882 AA096148 AA092327 H57062 R59098 R11247            F07659 Z44949 AF131829 L13835 T79889 AA252451 N28984 H85260 AL046384 AW995631 R58386 AI061651 AW376050            AW379789 W90347 AA450157 AI799939 AA461340 W02347 AA233095 N39675 AA659441 AW995284 W17060 R32252</p>

			AI042599 AL046385 AI970370 AA744764 AI249761 AI628106 R32668 AI863011 AI923998 AI186798 N26601 AI141864 N34992 AI377031 N23934 AI683466 BE219548 AA622032 AW089867 AA243717 N79547 R59099 AW241293 AI917545 AW103697 AI383179 AW517527 AW193642 W90348 AW381409 R11195 AA461166 AA836624 AA280285 AW242055 L13836 N89647 L13834 AI358605 AA452023 AI868391 H57063 AW075868 N20590 Z40695 R37603 R28484 AA251913 F03914 AA055772 N43752
5	323243	140566_2	W47525 AA134047 BE391212 AA330333 AA376355 BE304871 BE167342 H87402 AA631722 W45724 AA715517 AI925438 AI804849 AW241617 AW403807 AI653435 AA134048 AW747874 AI922327 AI814967 AI935895 AA228865 AW504076 AA225008 AW673858 C03914
10	323244	647858_1	AW675572 AI248270 T85161 AL133848 T70731 T69747
	323333	62251_1	AV651680 AA228883 AA367341 AW962458 AA628024 AW172426 AI767785 AA313012 AW963323
	323430	63341_1	AW062479 AW062488 AW062491 AW062480 AW938564 AW062478 AA322408 AA324351 AW938595 AW938598 BE162389 AW176556 AW938599 AW838792 AW938566 BE162305 BE162377 AW938570 AW062459 AW176555 AW938562 AW938568 AA251701 BE162320 AW938597
15	323465	193343_1	AA287406 AA261844 AA261845 AA287355 AA810895
	323479	194627_1	AA278246 AW292815 AA278703
	323538	217887_1	AW247696 BE265140 AW403615 AL037647 AA312336
	323632	333100_1	AL041844 AL040002 AL039950
	323731	226193_1	AA323414 AW664013 AI809377 AI276041 AW296883 AI798340
20	323753	12462_4	AK002161 AA327102 AI056868 AI743901 AI139018 AI199114 AI076003
	323835	506747_1	AL042005 AL042006 AA911481
	323898	243407_1	AA347566 AA346521 AI111169
	324048	267284_1	AA378739 AW964174 AA570564 AI076833 AW265063 AW006805 AA480656 AW004789
	324231	975669_1	W60827 AL079968 AL047234
25	324430	312113_1	AA464018 AA464079 AA468142
	324432	312487_1	AA464510 AA631257 AI740516 AI739132 AW972467 AI741376 AW068935 AI467852 AI752240 AI123717 AI754551 AW205510 AW044211 AW028889 AW198033 AI538632 AA513096
	324456	1155396_1	AW500954 AW501111 AW501394
	324512	1156071_1	AW502122 AW502125 AW501663 AW501720
30	324575	65704_1	AW502257 AI014241 AA100360 BE298534
	324609	333046_1	AW299534 AW299896 AA504765 AA505099 AA505100 AA584753 AW136415 AA768306
	324620	69834_1	BE397649 H14413 BE397689 BE514098 H53372 AA448021 R57944 AI307272 BE259369 H72331 BE251092 T27364 AA001666 AA044433 AA875998 AW075405 AW338356 AA001667 AW300173 AW514944 AW468914 AA604673 AA702749 AA805550 AA447621 AA934104 AI373527 AA604794 AI911203 AI500644 AI291383 AA731133 BE350633 AA044604 H95689 H14366 AW660983 AA912893 AI369587 AI382271 AA917508 AW138391 BE622560
35	324662	560496_1	AI376331 AI819150 AI097038 AI351100 AW504689
	324670	72231_1	AW503713 AA352950 AA044972 BE618246 AA335047 AW962269
	324692	351987_1	AA557952 AA677593 AA618150
	324715	290035_2	AI739168 AA426249 AI199636 AW505198 AW977291 AA824583 AA883419 AA724079 AI015524 AI377728 AW293682 AI928140 AA731438 AI092404 AI085630 AA731340
40	324728	210991_2	T85872 T48305
	324783	389615_1	AA640770 AI683112 AA913009
	324848	371388_1	AA602539 D59262 AI684171 N46711 AW021857 D19768
	324961	376239_1	AA613792 AW182329 T05304 AW858385
45	324988	22162_1	AK001379 AK001411 AW795711 T06997 AA287540 AA354538 AW957773 AI632268 AI651003 AI689650 AI809332 AW304483 AI805269 AA278506 AA862381 AA287875 AW628545 AI085761 AW025965 AI658615 AW628879 AW139496 AI214278 AA902745 AA991679 BE540102 AW593658 AI745602 AA744687 AI285441 AA807089 AI218314 AA721449 AI202987 AA432129 AI285502 AI281462 AA731319 BE082573
	325071	1562044_1	H09693 H09699 T09229
	325176	700767_1	T19142 AI351168 T52843 BE241963

**TABLE 1B**

Table 1B shows the genomic positioning for those primekeys in Table 1 that lack unigene ID's and accession numbers. For each predicted exon, the genomic sequence source used for prediction is listed. Nucleotide locations of each predicted exon are also listed.

Pkey: Unique number corresponding to an Eos probeset  
 Ref: Sequence source. The 7 digit numbers in this column are Genbank Identifier (GI) numbers. "Dunham I. et al." refers to the publication entitled "The DNA sequence of human chromosome 22," Dunham I. et al., Nature (1999) 402:489-495.  
 Strand: Indicates DNA strand from which exons were predicted.  
 Nt\_position: Indicates nucleotide positions of predicted exons.

	<b>15</b>	<b>Pkey</b>	<b>Ref</b>	<b>Strand</b>	<b>Nt_position</b>
	332792	Dunham, I. et.al.	Plus	73381-73768	
10	332908	Dunham, I. et.al.	Plus	1934283-1934366	
	332909	Dunham, I. et.al.	Plus	1946582-1946735	
20	332913	Dunham, I. et.al.	Plus	1963539-1963843	
	332952	Dunham, I. et.al.	Plus	2472864-2473012	
	332958	Dunham, I. et.al.	Plus	2516164-2516310	
25	332961	Dunham, I. et.al.	Plus	2521424-2521555	
	332975	Dunham, I. et.al.	Plus	2599641-2599702	
	332991	Dunham, I. et.al.	Plus	2686938-2687372	
30	333119	Dunham, I. et.al.	Plus	3288316-3288640	
	333131	Dunham, I. et.al.	Plus	3350064-3350170	
	333139	Dunham, I. et.al.	Plus	3369495-3369571	
35	333156	Dunham, I. et.al.	Plus	3617584-3617790	
	333222	Dunham, I. et.al.	Plus	3979706-3979803	
	333254	Dunham, I. et.al.	Plus	2521424-2521555	
40	333348	Dunham, I. et.al.	Plus	4711908-4712181	
	333349	Dunham, I. et.al.	Plus	4713940-4714084	
	333366	Dunham, I. et.al.	Plus	4798273-4798469	
45	333384	Dunham, I. et.al.	Plus	4907535-4907610	
	333385	Dunham, I. et.al.	Plus	4907928-4908032	
	333391	Dunham, I. et.al.	Plus	4916697-4916780	
	333488	Dunham, I. et.al.	Plus	5396233-5396310	
50	333520	Dunham, I. et.al.	Plus	5586133-5586296	
	333524	Dunham, I. et.al.	Plus	5612620-5612780	
	333532	Dunham, I. et.al.	Plus	5622804-5622937	
	333580	Dunham, I. et.al.	Plus	6142935-6143145	
	333585	Dunham, I. et.al.	Plus	6234778-6234894	
55	333597	Dunham, I. et.al.	Plus	6331421-6331536	
	333619	Dunham, I. et.al.	Plus	6562799-6562926	
	333671	Dunham, I. et.al.	Plus	7038849-7039193	
	333680	Dunham, I. et.al.	Plus	7071730-7071794	
	333682	Dunham, I. et.al.	Plus	7076641-7076760	
60	333763	Dunham, I. et.al.	Plus	7692491-7692630	
	333764	Dunham, I. et.al.	Plus	7693573-7693716	
	333769	Dunham, I. et.al.	Plus	7696625-7696707	
	333770	Dunham, I. et.al.	Plus	7700384-7700476	
	333849	Dunham, I. et.al.	Plus	8018323-8018472	
65	333875	Dunham, I. et.al.	Plus	8135505-8136179	
	333882	Dunham, I. et.al.	Plus	8153002-8153169	
	333922	Dunham, I. et.al.	Plus	8381385-8381444	
	333928	Dunham, I. et.al.	Plus	8468844-8469015	
	333947	Dunham, I. et.al.	Plus	8579888-8579966	
70	333949	Dunham, I. et.al.	Plus	8589634-8589791	
	333968	Dunham, I. et.al.	Plus	8681004-8681241	
	333983	Dunham, I. et.al.	Plus	8813593-8813668	
	333995	Dunham, I. et.al.	Plus	8855296-8855424	
	333997	Dunham, I. et.al.	Plus	8866668-8867255	
	334003	Dunham, I. et.al.	Plus	8892882-8892970	
	334012	Dunham, I. et.al.	Plus	9007456-9010221	
	334047	Dunham, I. et.al.	Plus	9428152-9428211	
	334063	Dunham, I. et.al.	Plus	9731991-9732085	
	334066	Dunham, I. et.al.	Plus	9739568-9739680	
	334078	Dunham, I. et.al.	Plus	9809783-9809863	
70	334118	Dunham, I. et.al.	Plus	10344273-10344384	

	334122	Dunham, I. et.al.	Plus	10411792-10411901
	334150	Dunham, I. et.al.	Plus	10529221-10529854
5	334220	Dunham, I. et.al.	Plus	12718720-12718857
	334298	Dunham, I. et.al.	Plus	13424763-13425914
	334324	Dunham, I. et.al.	Plus	13539210-13539323
	334335	Dunham, I. et.al.	Plus	13608488-13608705
	334433	Dunham, I. et.al.	Plus	14273261-14273429
	334532	Dunham, I. et.al.	Plus	14792798-14792901
10	334561	Dunham, I. et.al.	Plus	14987299-14987447
	334616	Dunham, I. et.al.	Plus	15176123-15176470
	334628	Dunham, I. et.al.	Plus	15310346-15310415
	334630	Dunham, I. et.al.	Plus	15322614-15322744
	334631	Dunham, I. et.al.	Plus	15325949-15326116
15	334661	Dunham, I. et.al.	Plus	15477716-15477786
	334677	Dunham, I. et.al.	Plus	15517449-15517560
	334696	Dunham, I. et.al.	Plus	15665919-15666002
	334714	Dunham, I. et.al.	Plus	15760702-15760767
	334718	Dunham, I. et.al.	Plus	15775491-15775599
20	334720	Dunham, I. et.al.	Plus	15792931-15793085
	334727	Dunham, I. et.al.	Plus	15942616-15942750
	334739	Dunham, I. et.al.	Plus	16004120-16004225
	334740	Dunham, I. et.al.	Plus	16009324-16009547
	334769	Dunham, I. et.al.	Plus	16170704-16170876
25	334872	Dunham, I. et.al.	Plus	19162417-19162565
	334876	Dunham, I. et.al.	Plus	19185336-19185400
	334883	Dunham, I. et.al.	Plus	19223107-19223253
	334891	Dunham, I. et.al.	Plus	19299770-19299944
	334900	Dunham, I. et.al.	Plus	19315678-19315743
30	334902	Dunham, I. et.al.	Plus	19317083-19317195
	334914	Dunham, I. et.al.	Plus	19495158-19495275
	334916	Dunham, I. et.al.	Plus	19572924-19573846
	335044	Dunham, I. et.al.	Plus	20842088-20842682
	335081	Dunham, I. et.al.	Plus	21113871-21113937
35	335158	Dunham, I. et.al.	Plus	21569610-21569666
	335164	Dunham, I. et.al.	Plus	21585912-21586014
	335166	Dunham, I. et.al.	Plus	21587100-21587213
	335170	Dunham, I. et.al.	Plus	21623383-21623967
	335188	Dunham, I. et.al.	Plus	21669118-21669328
40	335189	Dunham, I. et.al.	Plus	21673403-21673472
	335200	Dunham, I. et.al.	Plus	21743499-21743881
	335211	Dunham, I. et.al.	Plus	21774611-21774680
	335219	Dunham, I. et.al.	Plus	21875591-21875688
	335221	Dunham, I. et.al.	Plus	21882840-21882968
45	335225	Dunham, I. et.al.	Plus	21890315-21890448
	335255	Dunham, I. et.al.	Plus	22032258-22032661
	335287	Dunham, I. et.al.	Plus	22299047-22299299
	335361	Dunham, I. et.al.	Plus	22807292-22807445
	335364	Dunham, I. et.al.	Plus	22833430-22833586
50	335369	Dunham, I. et.al.	Plus	22843392-22843506
	335468	Dunham, I. et.al.	Plus	23787245-23787367
	335481	Dunham, I. et.al.	Plus	24082522-24084870
	335488	Dunham, I. et.al.	Plus	24118744-24118839
	335496	Dunham, I. et.al.	Plus	24164386-24164545
55	335497	Dunham, I. et.al.	Plus	24167666-24167869
	335499	Dunham, I. et.al.	Plus	24176698-24176869
	335504	Dunham, I. et.al.	Plus	24182110-24182199
	335599	Dunham, I. et.al.	Plus	25043628-25043775
	335623	Dunham, I. et.al.	Plus	25138489-25138547
60	335653	Dunham, I. et.al.	Plus	25329710-25329802
	335687	Dunham, I. et.al.	Plus	25445952-25446064
	335690	Dunham, I. et.al.	Plus	25455442-25455625
	335692	Dunham, I. et.al.	Plus	25468557-25468725
	335697	Dunham, I. et.al.	Plus	25481456-25481649
65	335701	Dunham, I. et.al.	Plus	25513366-25513807
	335708	Dunham, I. et.al.	Plus	25541777-25541907
	335739	Dunham, I. et.al.	Plus	25698550-25698826
	335742	Dunham, I. et.al.	Plus	25712654-25712771
	336003	Dunham, I. et.al.	Plus	28406289-28406759
	336015	Dunham, I. et.al.	Plus	28640586-28640673
70	336016	Dunham, I. et.al.	Plus	28646816-28646947
	336018	Dunham, I. et.al.	Plus	28660880-28660978
	336019	Dunham, I. et.al.	Plus	28663992-28664102
	336020	Dunham, I. et.al.	Plus	28683778-28683851
	336021	Dunham, I. et.al.	Plus	28686482-28686559
75	336023	Dunham, I. et.al.	Plus	28698240-28698343

	336071	Dunham, I. et.al.	Plus	29264457-29264684
	336090	Dunham, I. et.al.	Plus	29413020-29413162
	336107	Dunham, I. et.al.	Plus	29987731-29987869
	336121	Dunham, I. et.al.	Plus	30048054-30048129
5	336124	Dunham, I. et.al.	Plus	30053441-30053500
	336132	Dunham, I. et.al.	Plus	30107247-30107412
	336135	Dunham, I. et.al.	Plus	30123235-30123335
	336194	Dunham, I. et.al.	Plus	30443138-30443282
	336235	Dunham, I. et.al.	Plus	31122315-31122623
10	336367	Dunham, I. et.al.	Plus	33942937-33943058
	336379	Dunham, I. et.al.	Plus	33995071-33995243
	336439	Dunham, I. et.al.	Plus	34186130-34186215
	336502	Dunham, I. et.al.	Plus	34268953-34269083
15	336572	Dunham, I. et.al.	Plus	34446383-34446496
	336602	Dunham, I. et.al.	Plus	13424060-13424582
	336721	Dunham, I. et.al.	Plus	3371522-3371586
	336739	Dunham, I. et.al.	Plus	2599641-2599702
	336766	Dunham, I. et.al.	Plus	4905608-4905684
20	336833	Dunham, I. et.al.	Plus	6856506-6856634
	336836	Dunham, I. et.al.	Plus	7077262-7077326
	336878	Dunham, I. et.al.	Plus	9200300-9200399
	336880	Dunham, I. et.al.	Plus	9250034-9250123
	336902	Dunham, I. et.al.	Plus	10385555-10386053
25	336917	Dunham, I. et.al.	Plus	11228329-11228403
	336919	Dunham, I. et.al.	Plus	11351181-11351274
	336924	Dunham, I. et.al.	Plus	11525273-11525527
	336946	Dunham, I. et.al.	Plus	12337073-12337258
	336953	Dunham, I. et.al.	Plus	12988791-12988889
30	336979	Dunham, I. et.al.	Plus	14270748-14270816
	337169	Dunham, I. et.al.	Plus	23529987-23530214
	337175	Dunham, I. et.al.	Plus	23782209-23782374
	337182	Dunham, I. et.al.	Plus	23934889-23934962
	337238	Dunham, I. et.al.	Plus	27141465-27141776
	337303	Dunham, I. et.al.	Plus	29128849-29128974
35	337489	Dunham, I. et.al.	Plus	33295724-33295872
	337503	Dunham, I. et.al.	Plus	33385583-33385857
	337504	Dunham, I. et.al.	Plus	33386053-33386236
	337570	Dunham, I. et.al.	Plus	359309-359459
	337585	Dunham, I. et.al.	Plus	951744-952008
40	337629	Dunham, I. et.al.	Plus	2017380-2017517
	337670	Dunham, I. et.al.	Plus	3110593-3110760
	337674	Dunham, I. et.al.	Plus	3332616-3332697
	337692	Dunham, I. et.al.	Plus	3575105-3575299
	337740	Dunham, I. et.al.	Plus	3870165-3870223
45	337755	Dunham, I. et.al.	Plus	3971764-3971900
	337807	Dunham, I. et.al.	Plus	4444885-4444981
	337844	Dunham, I. et.al.	Plus	4993372-4993603
	337902	Dunham, I. et.al.	Plus	5682218-5682307
50	337904	Dunham, I. et.al.	Plus	5685819-5686012
	337919	Dunham, I. et.al.	Plus	6035207-6035326
	337951	Dunham, I. et.al.	Plus	6766321-6766382
	337958	Dunham, I. et.al.	Plus	6969162-6969270
	337964	Dunham, I. et.al.	Plus	7032720-7032802
55	338008	Dunham, I. et.al.	Plus	7697068-7697236
	338053	Dunham, I. et.al.	Plus	8412742-8412823
	338057	Dunham, I. et.al.	Plus	8526397-8526522
	338059	Dunham, I. et.al.	Plus	8540638-8540712
	338120	Dunham, I. et.al.	Plus	10765673-10765820
60	338124	Dunham, I. et.al.	Plus	10860311-10860471
	338178	Dunham, I. et.al.	Plus	12800037-12800181
	338196	Dunham, I. et.al.	Plus	13629317-13629466
	338204	Dunham, I. et.al.	Plus	13870980-13871152
	338239	Dunham, I. et.al.	Plus	14669918-14670016
	338249	Dunham, I. et.al.	Plus	14870864-14870944
65	338250	Dunham, I. et.al.	Plus	14874504-14874575
	338251	Dunham, I. et.al.	Plus	14963460-14963521
	338260	Dunham, I. et.al.	Plus	15458919-15459257
	338282	Dunham, I. et.al.	Plus	16240812-16241002
	338316	Dunham, I. et.al.	Plus	17089711-17089988
70	338364	Dunham, I. et.al.	Plus	18210049-18210226
	338374	Dunham, I. et.al.	Plus	18371200-18371282
	338454	Dunham, I. et.al.	Plus	20180035-20180113
	338494	Dunham, I. et.al.	Plus	21181818-21182009
	338596	Dunham, I. et.al.	Plus	23078273-23078348
75	338622	Dunham, I. et.al.	Plus	23546552-23546749

	338702	Dunham, I. et.al.	Plus	25219632-25219739
	338707	Dunham, I. et.al.	Plus	25266346-25266417
5	338716	Dunham, I. et.al.	Plus	25472519-25472686
	338765	Dunham, I. et.al.	Plus	26657278-26657346
	338852	Dunham, I. et.al.	Plus	28086911-28086971
	338862	Dunham, I. et.al.	Plus	28230332-28230444
	338962	Dunham, I. et.al.	Plus	29581892-29582020
	338997	Dunham, I. et.al.	Plus	30092658-30092730
10	339164	Dunham, I. et.al.	Plus	32207441-32207802
	339305	Dunham, I. et.al.	Plus	33334676-33334864
	339313	Dunham, I. et.al.	Plus	33383457-33383585
	339319	Dunham, I. et.al.	Plus	33410900-33410972
	339323	Dunham, I. et.al.	Plus	33418663-33418829
15	339356	Dunham, I. et.al.	Plus	33573387-33573517
	339358	Dunham, I. et.al.	Plus	33577760-33577922
	339361	Dunham, I. et.al.	Plus	33580121-33580251
	339413	Dunham, I. et.al.	Plus	34268734-34268875
	339418	Dunham, I. et.al.	Plus	34353362-34353421
20	339436	Dunham, I. et.al.	Plus	34546469-34546834
	332813	Dunham, I. et.al.	Minus	318840-318777
	332854	Dunham, I. et.al.	Minus	1283611-1283053
	332858	Dunham, I. et.al.	Minus	1339607-1339397
	332863	Dunham, I. et.al.	Minus	1389980-1389884
25	332868	Dunham, I. et.al.	Minus	1413234-1413078
	332884	Dunham, I. et.al.	Minus	1573063-1572923
	332886	Dunham, I. et.al.	Minus	1574863-1574660
	332896	Dunham, I. et.al.	Minus	1631641-1631422
	332929	Dunham, I. et.al.	Minus	2020758-2020664
30	332930	Dunham, I. et.al.	Minus	2022565-2022497
	332931	Dunham, I. et.al.	Minus	2023651-2023562
	332932	Dunham, I. et.al.	Minus	2035348-2035282
	332965	Dunham, I. et.al.	Minus	2537457-2537396
	332995	Dunham, I. et.al.	Minus	2708847-2708685
35	333002	Dunham, I. et.al.	Minus	2537457-2537396
	333011	Dunham, I. et.al.	Minus	2769669-2769571
	333029	Dunham, I. et.al.	Minus	2885241-2885175
	333033	Dunham, I. et.al.	Minus	2889900-2889699
	333110	Dunham, I. et.al.	Minus	3244892-3244779
40	333126	Dunham, I. et.al.	Minus	3324305-3324184
	333217	Dunham, I. et.al.	Minus	3967830-3967716
	333220	Dunham, I. et.al.	Minus	3969363-3968789
	333242	Dunham, I. et.al.	Minus	4104544-4104259
	333243	Dunham, I. et.al.	Minus	4104961-4104728
	333259	Dunham, I. et.al.	Minus	4306769-4306639
45	333270	Dunham, I. et.al.	Minus	4373573-4373219
	333278	Dunham, I. et.al.	Minus	4414616-4414389
	333279	Dunham, I. et.al.	Minus	4415252-4414844
	333358	Dunham, I. et.al.	Minus	4732336-4732236
50	333408	Dunham, I. et.al.	Minus	4936879-4936661
	333441	Dunham, I. et.al.	Minus	2708847-2708685
	333444	Dunham, I. et.al.	Minus	5070077-5069643
	333447	Dunham, I. et.al.	Minus	2537457-2537396
	333466	Dunham, I. et.al.	Minus	2708847-2708685
55	333473	Dunham, I. et.al.	Minus	2537457-2537396
	333496	Dunham, I. et.al.	Minus	5404643-5404523
	333511	Dunham, I. et.al.	Minus	5557881-5557718
	333542	Dunham, I. et.al.	Minus	5861529-5861341
	333568	Dunham, I. et.al.	Minus	5965072-5964999
60	333582	Dunham, I. et.al.	Minus	6158522-6158322
	333665	Dunham, I. et.al.	Minus	6975471-6975215
	333673	Dunham, I. et.al.	Minus	7054704-7054602
	333734	Dunham, I. et.al.	Minus	7535394-7535309
	333964	Dunham, I. et.al.	Minus	8626045-8625966
65	334156	Dunham, I. et.al.	Minus	10580883-10580765
	334172	Dunham, I. et.al.	Minus	11644142-11644008
	334183	Dunham, I. et.al.	Minus	11832582-11832508
	334184	Dunham, I. et.al.	Minus	11833848-11833757
	334223	Dunham, I. et.al.	Minus	12734365-12734269
70	334270	Dunham, I. et.al.	Minus	13249131-13249007
	334288	Dunham, I. et.al.	Minus	13295104-13294969
	334303	Dunham, I. et.al.	Minus	13454331-13454217
	334358	Dunham, I. et.al.	Minus	13724372-13724201
	334370	Dunham, I. et.al.	Minus	13782655-13782493
	334472	Dunham, I. et.al.	Minus	14391308-14391169
75	334474	Dunham, I. et.al.	Minus	14391920-14391809

	334487	Dunham, I. et.al.	Minus	14432191-14432132
	334500	Dunham, I. et.al.	Minus	14486730-14486621
	334537	Dunham, I. et.al.	Minus	14827542-14827354
5	334621	Dunham, I. et.al.	Minus	15190418-15190299
	334648	Dunham, I. et.al.	Minus	15363301-15363222
	334764	Dunham, I. et.al.	Minus	16151208-16151104
	334783	Dunham, I. et.al.	Minus	16293336-16293226
	334784	Dunham, I. et.al.	Minus	16294548-16294360
	334786	Dunham, I. et.al.	Minus	16297434-16297275
10	334789	Dunham, I. et.al.	Minus	16306095-16305996
	334806	Dunham, I. et.al.	Minus	16433227-16433125
	334823	Dunham, I. et.al.	Minus	16851360-16851189
	334850	Dunham, I. et.al.	Minus	17660892-17660787
	334924	Dunham, I. et.al.	Minus	19744615-19744229
15	334939	Dunham, I. et.al.	Minus	20131162-20131054
	334943	Dunham, I. et.al.	Minus	20135064-20134903
	334948	Dunham, I. et.al.	Minus	20141727-20141583
	334970	Dunham, I. et.al.	Minus	20195886-20195554
	334991	Dunham, I. et.al.	Minus	20341858-20341773
20	334993	Dunham, I. et.al.	Minus	20354277-20354174
	335062	Dunham, I. et.al.	Minus	20921289-20921087
	335207	Dunham, I. et.al.	Minus	21763011-21762880
	335288	Dunham, I. et.al.	Minus	22304275-22303770
	335289	Dunham, I. et.al.	Minus	22305950-22305708
25	335293	Dunham, I. et.al.	Minus	22316408-22316275
	335332	Dunham, I. et.al.	Minus	22557778-22557557
	335389	Dunham, I. et.al.	Minus	23043682-23043558
	335478	Dunham, I. et.al.	Minus	23924778-23924329
	335524	Dunham, I. et.al.	Minus	24237218-24236208
30	335547	Dunham, I. et.al.	Minus	24658526-24658460
	335671	Dunham, I. et.al.	Minus	25358629-25358533
	335682	Dunham, I. et.al.	Minus	25421215-25421093
	335684	Dunham, I. et.al.	Minus	25425165-25425096
	335698	Dunham, I. et.al.	Minus	25493029-25492767
35	335755	Dunham, I. et.al.	Minus	25763806-25763747
	335756	Dunham, I. et.al.	Minus	25764330-25764251
	335773	Dunham, I. et.al.	Minus	25880858-25880661
	335813	Dunham, I. et.al.	Minus	26318734-26318649
	335815	Dunham, I. et.al.	Minus	26320518-26320421
40	335817	Dunham, I. et.al.	Minus	26321875-26321750
	335827	Dunham, I. et.al.	Minus	26380557-26380472
	335829	Dunham, I. et.al.	Minus	26382348-26382251
	335836	Dunham, I. et.al.	Minus	26397823-26397694
	335857	Dunham, I. et.al.	Minus	26677208-26677096
45	335860	Dunham, I. et.al.	Minus	26684908-26684800
	335871	Dunham, I. et.al.	Minus	26734972-26734892
	335896	Dunham, I. et.al.	Minus	26977639-26977558
	335897	Dunham, I. et.al.	Minus	26978293-26978142
	335903	Dunham, I. et.al.	Minus	26985739-26985580
50	335914	Dunham, I. et.al.	Minus	27024197-27023994
	335916	Dunham, I. et.al.	Minus	27027028-27026912
	335935	Dunham, I. et.al.	Minus	27360288-27360058
	335940	Dunham, I. et.al.	Minus	27420194-27420000
	335999	Dunham, I. et.al.	Minus	28033986-28033848
55	336045	Dunham, I. et.al.	Minus	29044217-29044140
	336140	Dunham, I. et.al.	Minus	30134204-30133980
	336182	Dunham, I. et.al.	Minus	30371411-30371339
	336198	Dunham, I. et.al.	Minus	30459668-30459460
	336227	Dunham, I. et.al.	Minus	30902014-30901946
60	336246	Dunham, I. et.al.	Minus	31425669-31425253
	336262	Dunham, I. et.al.	Minus	31833610-31833533
	336292	Dunham, I. et.al.	Minus	32818035-32817927
	336347	Dunham, I. et.al.	Minus	33843218-33843104
	336397	Dunham, I. et.al.	Minus	34021504-34021389
65	336434	Dunham, I. et.al.	Minus	34073056-34072952
	336662	Dunham, I. et.al.	Minus	2158060-2157993
	336675	Dunham, I. et.al.	Minus	2020758-2020664
	336676	Dunham, I. et.al.	Minus	2022565-2022497
	336677	Dunham, I. et.al.	Minus	2023651-2023562
70	336678	Dunham, I. et.al.	Minus	2035348-2035282
	336684	Dunham, I. et.al.	Minus	2158060-2157993
	336703	Dunham, I. et.al.	Minus	5071373-5071278
	336707	Dunham, I. et.al.	Minus	2820219-2820111
	336722	Dunham, I. et.al.	Minus	3377722-3377590
75	336778	Dunham, I. et.al.	Minus	5071373-5071278

	336846	Dunham, I. et.al.	Minus	7566306-7566238
	336977	Dunham, I. et.al.	Minus	14110003-14109910
	336981	Dunham, I. et.al.	Minus	14478638-14478472
5	337056	Dunham, I. et.al.	Minus	17975104-17974976
	337094	Dunham, I. et.al.	Minus	20146915-20146778
	337102	Dunham, I. et.al.	Minus	20581738-20581628
	337161	Dunham, I. et.al.	Minus	23473450-23473375
	337366	Dunham, I. et.al.	Minus	30961904-30961787
10	337422	Dunham, I. et.al.	Minus	32030671-32030417
	337452	Dunham, I. et.al.	Minus	32415187-32415117
	337534	Dunham, I. et.al.	Minus	34193847-34193769
	337595	Dunham, I. et.al.	Minus	1020506-1020210
	337602	Dunham, I. et.al.	Minus	1282987-1282741
15	337603	Dunham, I. et.al.	Minus	1299296-1299194
	337645	Dunham, I. et.al.	Minus	5141462-5141329
	337760	Dunham, I. et.al.	Minus	4008389-4008037
	337772	Dunham, I. et.al.	Minus	4061918-4061782
	337776	Dunham, I. et.al.	Minus	4084555-4084460
20	337840	Dunham, I. et.al.	Minus	4940540-4940409
	337908	Dunham, I. et.al.	Minus	5697187-5697071
	337937	Dunham, I. et.al.	Minus	6556005-6555907
	337974	Dunham, I. et.al.	Minus	7153401-7153085
	337986	Dunham, I. et.al.	Minus	7296008-7295951
25	338000	Dunham, I. et.al.	Minus	7530875-7530793
	338221	Dunham, I. et.al.	Minus	14183649-14183568
	338510	Dunham, I. et.al.	Minus	21339584-21339508
	338535	Dunham, I. et.al.	Minus	21799696-21799274
	338546	Dunham, I. et.al.	Minus	22012448-22012383
30	338556	Dunham, I. et.al.	Minus	22179326-22179234
	338561	Dunham, I. et.al.	Minus	22311966-22311856
	338668	Dunham, I. et.al.	Minus	24500606-24500442
	338689	Dunham, I. et.al.	Minus	24893073-24892972
	338727	Dunham, I. et.al.	Minus	25926788-25926580
35	338759	Dunham, I. et.al.	Minus	26582475-26582199
	338763	Dunham, I. et.al.	Minus	26628148-26628009
	338779	Dunham, I. et.al.	Minus	27030151-27029795
	338876	Dunham, I. et.al.	Minus	28364326-28364071
	338889	Dunham, I. et.al.	Minus	28477552-28477412
40	339028	Dunham, I. et.al.	Minus	30574122-30573937
	339044	Dunham, I. et.al.	Minus	30721853-30721740
	339128	Dunham, I. et.al.	Minus	31692815-31692686
	339188	Dunham, I. et.al.	Minus	32347554-32347250
	339208	Dunham, I. et.al.	Minus	32491714-32491657
45	339215	Dunham, I. et.al.	Minus	32502559-32502383
	339230	Dunham, I. et.al.	Minus	32729004-32728929
	339256	Dunham, I. et.al.	Minus	32926055-32925967
	339280	Dunham, I. et.al.	Minus	33114230-33114010
	339370	Dunham, I. et.al.	Minus	33805912-33805797
	337895			
50	329598	3962482	Plus	39924-40220
	329563	3962490	Minus	410-635
	329557	3962492	Minus	53197-53647
	329539	3983503	Minus	1-326
	329526	3983506	Plus	12251-12325
55	329524	3983507	Minus	38025-38143
	329502	3983517	Plus	75-338
	329503	3983517	Minus	1801-1937
	329499	3983518	Plus	33463-33789
60	329479	3983526	Minus	7425-7561
	329625	4567169	Minus	85893-85984
	325363	5866920	Plus	700446-700516
	325366	5866920	Minus	920962-921713
	325433	5866936	Minus	480706-480826
65	325447	5866941	Minus	372480-372621
	325481	5866957	Plus	47590-47672
	325482	5866957	Plus	47957-48078
	325472	6017034	Minus	289581-289657
	325513	6017035	Minus	34295-34490
	325519	6017036	Minus	186804-186915
70	325587	6682462	Plus	126724-126967
	325585	6682462	Plus	73476-73574
	325594	5866992	Minus	470474-470566
	325609	5866996	Minus	981751-981849
	325622	5867000	Plus	69994-70075
75	325751	6682474	Plus	130437-130520

	325815	6682483	Minus	129273-130754
	329762	6048280	Plus	127744-127878
	329789	6469354	Minus	118977-119036
5	329797	6523160	Minus	10616-10894
	329838	6672062	Plus	33990-34098
	325864	5867069	Minus	110834-110904
	325885	5867087	Plus	193212-193377
	325892	5867088	Minus	10498-10652
10	325929	5867125	Minus	51715-51996
	325843	6552453	Minus	7126-7232
	329989	4567166	Plus	72861-73052
	329960	5091594	Minus	1031-1162
15	329959	5103803	Plus	188050-188193
	329936	6165200	Minus	82761-82920
	329919	6223624	Minus	103492-103681
	330004	6623963	Minus	78872-78999
	326025	5867176	Plus	70854-70915
	326054	5867184	Minus	146342-146469
20	326112	5867192	Plus	2151-2725
	326165	5867208	Minus	62787-62929
	326213	5867224	Minus	60751-60927
	326219	5867226	Minus	264008-264274
	326160	5867254	Minus	112000-112137
25	326257	5867264	Plus	222712-222819
	330057	6478962	Plus	75145-75287
	326359	5867293	Plus	9436-9494
	326393	5867341	Plus	41702-41841
	326399	5867353	Plus	6385-6536
30	326401	5867355	Plus	35165-35332
	326416	5867362	Minus	45283-45375
	326431	5867371	Plus	15855-15971
	326460	5867400	Minus	142633-142935
	326517	5867439	Plus	44732-46356
35	326519	5867439	Plus	166004-166243
	326596	6138928	Plus	133386-133563
	330081	6015314	Minus	5768-5835
	326714	5867595	Plus	124490-124568
	326752	5867615	Minus	1214-1562
40	326757	6249610	Plus	74531-74597
	326668	6552455	Plus	146726-146838
	326720	6552456	Plus	84525-84677
	326725	6552456	Minus	223005-223125
	326862	6552465	Plus	107702-107782
	326882	6682509	Minus	167988-168179
45	326892	6682511	Plus	119424-119500
	326996	5867660	Minus	63212-63404
	327010	5867664	Plus	941057-941139
	326919	6456782	Minus	40486-41046
50	327042	6531965	Minus	1380806-1381443
	327049	6531965	Minus	1924026-1924110
	327072	6531965	Minus	3796429-3797197
	327074	6531965	Plus	4039993-4040096
	327075	6531965	Plus	4041318-4041431
55	326981	6588016	Plus	105091-106038
	327133	6682522	Plus	38069-38938
	330137	4210430	Minus	21220-21377
	330138	4210430	Minus	22334-22460
	330143	4210430	Plus	184737-184848
60	330153	4325335	Plus	146951-147475
	330135	4456470	Minus	121583-121885
	327205	5867447	Plus	167335-167576
	327212	5867463	Minus	42308-42424
	327287	5867479	Minus	62838-63024
	327331	5867516	Minus	55606-55737
65	327364	6552412	Minus	115235-115396
	327413	5867750	Plus	101410-101508
	327481	5867783	Plus	104472-104673
	327458	6004455	Plus	173257-173378
	327516	6117815	Plus	199078-199216
70	327527	6381882	Minus	98950-99040
	327548	5867797	Minus	81067-81130
	327554	5867801	Minus	23092-23191
	327565	5867811	Plus	32516-32778
	327600	6004462	Minus	2621-2862
75	327687	5867847	Minus	169293-169362

	330182	5123954	Plus	120156-120245
	327742	5867944	Minus	143307-143512
	327805	5867968	Plus	19952-20019
	327809	5867968	Plus	54610-54761
5	327814	5867968	Plus	69377-70566
	327815	5867968	Plus	70804-71401
	327791	5867977	Plus	22491-22610
	327745	6531959	Minus	229066-229124
10	330211	6013592	Plus	59158-59215
	330207	6013606	Minus	109912-110004
	330257	6671881	Minus	143228-143393
	330262	6671884	Plus	67913-68053
	330286	6671913	Minus	31050-31171
15	328105	5868020	Minus	301705-301784
	328113	5868024	Minus	80378-80491
	328142	5868050	Minus	9656-9778
	328152	5868060	Minus	73981-74203
	328170	5868071	Plus	93170-93295
20	327910	5868162	Plus	21622-21748
	327919	5868165	Plus	547701-547800
	327990	5868218	Minus	36225-36503
	328249	6381891	Minus	96352-96527
	328251	6381891	Plus	124444-124557
	328253	6381894	Minus	4411-4509
25	328084	6469819	Minus	155366-155459
	328274	5868219	Minus	31244-31439
	328615	5868239	Plus	35214-35347
	328632	5868247	Plus	76734-76853
30	328779	5868309	Plus	41570-41639
	328783	5868309	Minus	73658-73822
	328801	5868321	Minus	44492-44609
	328820	5868330	Plus	90446-90602
	328835	5868339	Plus	88053-88461
35	328290	5868363	Minus	127366-127496
	328321	5868373	Minus	1029614-1029673
	328332	5868375	Plus	280154-280289
	328333	5868375	Plus	282506-282664
	328349	5868383	Minus	260704-260804
40	328450	5868425	Minus	209192-209321
	328466	5868434	Minus	15643-15900
	328479	5868449	Minus	331-560
	328481	5868449	Minus	8987-9180
	328546	5868487	Minus	17547-17722
45	328662	6004473	Plus	1184773-1184855
	328767	6017031	Minus	35625-35723
	328768	6017031	Minus	223741-224238
	328857	6381927	Minus	80557-81051
	328878	6552423	Plus	105580-105774
50	328882	6552423	Minus	157669-157826
	328690	6588001	Minus	571207-571274
	328691	6588001	Minus	579598-579664
	330307	4877982	Plus	107384-107559
	328903	5868514	Plus	23625-24468
55	328987	5868535	Minus	25705-25764
	328998	5868538	Plus	40996-41104
	329062	5868590	Minus	58977-59094
	329086	5868604	Minus	35489-35588
	329154	5868686	Minus	200851-201356
60	329156	5868686	Minus	202013-202341
	329164	5868691	Plus	62305-62517
	329170	5868693	Plus	67924-68019
	329179	5868704	Plus	181639-181815
	329193	5868716	Plus	168095-168181
65	329254	5868733	Plus	4133-4214
	329369	5868842	Minus	121148-121516
	329367	5868842	Minus	87201-87587
	329141	6017060	Plus	343924-343997
	329347	6456785	Plus	18433-18897
70	329017	6682532	Minus	255591-255672
	329434	5868883	Minus	31124-31263

**TABLE 2 DNA AND PROTEIN SEQUENCES FOR CBF9 AND BFO8**

Table 2 provides the nucleic acid and protein sequence of the CBF9 and BFO8 genes as well as the Unigene and Exemplar accession numbers for CBF9 and BFO8.

5

**CBF9 DNA SEQUENCE**

Gene name:	ESTs
Unigene number:	Hs.157601
Probeset Accession #:	W07459
Nucleic Acid Accession #:	AC005383
Coding Sequence:	328-2751 (underlined sequences correspond to start and stop codons)

	1	11	21	31	41	51	
	GACAGTGTTC	GCGGCTGCAC	CGCTCGGAGG	CTGGGTGACC	CGCGTAGAAC	TGAAGTACTT	60
	TTTTATTTGC	AGACCTGGGC	CGATGCCGCT	TTAAAAAAACG	CGAGGGGCTC	TATGCACCTC	120
	CCTGGCGGTA	GTTCCCTCGA	CCTCAGCCGG	GTGGGGTCTG	GCCGCCCTCT	CCCAGGAGAG	180
10	ACAAACAGGT	GTCCCCACGTG	GCAGCCGCCG	CCCGGGCGCC	CCTCCTGTGA	TCCCCTAGCG	240
	CCCCCTGGCC	CGAGCCGCGC	CGGGGTCTGT	GAGTAGAGCC	GCCCAGGGCAC	CGAGCGCTGG	300
	TCGCCGCTCT	CCTTCGCTTA	TATCAACATG	CCCCCTTCTC	TGTTGCTGGA	GGCCGTCTGT	360
	GTTTTCCTGT	TTTCCAGAGT	GCCCCCATCT	CTCCCTCTCC	AGGAAGTCCA	TGTAAGCAAA	420
15	GAAACCATCG	GGAAAGATTT	AGCTGCCAGC	AAAATGATGT	GGTGTCTGGC	TGCAGTGGAC	480
	ATCATGTTTC	TGTTAGATGG	GTCTAACAGC	GTGGGAAAG	GGAGCTTTGA	AAGGTCCAAG	540
	CACTTGCAC	TCACAGTC	TGACGGTCTG	GACATCAGCC	CCGAGAGGGT	CAGAGTGGGA	600
20	GCATTCCAGT	TCAGTCCAC	TCCTCATCTG	GAATTCCCC	TGGATTCTATT	TTCAACCCAA	660
	CAGGAAGTGA	AGGCAGAACAT	CAAGAGGATG	GTTTTCAAAG	GAGGGCCGAC	GGAGACGGAA	720
	CTTGCTCTGA	AATAACCTTCT	GCACAGAGGG	TTGCTGGAG	GCAGAAATGC	TTCTGTGCC	780
25	CAGATCCTCA	TCATCGTCAC	TGATGGGAAG	TCCCAGGGGG	ATGTGGCACT	GCCATCCAAG	840
	CAGCTGAAGG	AAAGGGGTGT	CACTGTGTTT	GCTGTGGGGG	TCAGGTTTC	CAGGTGGGAG	900
	GAGCTGCATG	CACTGGCCAG	CGAGCCTAGA	GGGCAGCACG	TGCTGTTGGC	TGAGCAGGTG	960
30	GAGGATGCCA	CCAACGGCCT	CTTCAGCACC	CTCAGCAGCT	CGGCCATCTG	CTCCAGCGCC	1020
	ACGCCAGACT	GCAGGGTCGA	GGCTCACCCC	TGTGAGCACA	GGACGCTGGA	GATGGTCCGG	1080
	GAGTCGCTG	GCAATGCC	ATGCTGGAGA	GGATCGCGC	GGACCCCTTG	GGTGTGGCT	1140
35	GCACACTGTC	CCTTCTACAG	CTGGAAGAGA	GTGTTCTAA	CCCCACCTGC	CACCTGCTAC	1200
	AGGACCACCT	GCCCAGGCC	CTGTGACTCG	CAGCCCTGCC	AGAATGGAGG	CACATGTGTT	1260
	CCAGAAGGAC	TGGACGGCTA	CCAGTGCCTC	TGCCCCTG	CCTTGGAGG	GGAGGCTAAC	1320
40	TGTGCCCCG	AGCTGAGCCT	GGATGCGAGG	GTGCACTC	TCTTCCTGCT	GGACAGCTCT	1380
	GCGGGCACCA	CTCTGGACGG	CTTCCTGCGG	GCCAAAGTCT	TCGTGAAGCG	TGTTGTGCGG	1440
	GCCGTGCTG	GGCAGGACTC	TCGGGGCCGA	GTGGGTGTGG	CCACATACAG	CAGGGAGCTG	1500
	CTGGTGGCGG	TGCGCTGTGG	GGAGTACCAAG	GATGTGCCTG	ACCTGGTCTG	GAGCCTCGAT	1560
	GGCATTCCCT	TCCGTGGTGG	CCCCACCC	ACGGGCAGTG	CCTTGC	GGCGGAGAG	1620
45	CGTGGCTTCG	GGAGCGCCAC	CAGGACAGGC	CAGGACCGGC	CACGTAGAGT	GGTGGTTTG	1680
	CTCACTGAGT	CACACTCCG	GGATGAGGTT	CGGGGCCAG	CGCGTCACGC	AAGGGCGCGA	1740
	GAGCTGCTCC	TGCTGGGTGT	AGGCAGTGAG	GCCGTGCGGG	CAGAGCTGGA	GGAGATCACA	1800
	GGCAGCCCAA	AGCATGTGAT	GGTCTACTCG	GATCCTCAGG	ATCTGTTCAA	CCAAATCCCT	1860
	GAGCTGCAGG	GGAAAGCTGTG	CAGCCGGCAG	CGGGCAGGGT	CCCGGACACA	AGCCCTGGAC	1920
50	CTCGTCTTC	TGTTGGACAC	CTCTGCCTCA	GTAGGGCCG	AGAATTGTC	TCAGATGCAG	1980
	AGCTTTGTA	GAAGCTGTGC	CCTCTGAGTT	GGGTGAACC	CTGACGTC	ACAGGTGGC	2040
	CTGGTGGGT	ATGGCAGCCA	GGTCAGACT	GGCTTCGGGC	TGGACACCAA	ACCCACCCGG	2100
	GCTGCGATG	TGCGGGCCAT	TAGCCAGGCC	CCCTACCTAG	GTGGGGTGGG	CTCAGCCGGC	2160
	ACCGCCCTGC	TGCACATCTA	TGACAAAGTG	ATGACGTC	AGAGGGGTG	CCGGCCTGGT	2220
55	GTCCCCAAAG	CTGTGGTGGT	GTCACAGGC	GGGAGGAGGCG	CAGAGGATG	AGCCGTTCC	2280
	GCCCAGAAC	TGAGGAACAA	TGGCATCTC	GTCTGGTCG	TGGGCGTGGG	GCCTGCTTA	2340
	AGTGAGGTC	TGCGGAGGCT	TGCAAGTC	CGGGATTCCC	TGATCCACGT	GGCAGCTTAC	2400
	GCCGACCTGC	GGTACCAACCA	GGACGTGCTC	ATTGAGTGGC	TGTTGAGGAGA	AGCCAAGCAG	2460
	CCAGTCAC	TCTGAAAC	CAGCCCGTGC	ATGAATGAGG	GCAGCTGCGT	CCTGCAGAAT	2520
60	GGGAGCTACC	GCTGCAAGTG	TCGGGATGGC	TGGGAGGGCC	CCCACTGCGA	GAACCGTGAG	2580
	TGGAGCTCTT	GCTCTGTATG	TGTGAGCCAG	GGATGGATT	TTGAGACGCC	CCTGAGGCCAC	2640
	ATGGCTCCCG	TGCAGGAGGG	CAGCAGCCG	ACCCCTCCCA	GCAACTACAG	AGAAGGCC	2700
	GGCACTGAAA	TGGTGCCTAC	CTTCTGGAAT	GTCTGTGCC	CAGGTCTTAA	GAATGTCTG	2760
	TTCCCGCCGT	GGCCAGGACC	ACTATTCTA	CTGAGGGAGG	AGGATGTCCC	AACTGCAGCC	2820

ATGCTGCTTA GAGACAAGAA AGCAGCTGAT GTCACCCACA AACGATGTTG TTGAAAAGTT 2880  
 TTGATGTGTA AGTAAATACC CACTTCTGT ACCTGCTGTG CCTTGGTGTAG GCTATGTCAT 2940  
 CTGCCACCTT TCCCTTGAGG ATAAACAAGG GGTCCTGAAG ACTTAAATTG AGCGGCCTGA 3000  
 5 CGTCCCTTGT CACACAATCA ATGCTGCCA GAATGTTGTT GACACAGTAA TGCCCAGCAG 3060  
 AGGCCCTTAC TAGAGCATCC TTTGGACGGC GAAGGCCACG GCCTTCAAG ATGAAAGCA 3120  
 GCAGCTTTC CACTTCCCCA GAGACATTCT GGATGCATT GCATTGAGTC TGAAAGGGGG 3180  
 CTTGAGGGAC GTTGTGACT TCTTGGCGAC TGCCCTTGT GTGTGGAAGA GACTTGGAAA 3240  
 GGTCTCAGAC TGAATGTGAC CAATTAACCA GCTTGGTTGA TGATGGGGGA GGGGCTGAGT 3300  
 10 TGTGCATGGG CCCAGGCTCG GAGGGCCACG TAAAATCGTT CTGAGTCGTG AGCAGTGTCC 3360  
 ACCTTGAAGG TCTTC

**CBF9 Protein sequence**

Gene name: ESTs  
 Unigene number: Hs.157601

Protein Accession #: none found

15 Signal sequence: 1-17  
 Transmembrane domains: none found  
 VGW domains: 49-223; 341-518; 529-706  
 EGF domains: 298-333; 715-748  
 20 Cellular Localization: plasma membrane

1	11	21	31	41	51	
25 MPPFLLLEAV	CVFLFSRVPP	SLPLQEVHVS	KETIGKISAA	SKMMWCSAAV	DIMFLLDGSN	60
SVGKGSFERS	KHFAITVCVG	LDISPERVRV	GAFQFSSTPH	LEFFPLDSFST	QQEVKARIKR	120
MVFKGGRRET	ELALKYLLHR	GLPGGRNHSV	PQILIIVTDG	KSQGDVALPS	KQLKERGVTV	180
FAVGVRFPWR	EELHALASEP	RGQHVLLAEQ	VEDAINGLFS	TLSSSAICSS	ATPDCRVEAH	240
PCEHRTLEMV	REFAGNAPCW	RGSRRRTLAVL	AAHCDFYSWK	RVLTHPATC	YRTTCPGPCD	300
30 SQPCQNGGT	VPEGLDGYQC	LCPLAFIGGEA	NCALKSLEC	RVDLLFLLDS	SAGTTLDGFL	360
RAKVFVKRFV	RAVLSEDSRA	RVGVATVYSRE	LLVAVPVGEY	QDVFDLVWSL	DGIPFRGGPT	420
LTGSALRQAA	ERGFGSATRT	GQDRPRRVVV	LLTESHSEDE	VAGPARHARA	RELLLLGVGS	480
EAVRAELEI	TGSPKHVMVY	SDPQDLFNQI	PELQKLCQR	QRPGCRTQAL	DLVFMLDTSA	540
35 SVGPNENFAQM	QSFRVSCALQ	FEVNPDVTQV	GLVVYGSVQ	TAFGLDTKPT	RAAMLRAISQ	600
APYLGGVGSA	GTALLHIYDK	VMTVQRGARP	GVPKAVVVL	GGRGAEDAAB	PAQKLRNNGI	660
SVLVVGVPV	LSEGRLRLAG	PRDSLHVA	YADLRYHQDV	LIEWLGEAK	QPVNLCKPSP	720
CMNEGSCVLO	NGSYRCKCRD	GWEGPHCENR	EWSSCSVCVS	QGWILETPLR	HMAPVQEGSS	780
RTPPSNYREG	LGTEMVPTFW	NVCAPGP				

40

**BFO8 DNA SEQUENCE**

Gene name: TMPRSS3a  
 Unigene number: Hs.298241  
 45 Probeset Accession #: AI538613  
 Nucleic Acid Accession #: AB038157  
 Coding sequence: 202-1566 (underlined sequences correspond to start and stop codons)

50

1	11	21	31	41	51	
55 ACCGGGCACC	GGACGGCTCG	GGTACTTTCG	TTCTTAATT	GGTCATGCC	GTGTGAGCCA	60
GGAAAGGGCT	GTGTTTATGG	GAAGCCAGTA	ACACTGTGGC	CTACTATCTC	TTCCGTGGTG	120
CCATCTACAT	TTTGGGACT	CGGAATTAT	GAGGTAGAGG	TGGAGGCCA	GCCGGATGTC	180
AGAGGTCTG	AAATAGTCAC	<u>CATGGGGAA</u>	AATGATCCGC	CTGCTGTTGA	AGCCCCCTTC	240
TCATTCCGAT	CGCTTTTGG	CCTTGATGAT	TTGAAAATAA	GTCCTGTTGC	ACCAGATGCA	300
60 GATGCTGTTG	CTGCACAGAT	CCTGTCACTG	CTGCCATTGA	AGTTTTTCC	AATCATCGTC	360
ATTGGGATCA	TTGCATTGAT	ATTAGCACTG	GCCATTGGTC	TGGGCATCCA	CTTCGACTGC	420
TCAGGGAAGT	ACAGATGTG	CTCATCC	AAGTGTATCG	AGCTGATAGC	TCGATGTGAC	480
GGAGTCTCGG	ATTGCAAAGA	CGGGGAGGAC	GAGTACCGCT	GTGTCCGGGT	GGGTGGTCAG	540
AATGCCGTG	TCCAGGTGTT	CACAGCTGCT	TCGTGGAAGA	CCATGTGCTC	CGATGACTGG	600
65 AAGGGTCACT	ACGCAAATGT	TGCGCTGTG	CAACTGGGTT	TCCCAAGCTA	TGTGAGTTCA	660
GATAACCTCA	GAGTGAGCTC	GCTGGAGGGG	CAGTCCGGG	AGGAGTTG	GTCCCATCGAT	720
CACCTCTTGC	CAGATGACAA	GGTGA	TTACACCA	CAGTATATGT	GAGGGAGGGA	780

	TGTGCCTCTG	GCCACGTGGT	TACCTTGCAG	TGCACAGCCT	GTGGTCATAG	AAGGGGCTAC	840
	AGCTCACGCA	TCGTGGGTGG	AAACATGTCC	TTGCTCTCGC	AGTGGCCCTG	GCAGGCCAGC	900
5	CTTCAGTTCC	AGGGCTACCA	CCTGTGCGGG	GGCTCTGTCA	TCACGCCCT	GTGGATCATC	960
	ACTGCTGCAC	ACTGTGTTA	TGACTTGTAC	CTCCCCAAGT	CATGGACCAT	CCAGGTGGGT	1020
	CTAGTTCCC	TGTTGGACAA	TCCAGCCCCA	TCCCCACTTGG	TGGAGAAGAT	TGTCTACCAC	1080
	AGCAAGTACA	AGCCAAGAG	GCTGGGCAAT	GACATGCC	TTATGAAGCT	GGCCGGGCCA	1140
	CTCACGTTCA	ATGAAATGAT	CCAGCCTGTG	TGCCTGCCA	ACTCTGAAGA	GAACCTCCCC	1200
	GATGGAAAAG	TGTGCTGGAC	GTCAGGATGG	GGGGCCACAG	AGGATGGAGC	AGGTGACGCC	1260
10	TCCCCGTCTC	TGAACCAACGC	GGCCGTCCCT	TTGATTTCGA	ACAAGATCTG	CAACCCACAG	1320
	GACGTGTAGC	GTGGCATCAT	CTCCCCCTCC	ATGCTCTGCG	CGGGTACCT	GACGGGTGGC	1380
	GTGGACAGCT	GCCAGGGGGG	CAGCGGGGGG	CCCCCTGGTGT	GTCAAGAGAG	GAGGCTGTGG	1440
	AAGTTAGTGG	GAGCGACAG	CTTTGGCATC	GGCTGCGCAG	AGGTGAACAA	GCCTGGGTG	1500
	TACACCCGTG	TCACCTCCTT	CCTGGACTGG	ATCCACGAGC	AGATGGAGAG	AGACCTAAA	1560
15	ACCTGAAGAG	GAAGGGGACA	AGTAGGCCACC	TGAGTTCCCTG	AGGTGATGAA	GACAGCCGA	1620
	TCCTCCCCCTG	GACTCCCGTG	TAGGAACCTG	CACACGAGCA	GACACCCTG	GAGCTCTGAG	1680
	TTCCGGCACC	AGTAGCAGGC	CCGAAAGAGG	CACCCCTCCA	TCTGATTCCA	GCACAACCTT	1740
	CAAGCTGCTT	TTTGTTTTTT	GTTTTTTTGA	GGTGGAGTCT	CGCTCTGTTG	CCCAGGCTGG	1800
	AGTGCAGTGG	CGAAATCCCT	GCTCACTGCA	GCCTCCGCTT	CCCTGGTTCA	AGCGATTCTC	1860
20	TTGCCTCAGC	TTCCCCAGTA	GCTGGGACCA	CAGGTGCCCG	CCACACACCC	CAACTAATT	1920
	TTGTATTTTTT	AGTAGAGACA	GGGTTTCACC	ATGTTGGCCA	GGCTGCTCTC	AAACCCCTGA	1980
	CCTCAAATGA	TGTGCCCTGCT	TCAGCCTCCC	ACAGTGCCTGG	GATTACAGGC	ATGGGCCACC	2040
	ACGCCTAGCC	TCACGCTCTT	TTCTGATCTT	CACTAAGAAC	AAAAGAAGCA	GCAACTTGCA	2100
	AGGGCGGCC	TTCCCACTGG	TCCATCTGGT	TTTCTCTCCA	GGGGTCTTGC	AAAATTCTG	2160
25	ACGAGATAAG	CAGTTATGTG	ACCTCACGTG	CAAAGCCACC	AACAGCCACT	CAGAAAAGAC	2220
	GCACCAGCCC	AGAAGTGCAG	AACTGCAGTC	ACTGCACGTT	TTCATCTCTA	GGGACCAGAA	2280
	CCAAACCCAC	CCTTCTACT	TCCAAGACTT	ATTTTCACAT	GTGGGGAGGT	TAATCTAGGA	2340
	ATGACTCGTT	TAAGGCCTAT	TTTCATGATT	TCTTTGTAGC	ATTTGGTGCT	TGACGTATTA	2400
	TTGTCCTTTG	ATTCCAAATA	ATATGTTTCC	TTCCCTCAAA	AAAAAAAAAA	AAAAAAAAAA	2460
30	AAAAAAA						

**BFO8 Protein sequence:**

35	Gene name:	TMPRSS3a					
	Unigene number:	Hs.298241					
	Probeset Accession #:	AI538613					
	Protein Accession #:	BAB20077					
40	Signal sequence:	none found					
	Transmembrane domains:	43-65, 239-261					
	Tryp_SPC domain:	216-444					
	Cellular Localization:	not determined					
45	1	11	21	31	41	51	
	MGENDPPEAVE	APFSFRSLFG	LDDLKISPVA	PDADAVAAQI	LSLLPLKFFP	IIVIGIIIALI	60
	LALAIGLGIH	FDCSGKYRCR	SSFKIELIA	RCDGVSDCKD	GEDEYRCVRV	GGQNAVLQVF	120
50	TAASWKTMC	DDWKGHYANV	ACAQLGFPSY	VSSDNLRVSS	LEGQFREEFV	SIDHLLPDDK	180
	VITALHHHSVY	REGCASGHVV	TLOCTACGHR	RGYSSRIVGG	NMSILLSQWPW	QASLQFQGYH	240
	LCGGSVITPL	WIITAACCVY	DLYLPKSWTI	QVGLVSLLDN	PAPSHLVEKI	VYHSKYKPKR	300
	LGNDIALMKL	AGPLTFNEMI	QPVCPLPNSEE	NFPDGKVCWT	SGWGATEDGA	GDASPVLNHA	360
	AVPLISNKIC	NHRDVYGGII	SPSMILCAGYL	TGGVDSCQGD	SGGPLVCQER	RLWKLVVGATS	420
55	FGIGCAEVNK	PGVYTRVTSF	LDWIHEQMER	DLKT			

WHAT IS CLAIMED IS:

- 1           1. A method of screening drug candidates comprising:
  - 2               a) providing a cell that expresses an expression profile gene selected from the group consisting of an expression profile gene set forth in Table 1 or Table 2 or fragment thereof;
  - 5               b) adding a drug candidate to said cell; and
  - 6               c) determining the effect of said drug candidate on the expression of said expression profile gene.
- 1           2. A method according to claim 1 wherein said determining comprises comparing the level of expression in the absence of said drug candidate to the level of expression in the presence of said drug candidate.
- 1           3. A method of screening for a bioactive agent capable of binding to a colorectal cancer modulator protein (colorectal cancer modulator protein), wherein said colorectal cancer modulator protein is encoded by a nucleic acid selected from the group consisting of a nucleic acid of Table 1 or Table 2 or a fragment thereof, said method comprising:
  - 6               a) combining said colorectal cancer modulator protein and a candidate bioactive agent; and
  - 8               b) determining the binding of said candidate agent to said colorectal cancer modulator protein.
- 1           4. A method for screening for a bioactive agent capable of modulating the activity of a colorectal cancer modulator protein, wherein said colorectal cancer modulator protein is encoded by a nucleic acid selected from the group consisting of a nucleic acid of Table 1 or Table 2 or a fragment thereof, said method comprising:
  - 5               a) combining said colorectal cancer modulator protein and a candidate bioactive agent; and

7                   b) determining the effect of said candidate agent on the bioactivity of said  
8                   colorectal cancer modulator protein.

1                   5. A method of evaluating the effect of a candidate colorectal cancer drug  
2                   comprising:

3                   a) administering said drug to a patient;

4                   b) removing a cell sample from said patient; and

5                   c) determining the expression of a gene selected from the group consisting of a  
6                   nucleic acid of Table 1 or Table 2.

1                   6. A method according to claim 5 further comprising comparing said  
2                   expression profile to an expression profile of a healthy individual.

1                   7. A method of diagnosing colorectal cancer comprising:

2                   a) determining the expression of one or more genes selected from the group  
3                   consisting of a nucleic acid of Table 1 or Table 2 or a fragment thereof or a polypeptide  
4                   encoded thereby in a first tissue type of a first individual; and

5                   b) comparing said expression of said gene(s) from a second normal tissue type  
6                   from said first individual or a second unaffected individual;

7                   wherein a difference in said expression indicates that the first individual has  
8                   colorectal cancer.

1                   8. A method for screening for a bioactive agent capable of interfering with the  
2                   binding of a colorectal cancer modulator protein (colorectal cancer modulator protein) or a  
3                   fragment thereof and an antibody which binds to said colorectal cancer modulator protein or  
4                   fragment thereof, said method comprising:

5                   a) combining a colorectal cancer modulator protein or fragment thereof, a  
6                   candidate bioactive agent and an antibody which binds to said colorectal cancer modulator  
7                   protein or fragment thereof; and

8                   b) determining the binding of said colorectal cancer modulator protein or  
9                   fragment thereof and said antibody.

1               9. A method for inhibiting the activity of a colorectal cancer modulator  
2 protein (colorectal cancer modulator protein), wherein said colorectal cancer modulator  
3 protein is encoded by a nucleic acid selected from the group consisting of a nucleic acid of  
4 Table 1 or Table 2 or a fragment thereof, said method comprising binding an inhibitor to said  
5 colorectal cancer modulator protein.

1               10. A method according to claim 9 wherein said inhibitor is an antibody.

1               11. A method of treating colorectal cancer comprising administering to a  
2 patient an inhibitor of a colorectal cancer modulator protein, wherein said colorectal cancer  
3 modulator protein is encoded by a nucleic acid selected from the group consisting of a  
4 nucleic acid of Table 1 or Table 2 or a fragment thereof.

1               12. A method according to claim 11 wherein said inhibitor is an antibody.

1               13. A method of neutralizing the effect of a colorectal cancer modulator  
2 protein, or a fragment thereof, comprising contacting an agent specific for said protein with  
3 said protein in an amount sufficient to effect neutralization.

1               14. A method for localizing a therapeutic moiety to colorectal cancer tissue  
2 comprising exposing said tissue to an antibody to a colorectal cancer modulator protein or  
3 fragment thereof conjugated to said therapeutic moiety.

1               15. The method of Claim 14, wherein said therapeutic moiety is a cytotoxic  
2 agent.

1               16. The method of Claim 14, wherein said therapeutic moiety is a  
2 radioisotope.

1               17. A method for inhibiting colorectal cancer in a cell, wherein said method  
2 comprises administering to a cell a composition comprising antisense molecules to a nucleic  
3 acid of Table 1 or Table 2.

1               18. An antibody which specifically binds to a protein encoded by a nucleic  
2 acid of Table 1 or Table 2 or a fragment thereof.

1                   19. The antibody of Claim 18, wherein said antibody is a monoclonal  
2                   antibody.

1                   20. The antibody of Claim 18, wherein said antibody is a humanized  
2                   antibody.

1                   21. The antibody of Claim 18, wherein said antibody is an antibody fragment.

1                   22. A biochip comprising one or more nucleic acid segments selected from  
2                   the group consisting of a nucleic acid of Table 1 or Table 2 or a fragment thereof, wherein  
3                   said biochip comprises fewer than 1000 nucleic acid probes.

1                   23. A nucleic acid having a sequence at least 95% homologous to a sequence  
2                   of a nucleic acid of Table 1 or Table 2 or its complement.

1                   24. A nucleic acid which hybridizes under high stringency to a nucleic acid of  
2                   Table 1 or Table 2 or its complement.

1                   25. A polypeptide encoded by the nucleic acid of Claim 23 or 24.

1                   26. A method of eliciting an immune response in an individual, said method  
2                   comprising administering to said individual a composition comprising the polypeptide of  
3                   Claim 25 or a fragment thereof.

1                   27. A method of eliciting an immune response in an individual, said method  
2                   comprising administering to said individual a composition comprising a nucleic acid  
3                   comprising a sequence of a nucleic acid of Table 1 or Table 2 or a fragment thereof.

1                   28. A method of determining the prognosis of an individual with colorectal  
2                   cancer comprising:

3                   a) determining the expression of one or more genes selected from the group  
4                   consisting of a nucleic acid of Table 1 or Table 2 or a fragment thereof in a first tissue type of  
5                   a first individual; and

6                   b) comparing said expression of said gene(s) from a second normal tissue type  
7                   from said first individual or a second unaffected individual;

8                   wherein a substantial difference in said expression indicates a poor prognosis.

1                   29. A method of treating colorectal cancer comprising administering to an  
2 individual having colorectal cancer an antibody to a colorectal cancer modulator protein or  
3 fragment thereof conjugated to a therapeutic moiety.

1                   30. The method of Claim 29, wherein said therapeutic moiety is a cytotoxic  
2 agent.

1                   31. The method of Claim 29, wherein said therapeutic moiety is a  
2 radioisotope.